Drug Facts

Active ingredients

Purpose

Benzocaine 7%......Oral pain reliever Menthol 1%.....Oral pain reliever

Uses temporarily relieves occasional minor irritation and pain associated with

- sore throat sore mouth
- canker sores

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
 rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other 'caine' anesthetics.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

■ for teething ■ in children under 6 years of age

When using this product

do not exceed recommended dosage

Drug Facts (continued)

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and older	apply to affected area (one spray); gargle, swish around in the mouth, or allow to remain in place at least 1 minute then spit out. Use up to 4 times daily or as directe by a doctor.
children 6 to under 12 years of age	supervise use
children under 6 years of age	do not use

Drug Facts (continued)

Other information

■ store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, cetylpyridinium chloride, dibasic sodium phosphate, edetate disodium, flavor, hydrochloric acid, polyethylene glycol, propylene glycol, purified water, sodium hydroxide, sucralose

