

## Drug Facts

### Active ingredients (in each 20 mL) Purposes **MUCINEX FAST-MAX COLD & FLU**

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Guaifenesin 400 mg.....Expectorant  
Phenylephrine HCl 10 mg.....Nasal decongestant

## Drug Facts (continued)

### Uses **MUCINEX FAST-MAX COLD & FLU**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - stuffy nose
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### **MUCINEX NIGHTSHIFT SEVERE COLD & FLU**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - sneezing
  - sinus congestion and pressure
  - runny nose
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- diabetes
- liver disease
- high blood pressure
- thyroid disease
- glaucoma (**Nightshift Severe Cold & Flu only**)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (**Nightshift Severe Cold & Flu only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nightshift Severe Cold & Flu only**)

### When using this product do not use more than directed

- excitability may occur, especially in children (**Nightshift Severe Cold & Flu only**)

## Drug Facts (continued)

### Active ingredients (in each 20 mL) Purposes **MUCINEX NIGHTSHIFT SEVERE COLD & FLU**

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Phenylephrine HCl 10 mg.....Nasal decongestant  
Triprolidine HCl 2.5 mg.....Antihistamine

## Drug Facts (continued)

- marked drowsiness may occur (**Nightshift Severe Cold & Flu only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nightshift Severe Cold & Flu only**)
- avoid alcoholic drinks (**Nightshift Severe Cold & Flu only**)
- use caution when driving a motor vehicle or operating machinery (**Nightshift Severe Cold & Flu only**)

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

### **MUCINEX FAST-MAX COLD & FLU**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### **MUCINEX NIGHTSHIFT SEVERE COLD & FLU**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

## Other information

- each 20 mL contains: sodium 12 mg (Fast-Max Cold & Flu only) and sodium 16 mg (Nightshift Severe Cold & Flu only)
- store at 20-25°C (68-77°F)
- do not refrigerate

### Inactive ingredients (**MUCINEX FAST-MAX COLD & FLU**)

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate<sup>1</sup>, xanthan gum

<sup>1</sup>may contain this ingredient

## Drug Facts (continued)

### Inactive ingredients (**MUCINEX NIGHTSHIFT SEVERE COLD & FLU**)

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

**Do not take Mucinex Fast-Max Cold & Flu and Mucinex Nightshift Severe Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinex liquid. TAKE ONLY AS DIRECTED.**