

## Drug Facts

### Active ingredients (in each liquid gel) Purposes Mucinex FAST-MAX DAY Cold & Flu

Acetaminophen 325 mg...Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine HCl 5 mg.....Nasal decongestant

### Active ingredients (in each liquid gel) Purposes Mucinex FAST-MAX NIGHT Cold & Flu

Acetaminophen 325 mg...Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses** ■ temporarily relieves these common cold and flu symptoms:  
■ sinus congestion and pressure  
■ cough ■ minor aches and pains  
■ headache ■ nasal congestion ■ sore throat  
■ runny nose (*NIGHT only*) ■ sneezing (*NIGHT only*)  
■ itching of the nose or throat (*NIGHT only*)  
■ itchy, watery eyes due to hay fever (*NIGHT only*)  
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (*DAY only*)  
■ controls cough to help you get to sleep  
■ temporarily reduces fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- high blood pressure ■ thyroid disease

## Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- glaucoma (*NIGHT only*)
- a breathing problem such as emphysema or chronic bronchitis (*NIGHT only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*NIGHT only*)

### When using this product ■ do not use more than directed

- excitability may occur, especially in children (*NIGHT only*)
- marked drowsiness may occur (*NIGHT only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*NIGHT only*)

- avoid alcoholic drinks (*NIGHT only*)

- be careful when driving a motor vehicle or operating machinery (*NIGHT only*)

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

### Other information ■ store at 20-25°C (68-77°F)

- avoid excessive heat

**Inactive ingredients (DAY only)** FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

**Inactive ingredients (NIGHT only)** D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide