# **Drug Facts**

# Active ingredients (in each 20 mL)

# **Purposes**

Dextromethorphan HBr 20 mg... Cough suppressant Guaifenesin 400 mg..... Expectorant Phenylephrine HCl 10 mg....Nasal decongestant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

### **Warnings**

#### Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# **Drug Facts** (continued)

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

# When using this product do not use more than directed

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.



# **Drug Facts** (continued)

### **Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over:
  20 mL in dosing cup provided every
  4 hours
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum \*may contain this ingredient

### Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

