

# Vetgene

## FARM CLEANER & SANITISER

### PRODUCT DESCRIPTION

Use Vetgene Farm Cleaner & Sanitiser wherever there is a risk of infection. Recommended for use in all types of livestock housing (e.g lambing pens, calf pens, broiler houses), on associated equipment for cattle, pigs, poultry, that can harbour numerous harmful micro-organisms and in veterinary medical facilities to clean, disinfect and sterilize even the most critical medical devices and instruments.

Vetgene Farm Cleaner & Sanitiser is formulated to provide a rapid and effective kill of a wide range of microorganisms. Containing Glutaldehyde and twin-chain-quats, this product works effectively in dirty conditions where other types of sanitisers would quickly exhaust.

Correct PPE is imperative. Standard PPE will not suffice. Elbow-length **butyl rubber** or **nitrile** gloves and **splash-proof** goggles or safety glasses with **full face shields**. Keep an eyebath close at hand and ready to use. Do not use with misting trigger, atomiser or sprayer without an **air-supplying respirator** (approved for use with Glutaraldehyde). Always ensure adequate ventilation.

Vetgene Farm Cleaner & Sanitiser will not damage sensitive equipment, e.g. endoscopes, when used according to the usage directions, (as well as the equipment manufacturer's instructions). Not suitable for use on rubber or plasticized PVC.

- HOSPITAL-GRADE
- CLEANS/DISINFECTS/STERILIZES
- ANTI-BACTERICIDAL
- ANTI-FUNGICIDAL
- ANTI-VIRUCIDAL
- STABLE
- REDUCED ODOUR
- REDUCED IRRITANT
- REDUCED SKIN DISCOLOURATION
- GENTLE ON EQUIPMENT

### INGREDIENTS

- 10% Glutaraldehyde.
- 10% Quat.
- 1% Phosphoric Acid.
- Balance comprises non hazardous ingredients.

### AVAILABLE SIZES

Available in 1L, 5L, 20L & 200L.

### PRODUCT USES

- Cleans hard surfaces.
- Disinfects hard surfaces.
- Protects from a wide range of disease-causing microorganisms.

### DIRECTIONS FOR USE

Livestock housing:

- Remove all livestock, portable equipment, litter & manure.
- Rinse out any gross soils.
- Dilute 1:50L - 1:200L in potable water depending on soil.
- Apply to all surfaces.
- Leave to penetrate for at least 5 minutes.
- Rinse off with clean water.
- Allow the area to dry before reintroducing animals or bedding.

Medical devices and instruments:

- Fully immerse instruments in the solution.
- Flush, then fill needles and other hollow parts with the solution.
- Soak in covered containers to prevent contamination and/or evaporation.

As a pre-clean (Dilute 1:20):

- Soak until visible soil loosens.
- Agitate for 1 minute.
- Drain solution and rinse with clean water.
- Dry completely.

Sterilization (dilute 1:5):

- Soak instruments at 35°C for 8 hours.
- Rinse with 5 times the amount of STERILE water required to immerse the instrument.
- Dry completely.
- Store for future use.

High level disinfection (dilute 1:5):

- Soak instruments at 35°C for 5 minutes, or at 25°C for 20 minutes.
- Rinse with 5 times the amount of STERILE water required to immerse the instrument.
- Dry completely.
- Store for future use.

### STORAGE

- Store in a cool dry place, out of direct sunlight.
- Store away from foodstuffs.
- Store away from heat and flames.

Hospital-grade disinfectant. bactericidal, fungicidal & virucidal, it protects against a wide range of microorganisms.



### SAFETY INSTRUCTIONS



**UN-2927: TOXIC LIQUID, CORROSIVE, ORGANIC, N.O.S.**



**DANGER**

Read label/SDS before use. Keep from children. Keep packaging in case medical advice is required.

- Toxic if swallowed.
- Causes severe skin burns & serious eye damage.
- Toxic to the environment.
- Combustible liquid.
- Harmful if inhaled.
- May cause an allergic reaction.
- May cause lung-damage.
- Keep away from flames/hot surfaces.
- Wash hands thoroughly after handling.
- Wear protective gloves & goggles/face shield.

### FIRST AID

- If on skin, wash with soap & water for at least 15 minutes.
- If in eyes, rinse well with water. Remove any contact lenses. Continue rinsing for 15 minutes.
- If inhaled, move to fresh air & rest where you can breathe comfortably.
- If ingested, do not induce vomiting. Drink a glass of water.

Seek immediate medical attention if you feel unwell or any irritation persists.

**POISON INFO CENTRES:**  
**NEW ZEALAND 0800 764 766**  
**AUSTRALIA 13 1126**

