

Use this product in: Bathrooms, Homes, Kitchens, Dental offices, Hospitals, Medical Offices, Nursing homes, Day care centers, Nurseries, Bars, Cafeterias, Institutional kitchens, USDA inspected food processing facilities, Colleges, Correctional facilities, Dressing rooms, Exercise facilities, Factories, Health clubs, Hotels, Institutional facilities, Institutions, Laundry rooms, Locker rooms, Motels, Prisons, Public facilities, Public rest rooms, Salons, Schools, Shower and bath areas, Sick rooms, Airplanes, Airports, Ambulances, Buses, Emergency vehicles, Ships, Taxis, Trailers, Trains and Transport terminals.

Use this product on washable hard, nonporous surfaces of: Bathroom fixtures, Bathtubs, Cabinets, Cages, Chairs, Computer keyboards, Counters, Cupboards, Desks, Diaper changing tables, Diaper pails, Doorknobs, Equipment tables, Faucets, Faucet Handles, Floors, Finished floors, Garbage cans, Glass, Highchairs, Lab benches, Linoleum, Light switches, Laminate flooring, Refrigerators - exterior, Refrigerated storage and display equipment<sup>8</sup>, Sealed floors, Showers, Shopping cart handles, Shower walls, Sinks, Storage areas, Stoves<sup>8</sup>, Tables, Tanning Beds, Exterior Toilet bowl surfaces, Toilet seats, Toys<sup>3</sup>, Tub walls, Tubs, Tiles, glazed, Walls.

Other hard nonporous surfaces made of: ABS, Acrylic, Aluminum, Brass, Chrome, Finished woodwork, Formica®, Glass, Glazed ceramic, Glazed enameled surfaces, Glazed porcelain, LDPE, HDPE, Metal, Plastic, Polycarbonate, PVC, Sealed granite, Sealed marble, Sealed quartz, Sealed stone, Stainless steel, Upholstery- vinyl and plastic.

Use this product to clean and disinfect nonporous personal protection equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, silicone rubber or PVC hearing protectors and vinyl covered earmuffs. Rinse all equipment that comes in contact with skin before reuse with clean warm water, and allow to air dry.

This product kills: **Bacteria:** Pseudomonas aeruginosa (Pseudomonas), Salmonella enterica (Salmonella),

Staphylococcus aureus (Staph), Acinetobacter baumannii, Burkholderia cepacia, Campylobacter jejuni (Campylobacter), Corynebacterium ammoniagenes, Enterobacter aerogenes, Enterococcus faecalis (Enterococcus), Enterococcus faecalis - Vancomycin Resistant (VRE), Escherichia coli (E. coli), Escherichia coli O157:H7, ESBL Escherichia coli - (Extended spectrum beta-lactamase producing E. coli), Klebsiella pneumoniae (Klebsiella), Klebsiella pneumoniae (NDM-1), Legionella pneumophila, Listeria monocytogenes, Salmonella schottmuelleri (Salmonella), Salmonella typhi (Salmonella), Serratia marcescens (Serratia), Shigella dysenteriae (Shigella), Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA)(NRS384)(USA300), Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA) (NRS123)(USA400) Staphylococcus aureus - Methicillin-Resistant (MRSA), Staphylococcus aureus - Multi-Drug Resistant (resistant to tetracycline [Tc], penicillin [Pc], streptomycin [Sm] and erythromycin [Em] and susceptible to chloramphenicol [Cm] *in vitro*), Staphylococcus aureus-Vancomycin Intermediate Resistant - (VISA), Streptococcus pyogenes (Strep), Vibrio cholerae.

**Viruses:** \*Hepatitis B Virus (HBV), \*Hepatitis C Virus (HCV), \*Herpes Simplex Virus Type 1, \*Herpes Simplex Virus Type 2, \*HIV-1 (AIDS Virus), \*Human Coronavirus, \*Influenza Type A Virus / Brazil Influenza, \*Influenza A Virus H1N1, \*Norwalk Virus (Norovirus), \*Respiratory Syncytial Virus (RSV), \*Rotavirus, \*SARS Associated Coronavirus (SARS) (cause of Severe Acute Respiratory Syndrome), \*SARS-Related Coronavirus 2 (SARS-CoV-2) (cause of COVID-19), \*Vaccinia Virus.

**Animal Viruses:** Avian Influenza Virus H3N2, Avian Influenza Virus H5N1, Canine Distemper Virus, Feline Calicivirus, Newcastle's Disease Virus, and Pseudorabies Virus.

**To Open Package:**

Remove cover. Pull up wipe corner from center of roll. Thread through hole in the cover. Replace cover. Open center cap on cover to remove wipe. When not in use, close center cap to prevent moisture loss.



# SPec 4

Clean and Disinfect in Just 4 Minutes

- FAST DRYING; LEAVES NO DULL OR STICKY RESIDUE
- BLEACH FREE • DEODORIZES
- KILLS 99.9% OF BACTERIA AND VIRUSES\*
- KILLS SARS-CoV-24, WHICH CAUSES COVID-19

**Active Ingredients:**

|   |                        |
|---|------------------------|
| Octyl decyl dimethyl ammonium chloride.....                                 | 0.0909%                |
| Dioctyl dimethyl ammonium chloride.....                                     | 0.0364%                |
| Didecyl dimethyl ammonium chloride.....                                     | 0.0545%                |
| Alkyl (C14, 50%; C12, 40%; C16, 10%) dimethyl benzyl ammonium chloride..... | 0.1212%                |
| <b>Other Ingredients:</b> .....   | 99.6970%               |
| <b>Total</b> .....  | 100.0000% <sup>1</sup> |

<sup>1</sup>Includes weight of absorbent wipe.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
 See side panel for Precautionary Statements and First Aid.

EPA Reg. No. 6836-340-68562 EPA Est. No. 10088-WI-2

**Net Contents: 1 lb. 9 ozs.**  
**180 Wipes 6" x 7" (15.24cm x 17.78cm)**

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use as a diaper wipe or for personal cleansing. This is not a baby wipe!  
 Do not flush in toilet.  
 For plastic and painted surfaces, spot test on an inconspicuous area before use.  
 This product will not harm most surfaces: including acrylic, sealed fiberglass and vinyl.

**TO CLEAN AND DISINFECT/ VIRUCIDAL\* DIRECTIONS:** Wipe hard, nonporous surface with wipe until surface is visibly wet. Allow surface to remain wet for 4 minutes. **For Norwalk Virus, Feline Calicivirus, Norovirus and Rotavirus,** treated surfaces must remain wet for 10 minutes. Let air dry. If surfaces are visibly dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush.

**Mold and mildew stain control,** this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Wipe surface and let it remain wet for 4 minutes. Repeat treatment every seven days, or more often if new growth appears.

This product is not for use on medical device surfaces.

**\*KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.**

**Personal protection:** Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

**Cleaning procedure:** Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of this product. This product can be used for this purpose.

**Contact time:** Leave surface wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses and bacteria claimed.

**Disposal of infectious material:** Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

**SANITIZING, DEODORIZING:** To sanitize, deodorize hard nonporous, non-food contact surfaces; wipe surface, using enough wipes to allow treated surface to remain wet for 15 seconds for Staphylococcus aureus (Staph) (ATCC 6538), Klebsiella pneumoniae (ATCC 4352), and Campylobacter jejuni (ATCC 29428). Let surface dry. For visibly soiled surfaces, clean excess dirt first.

**STORAGE AND DISPOSAL**

Store in original container in areas inaccessible to children. Keep securely closed. Nonrefillable container. Do not reuse or refill this container. Discard empty container in trash or recycle if available.

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

<sup>3</sup>Rinse toys with clean, potable water after disinfecting.

<sup>4</sup>On hard, non-porous, non-food surfaces

<sup>8</sup>Allow the surface to adjust to room temperature before disinfection

Formica® is a registered trademark of the Diller Corporation 1567.080320