

Cleans, Disinfects, Deodorizes Germicidal, Tuberculocidal, Fungicidal, and Virucidal\*

Active Ingredient:	Citric Acid	0.60%
Inert Ingredients:		. 99,40%
Total		100.00%

## KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS. CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating. drinking. chewing gum. using tobacco or using the toilet. FIRST AID: IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses. if present, after the first five minutes. then continue rinsing. Call a Poison Control Center or doctor for further treatment advice.

In the presence of 5% blood serum and a contact time of S minutes, is effective against: BACTERICIDAL: Pseudomonas aeruginosa, Slaphylococcus aureus. Salmonella enterica (formerly Salmonella choleraesuis), Carbapenem-resistani Klebsiella pneumoniae, Lisleria monocytogenes, Escherichia Coli 0157:H7, Staphylococcus aureus - MRSA. Staphylococcus epidermidis - MRSE, Enterococcus faecalis Vancomycin Resistant (VRE) VIRUCtDAL\*: Respiratory Syncytial Virus, Rotavirus, Human Immunodeficiency Virus Type 1, Herpes Simplex Type 1 Virus, Herpes Simplex Type 2 Virus, Vaccinia Virus, Influenza A Virus. Pandemic 2009 H1N1 Influenza A Virus (formerly called swine flu), Adenovirus Type 2, Feline Calicivirus (surrogate for Norovirus) FUNGICIDAL: Trichophyton interdigitale TUBERCULOCtDAL: Effective against Mycobacterium bovis In 5 minutes at 20°c / 68'F on previously cleaned hard non-porous inanimate surfaces.

AREAS OF USE: Hospitals. Health care facilities, Medical offices, Dental offices, Day care centers, Nursing homes, Schools. Bathrooms, Nurseries, Institutional facilities. TYPES OF **SURFACES:** Suitable for use on Floors, Walls, Fixtures, Countertops, Kitchen surfaces, Medical equipment surfaces, Stainless steel, Chrome, Glass, Vinyl, CuVerro® antimicrobial copper, Glazed ceramic tile, Plaslic, Appliances, Children's toys, Changing tables, High chairs.

EPA Reg. No. 34810-36 EPA Est. No. 34810-MO-1 Net Contents: 6 lbs. 7 oz. 400 Wet Wipes (7.1" x 7.1") DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with Its labeling. Product Is ready to use. Preclean areas with gross filth and visible soil prior to disin, fection. DISPENSER DIRECTIONS: Open lid. Pull out wipe from center of roll and Ihrough lid. Close lid to retain moisture. TO CLEAN: Unfold premoislened cloth and wipe area to be cleaned for 30 seconds or until clean. TO DISINFECT: Unfold premoislened cloth and wipe hard non-porous area or item 10 be di:'tiinfcctcd. Wet all ::;urfooc3 thoroughly. Leovc for 5 minute::;. Wipe or let ::iir dry. No rin::jing or wiping is required, except on direct food contact surfaces which require a potable water rinse after treatment. FUNGICIDAL DIRECTIONS: This product is effective against Trichophylon interdigitale (athlele's fool fungus and a cause of ringworm) on inanimate surfaces in locker rooms, dressing rooms, exercise facilities, and shower and bath areas. Follow disinfection directions.

DECONTAMINATION OF CRJTICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERIL-IZATION/HIGH LEVEL DISINFECTION: This product is not lo be used as a terminal sterilan/high level disinfectant on any surface or instrument that (1) is introduced directly into lhe human body, either into or ln contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes bul which does not ordinarily penetrate the blood barrier or otherwise enler normally slerile areas of the body. This product may be used to preclean or decontaminale critical or semi-critical medical devices prior to slerilization or high-level disinfection.

CLEANCIDE WIPES KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS

PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS In all settings in which !here is an expected likelihood of soiling of hard, non-porous surfaceslobjects with blood or body fluids and in which the surfaceslobjects likely to be soiled with blood or body fluids can be associated with lhe potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS virus).

\*\*SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS\*\*

\*\*SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HfV-1 (AIDS virus). PERSONAL PROTECTION: When handling objects soiled with blood or body fluids use disposable latex gloves, gowns. mask and eye coverings. CLEANING PROCEDURE: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant. CONTACT TIME: Leave precleaned surfaces wet for five minutes using mop. cloth or sponge to mitigate HIV-1 virus. Use a 5 minute contact lime to kill all bacteria, viruses\* and fungi listed on lhe label, DISPOSAL OF INFECTIOUS MATERIAL: Blood and other fluids should be autoclaved and disposed of according to Federal, State and Local regulations for infectious waste disposal.

**STORAGE AND DISPOSAL:** Do no! leave containers open during storage. No special handling is required. Towelette: Do not reuse towelette. Dispose of used towelette in trash. **Dispenser:** Nonrefillable container. Do not reuse or refill this container. Offer for recyclin9 if available.

Distributed by: Metropolitan Compounds 3343 N University Drive Hollywood, FL 33024 855-474-6781