

 **impello**<sup>®</sup>  
The Biostimulant Company™



**NERO**<sup>™</sup> **SC**

**For use on specialty crops, fruits, and vegetables**

**NET VOLUME** 1 gal (3.79 L)  
**NET WT.** 8.54 lbs/gallon (3.87kg)

Nero™ SC contains a selected duo of microbial strains that allows for better crop nutrition by fixing nitrogen (*Azospirillum brasilense*) and improving nutrient uptake by crops.

Nero™ SC is a soluble liquid and can be mixed with irrigation water or added to fertilization tanks and applied by drip irrigation.

## STORAGE AND DISPOSAL

Nero™ SC is stable for at least 12 months when stored in a cool, dry location in a tightly sealed container. Exposure to high heat and humidity are not recommended.

## SAFETY AND HANDLING

In case of contact with skin or eyes, promptly rinse with water at least 15 minutes. Contains microorganisms. May cause sensitization reactions. For detailed handling information, refer to [impellobio.com](http://impellobio.com) or contact [info@impellobio.com](mailto:info@impellobio.com).

## INCOMPATIBILITIES

Nero™ SC is not recommended to mix with strong alkaline or acidic products. For mixtures, please consult with the manufacturer or the supplier.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

### GUARANTEED BY:

**Agronova Biotech USA, Corp.**  
5323 Millenia Lakes Blvd. Suite 300  
Orlando, Florida 32839

### DISTRIBUTED BY:

**Impello Biosciences, LLC.**  
900 Josephine Court, Unit 2  
Loveland, CO 80537  
[info@impellobio.com](mailto:info@impellobio.com) • [www.impellobio.com](http://www.impellobio.com)

FL License # F003060

## CONTAINS NON-PLANT FOOD INGREDIENT(S) GUARANTEED ANALYSIS - SOIL AMENDING INGREDIENTS

*Azospirillum brasilense*.....1x10<sup>9</sup> CFU/mL  
*Pantoea dispersa*.....1x10<sup>8</sup> CFU/mL  
CFU/mL = Colony Forming Units per milliliter

## DOSAGE RECOMMENDATIONS

Nero™ SC is usually applied through irrigation systems. It must be applied to the root zone of the plant to maximize microbial activity in the rhizosphere. Do not exceed the recommended rates.

CROP	Max. no. of applications	Rate of application	
		fl.oz/acre	L/ha
Vegetable Crops	4-6	34.0-68.0	2.5-5
Row Crops			
Nut Trees			
Extensive herbaceous crops	3-6	34.0-68.0	2.5-5
Fruit trees	5-8	34.0	2.5

*\*For other crops and/or more information, consult with Impello's technical support department.*

BATCH no.

MANUFACTURED DATE

EXPIRATION DATE

100% SATISFACTION GUARANTEED