

www.antibodiesinc.com orders@antibodiesinc.com 530-758-4400

Product Datasheet

Anti-TRIP8b (Exon 4) Antibody FL650 Conjugate



KO Validated

Overview

C-1-1#	
Catalog #	75-208-FL650
Conjugate	FL650 Ex: 655 nm, Em: 676 nm
lsotype	lgG1
Clone Number	N212/3
Size	200 μL
Concentration	0.5 mg/mL
Host Species	Mouse Monoclonal
Format	Purified by Protein A chromatography
Buffer	PBS with 0.09% azide
Applications	ICC, IHC
Species Reactivity	Human, Mouse, and Rat
Immunogen	Fusion protein amino acids 1-192 (exon 1a, exon 4 and constant region) of rat TRIP8b (accession number Q925N3) produced recombinantly in E. Coli
Molecular Weight	70 kDa
Cite this Antibody	Antibodies Inc Cat# 75-208-FL650, RRID: AB_2939775
Details	
Details Target Description	PEX5-related protein, Peroxisomal Biogenesis Factor 5 Like, TPR-containing Rab8b-interacting protein or TRIP8b is encoded by the gene Pex5l. It is a member of the peroxisomal targeting signal receptor family. TRIP8b is a cytoplasmic accessory subunit of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels. TRIP8b interacts with the carboxyl-terminal region of HCN channels and regulates their cell-surface expression to the dendrites in many types of neurons. TRIP8b plays a role in regulation of HCN channel cyclic nucleotide kinetics. TRIP8b is expressed in the brain. Diseases associated with PEX5L include Rhizomelic Chondrodysplasia Punctata, Type 5 and Rhizomelic Chondrodysplasia Punctata, Type 2.
	protein or TRIP8b is encoded by the gene Pex5l. It is a member of the peroxisomal targeting signal receptor family. TRIP8b is a cytoplasmic accessory subunit of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels. TRIP8b interacts with the carboxyl-terminal region of HCN channels and regulates their cell-surface expression to the dendrites in many types of neurons. TRIP8b plays a role in regulation of HCN channel cyclic nucleotide kinetics. TRIP8b is expressed in the brain. Diseases associated with PEX5L include Rhizomelic Chondrodysplasia Punctata, Type 5
Target Description	protein or TRIP8b is encoded by the gene Pex5I. It is a member of the peroxisomal targeting signal receptor family. TRIP8b is a cytoplasmic accessory subunit of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels. TRIP8b interacts with the carboxyl-terminal region of HCN channels and regulates their cell-surface expression to the dendrites in many types of neurons. TRIP8b plays a role in regulation of HCN channel cyclic nucleotide kinetics. TRIP8b is expressed in the brain. Diseases associated with PEX5L include Rhizomelic Chondrodysplasia Punctata, Type 5 and Rhizomelic Chondrodysplasia Punctata, Type 2.

Storage

Aliquot and store at \leq -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to removing the cap.

Our Guarantee

As an original manufacturer, we are dedicated to creating quality and reproducible antibodies that further your research. We provide personalized customer support from the scientists that made the antibody and offer a free replacement or 100% refund if we cannot resolve an issue. Order today and experience our 50+ year passion for science.

Note: For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, viewable on our website.