

www.antibodiesinc.com orders@antibodiesinc.com 530-758-4400

## **Product Datasheet**

## Anti-Mad3 Antibody FL650 Conjugate



## Overview

Catalog #	75-227-FL650
Conjugate	FL650 Ex: 655 nm, Em: 676 nm
Isotype	lgG2b
Clone Number	N129A/6
Size	200 μL
Concentration	0.5 mg/mL
Host Species	Mouse Monoclonal
Format	Purified by Protein A chromatography
Buffer	PBS with 0.09% azide
Applications	ICC
Species Reactivity	Human and Mouse
Immunogen	Fusion protein amino acids 116-206 (C-terminus) of human Mad3 (accession number Q9BW11) produced recombinantly in E. Coli
Molecular Weight	25 kDa
Cite this Antibody	Antibodies Inc Cat# 75-227-FL650, RRID: AB_2939831
Details	
Details Target Description	MAX Dimerization Protein 3 is encoded by the gene MXD3. MXD3 is a member of the Myc superfamily. MXD3 is a transcriptional repressor which forms a heterodimer with the cofactor MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'- CAC[GA]TG-3'. MXD3 is expressed in lung and blood. Diseases associated with MXD3 include Perianal Hematoma and Bartholin's Duct Cyst.
	superfamily. MXD3 is a transcriptional repressor which forms a heterodimer with the cofactor MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. MXD3 is expressed in lung and blood. Diseases associated with MXD3 include
Target Description	superfamily. MXD3 is a transcriptional repressor which forms a heterodimer with the cofactor MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. MXD3 is expressed in lung and blood. Diseases associated with MXD3 include Perianal Hematoma and Bartholin's Duct Cyst.
Target Description Specificity	superfamily. MXD3 is a transcriptional repressor which forms a heterodimer with the cofactor MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. MXD3 is expressed in lung and blood. Diseases associated with MXD3 include Perianal Hematoma and Bartholin's Duct Cyst. No cross-reactivity reported Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity

## **Our Guarantee**

As an original manufacturer, we are dedicated to creating quality and reproducible antibodies that further your research. We provide personalized customer support from the scientists that made the antibody and offer a free replacement or 100% refund if we cannot resolve an issue. Order today and experience our 50+ year passion for science.

Note: For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, viewable on our website.