

Product Datasheet

Anti-Slo1/BKAlpha Potassium Channel Antibody FL490 Conjugate



Overview

Catalog #	75-421-FL490
Conjugate	FL490 Ex: 491 nm, Em: 515 nm
Isotype	lgG1
Clone Number	L6/48
Size	200 μL
Concentration	0.5 mg/mL
Host Species	Mouse Monoclonal
Format	Purified by Protein A chromatography
Buffer	PBS with 0.09% azide
Applications	ICC, IHC
Species Reactivity	Mouse and Rat
Immunogen	Fusion protein amino acids 690-1196 (cytoplasmic C-terminus) of mouse Slo1 (accession number Q08460) produced recombinantly in E. Coli
Molecular Weight	110-130 kDa
Cite this Antibody	Antibodies Inc Cat# 75-421-FL490, RRID: AB_2940396
Details	
Details Target Description	Calcium-activated potassium channel subunit alpha-1, Potassium Calcium-Activated Channel Subfamily M Alpha 1 or Slo1/BKalpha is encoded by the gene KCNMA1. BKalpha binds to one of 4 different beta subunits to form a channel in the calcium activated large conductance (MaxiK channel) family. This potassium channel is activated by membrane depolarization or an increase in cytosolic ca2+ and mediated the export of K+. BKalpha is involved with various cell processes including contraction of smooth muscle, regulation of transmitter release, innate immunity and regulation of membrane potential. Diseases associated with KCNMA1 include forms of Epilepsy and Developmental Delay Cerebellar Atrophy and Seizures.
	Subfamily M Alpha 1 or Slo1/BKalpha is encoded by the gene KCNMA1. BKalpha binds to one of 4 different beta subunits to form a channel in the calcium activated large conductance (MaxiK channel) family. This potassium channel is activated by membrane depolarization or an increase in cytosolic ca2+ and mediated the export of K+. BKalpha is involved with various cell processes including contraction of smooth muscle, regulation of transmitter release, innate immunity and regulation of membrane potential. Diseases associated with KCNMA1 include forms of Epilepsy and
Target Description	Subfamily M Alpha 1 or Slo1/BKalpha is encoded by the gene KCNMA1. BKalpha binds to one of 4 different beta subunits to form a channel in the calcium activated large conductance (MaxiK channel) family. This potassium channel is activated by membrane depolarization or an increase in cytosolic ca2+ and mediated the export of K+. BKalpha is involved with various cell processes including contraction of smooth muscle, regulation of transmitter release, innate immunity and regulation of membrane potential. Diseases associated with KCNMA1 include forms of Epilepsy and Developmental Delay Cerebellar Atrophy and Seizures.

Storage

Aliquot and store at \leq -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to removing the cap.

Our Guarantee

As an original manufacturer, we are dedicated to creating quality and reproducible antibodies that further your research. We provide personalized customer support from the scientists that made the antibody and offer a free replacement or 100% refund if we cannot resolve an issue. Order today and experience our 50+ year passion for science.

Note: For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, viewable on our website.