

## Product Datasheet

## Anti-Glycine Receptor Alpha3L Antibody FL650 Conjugate



## Overview

<b>Catalog #</b>	75-427-FL650
<b>Conjugate</b>	FL650 Ex: 655 nm, Em: 676 nm
<b>Isotype</b>	IgG1
<b>Clone Number</b>	N424/48
<b>Size</b>	200 µL
<b>Concentration</b>	0.5 mg/mL
<b>Host Species</b>	Mouse Monoclonal
<b>Format</b>	Purified by Protein A chromatography
<b>Buffer</b>	PBS with 0.09% azide
<b>Applications</b>	ICC, IHC
<b>Species Reactivity</b>	Human
<b>Immunogen</b>	Fusion protein amino acids 342-431 (cytoplasmic loop) of human GlyRA3L (accession number O75311) produced recombinantly in E. Coli
<b>Molecular Weight</b>	55 kDa
<b>Cite this Antibody</b>	Antibodies Inc Cat# 75-427-FL650, RRID: AB_2940423

## Details

<b>Target Description</b>	Glycine is an important inhibitory transmitter in the brainstem and spinal cord. Glycine receptors are members of the ligand-gated ion channel family (LGICs) that mediate rapid chemical neurotransmission (Schofield et al., 2003). The binding of glycine to its receptor produces a large increase in chloride conductance, which causes membrane hyperpolarization. Glycine receptors are anchored at inhibitory chemical synapses by a cytoplasmic protein, gephyrin (Fischer et al., 2000). The glycine receptor has been used to great advantage in the identification of the binding sites for alcohol on the LGIC family of proteins (Beckstead et al., 2001; Mihic et al., 1997). These receptors have also been extremely useful in studies of synaptic clustering of receptors (Craig and Lichtman, 2001). The glycine receptor may also act in concert with an NMDAR subunit to form an excitatory receptor (Chatterton et al., 2002).
<b>Specificity</b>	Does not cross-react with GlyRA3K
<b>Purification Method</b>	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and conjugation of purified mAb. Purified mAbs are >90% specific antibody.

**Quality Control Tests**

Each new lot of antibody is quality control tested on cells overexpressing target protein and confirmed to give the expected staining pattern.

**Storage**

Aliquot and store at  $\leq -20^{\circ}\text{C}$  for long term storage. For short term storage, store at  $2-8^{\circ}\text{C}$ . For maximum recovery of product, centrifuge the vial prior to removing the cap.

**Our Guarantee**

---

As an original manufacturer, we are dedicated to creating quality and reproducible antibodies that further your research. We provide personalized customer support from the scientists that made the antibody and offer a free replacement or 100% refund if we cannot resolve an issue. Order today and experience our 50+ year passion for science.

**Note:** For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, viewable on our website.