

Anti-Tubulin, Beta-III Immunohistofluorescence Protocol

Catalog #: 2020-TUB

Species: mouse

Tissue: Mouse dentate gyrus

Fixation: 4% paraformaldehyde overnight

Antibody incubation: Primary Antibody: RT, 1 hour Secondary Antibody: RT, 1 hour

Antigen Retrieval: 10mM citrate buffer (pH 6.0, 0.05% Tween 20)

Materials Required

- ✓ **Fixative:** 4% paraformaldehyde in PBS
 - ✓ **1X PBS:** 137 mM NaCl, 28 mM Na₂HPO₄, 5.4 mM KCl, 2.9 mM KH₂PO₄, pH 7.6
 - ✓ **PBST:** 0.4% Triton-X in 1X PBS
 - ✓ **Blocking buffer:** 10% goat serum in PBST
 - ✓ **Incubation buffer:** 2% goat serum in PBST
 - ✓ **Secondary Antibody:** example used is Goat-Anti-Mouse Alexa Fluor 488, Molecular Probes ([catalog # A-11001](#))
 - ✓ **Mounting media:** Permount, Fisher Scientific ([catalog # SP15-100](#))
 - ✓ **Counterstain:** DAPI, ThermoFisher ([catalog # D1306](#))
-

Before you begin

This protocol was used for tissues fixed without perfusion following standard FFPE protocol. Cut tissue into 3-5mm sections and place in 4% paraformaldehyde in PBS overnight at 4C. For proper fixation, submerge sections into a 20x volume of fixative based on the mass of the tissue. After fixation of tissue, dehydrate and embed tissue into paraffin blocks according to standard protocol. Then section the blocks at 8 microns. Finally, transfer the sections onto positively charged slides (example: [SFH1103](#), BioCare Medical) and dry overnight at room temperature.

Deparaffinize

1. Warm slides for 10 minutes in a 60C oven.
2. Incubate slides in the following dehydrants in this order
 - I. Xylene: 3 times, 10 minute intervals
 - II. 100% ethanol: 2 times, 5 minute intervals
 - III. 95% ethanol: 2 times, 3 minute intervals
 - IV. 80% ethanol: 2 times, 1 minute interval
 - V. H₂O: 2 times, dip to rinse.

Antigen Retrieval

1. Place slides in 10mM citrate buffer (pH 6.0, room temperature) for 30 minutes.
2. Wash slides with 1X PBS.

Immunohistochemistry

1. Block slides with blocking buffer for 30 minutes at RT.
2. Wash slides with PBST 3 times, in 15 minute intervals.
3. Dilute Anti-Tubulin, Beta-III (Cat. # 2020-TUB) to 1:1000 in incubation buffer. Incubate sections for 1 hour at room temperature.

4. Wash slides with PBST 3 times, in 15 minute intervals.
5. Dilute secondary antibody in incubation buffer per manufacturer's recommendation. Incubate sections for 1 hour at room temperature.

Tech Tip:


- a. A goat anti-mouse Alexa Flour 488 antibody was used to produce the image below, ThermoFisher ([catalog # A-11001](#), 1:1000). Any anti-rabbit secondary can be used.
6. Remove secondary antibody and wash slides with PBST 3 times, in 15 minute intervals.
7. Prepare fresh DAPI solution per manufacturer's recommendation. Apply to slide and rinse 5 times with PBS.

Tech Tip:

- a. Any nuclear counterstain can be used, for this protocol DAPI was used, [catalog # D1306](#).
8. Apply mounting medium onto slide and gently place glass cover slip before viewing under the microscope.

Tech Tip:

- a. Any mounting media can be used, for this protocol Permount was used, [catalog # SP15-100](#).



Immunostaining of mouse dentate gyrus with anti-tubulin, beta-III (cat# 2020-TUB, red, 1:1000). The blue is DAPI staining nuclear DNA. The image was kindly provided by Robert Wine.