



**0.172% Dithiopyr  
Professional Fertilizer  
0-0-7**

Item # 02022272

**TURF AND ORNAMENTAL HERBICIDE**

For Season-Long Control of Crabgrass and Control or Suppression of Listed Annual Grasses and Broadleaf Weeds in Established Lawns and Ornamental Turf, including Golf Course Fairways, Roughs and Tee Boxes.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

**In the state of New York this product may be applied only by commercial applicators at no more than 288 lbs. (0.5 pounds AI) per acre (6.67 lbs. per 1,000 sq. ft.) per year. Not for use in Nassau or Suffolk Counties.**

**Active Ingredient:**

Dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate ..... 0.172%

**Inert Ingredients:**.....99.828%

TOTAL:..... 100.000%

EPA REG NO. 53883-209

EPA EST. NO. 9404-FL-1

Guaranteed Fertilizer Analysis: 0-0-7	
Total Nitrogen (N) .....	0.00%
0.00% Ammonical Nitrogen	
0.00% Water Soluble Nitrogen	
0.00% Water Insoluble Nitrogen	
Available Phosphate (P205).....	0.00%
Soluble Potash (K205).....	7.00%
Chlorine, not more than .....	7.00%
DERIVED FROM: ,Muriate of Potash.	

Net Weight: 45 lb bag

Distributed By:  
Sunniland Corporation  
P.O. Box 8001  
Sanford, FI 32772

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See [other] panel for further precautions and First Aid information

<b>FIRST AID</b>	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or physician immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or physician.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if inhaled. Causes moderate eye irritation. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, sock, shoes and chemical-resistant gloves such as natural rubber. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL INFORMATION**

Dithiopyr 0.172% Plus Fertilizer provides season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

**SPREADER SETTINGS** to apply at the listed pounds per 1,000 square feet

Spreader	Cone setting	1.66 lbs	2.45 lbs	3.33 lbs	4.89 lbs	6.67 lbs
Accu-Pro 2000	6	K ½	L 1/2	M1/2	N ½	P
SR2000	6	K ½	L 1/2	N	O ½	Q
Lesco 021820	0.25 Closed	C 1/2	D ½	E 1/2	F ½	H
Spyker 76-22		4.1	4.3	4.5	4.7	5.1
PSB Bigfoot	A.5	J	J ½	K1/2	L ½	M 1/2
Lely HR	II-B	4.75	5.25	6	6.5	7.5
Vicon 203		13	15	17	19	23

**FOR USE ON TURF**

Dithiopyr 0.172% Plus Fertilizer provides season-long control of crabgrass and control or suppression of other annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

Apply this product only to seeded, sodded, or sprigged turf that is well established. Before the use of Dithiopyr 0.172% Plus Fertilizer, roots must be developed and the turf mowed at least twice to a uniform height. Injury may result if the turf is not well established, or stressed by weather, pests or other conditions.

Dithiopyr 0.172% Plus Fertilizer may be applied to established turfgrass species listed on this label. When applied as directed, the following turfgrass species are tolerant to this product:

COOL-SEASON GRASSES	USE PRECAUTIONS
Creeping Bentgrass <sup>1</sup> ( <i>Agrostis palustris</i> ), Kentucky Bluegrass ( <i>Poa pratensis</i> ), Fine Fescue <sup>2</sup> ( <i>Festuca rubra</i> ), Tall Fescue ( <i>Festuca arundinacea</i> ), Perennial Ryegrass ( <i>Lolium perenne</i> ),	<sup>1</sup> Turfgrass injury may result from the use of this product on certain varieties of creeping bentgrass such as Cohansey, Carmen, Seaside and Washington. Not all varieties of creeping bentgrass have been testes. Do not apply this product to Colonial Bentgrass ( <i>Agrostis tenuis</i> ) varieties.  <sup>2</sup> Turfgrass injury may result from use of this product on the following varieties of Fescue such as Atlanta, Banner, Beauty, Bilgart, CF-2, Enjoy, HF-93, Highlight, Ivalo, Koket, Jamestown, Majenta, Mary, Pennilawn, Tamara, Tatiana, Walfort and Waldina. Not all varieties of fescue have been tested.
WARM-SEASON GRASSES	
Bahiagrass ( <i>Paspalum notatum</i> ), Bermudagrass ( <i>Cynodon dactylon</i> ), Buffalograss <sup>3</sup> ( <i>Buchloe dactyloides</i> ), Carpetgrass ( <i>Axonopus compressus</i> ), Centipedegrass ( <i>Eremochloa ophiuroides</i> ), Kikuyugrass ( <i>Pennisetum clandestinum</i> ), St. Augustinegrass ( <i>Stenotaphrum secundatum</i> ), Zousiagrass ( <i>Zoysia japonica</i> )	<sup>3</sup> Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

**NOTE:** It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response to turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

**RESEEDING, OVERSEEDING OR SPRIGGING**

Reseeding, overseeding, or sprigging of treated areas within 12 weeks after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use mechanical or power seeding equipment (slit seeders) for good seed-to-soil contact.

**USE RESTRICTIONS**

- Do not apply more than 6.67 lbs./1,000 sq. ft. (0.5 Lbs. Al/acre) per application.
- Do not apply more than 20 lbs./1,000 sq. ft. (1.5 lbs. Al/acre) per year.
- In New York, do not apply more than 6.67 lbs. /1,000 sq. ft. per year.
- Do not use clippings from treated turf for animal feed or for mulching around vegetables or fruit trees.
- Do not allow domestic animals to feed or forage on treated turf.
- Keep people and pets off treated areas until dust has settled.
- Do not use on golf course putting greens.

### **CRABGRASS CONTROL**

Dithiopyr 0.172% Plus Fertilizer provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. Dithiopyr 0.172% Plus Fertilizer provides “early post-emergence” control during the early stages of growth after crabgrass emerges from the ground. This early post-emergence control is limited and applications are only effective if applied before the crabgrass has its 3-leaf or first tier (which usually is about the time when you can first see the crabgrass in the lawn). With this post-emergence activity, the applicator has 2 to 4 weeks longer to make applications for control of crabgrass.

### **APPLICATION INSTRUCTIONS**

For best results, apply Dithiopyr 0.172% Plus Fertilizer within a few days after mowing and delay mowing again for a few days after application. Rainfall or irrigation activates the product. Improved weed control will result if the application is followed by ½ inch of rainfall or sprinkler irrigation. Weed control may be hampered if rainfall or irrigation does not occur with 30 days after application. Split, or half rate applications spaced 5 to 10 weeks apart may also enhance weed control. Cultural practices such as aerification and de-thatching should be completed and the turf allowed to recover before application of this product.

### **APPLICATION TIMING**

This product may be applied as a single application, split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described below.

**SPRING APPLICATIONS:** For single applications in the spring or early summer, follow the appropriate rate listed in the “Application Rates” table below. Rates are based on the location to be used (Region), the mowing height of the turfgrass, and whether the timing of the application is considered preemergence or postemergence. The length of residual weed control is dependent on the total rate applied, but may vary depending on weather, weed pressure and location.

Program 1: For preemergence control to turf mowed relatively high, such as home lawns, this program will provide 3 to 5 months of preemergence crabgrass control. Do not use this program for early postemergence control of crabgrass.

Program 2: For preemergence control at sites where the turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year. This program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass at locations where turf is cut high.

Program 3: For preemergence control at locations where turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year. This program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass at locations where the turf is cut relatively low.

**FALL APPLICATIONS:** Dithiopyr 0.172% Plus Fertilizer may be applied in late summer or early fall (late August through November). Follow the instructions for “Program 3” for crabgrass control through the early part of the next spring. Fall applications may be followed by an appropriate spring application to provide season-long control.*[Optional Language:*

**TURF GRASS REGIONS**



**TABLE 1: APPLICATION RATES FOR CRABGRASS CONTROL**

<b>REGION</b>	<b>PROGRAM 1</b> Preemergence (high-cut turf)	<b>PROGRAM 2</b> Preemergence (low-cut turf) Postemergence (high-cut turf)	<b>PROGRAM 3</b> Preemergence (low-cut turf) Postemergence (low-cut turf)
North (all areas)	1.66 lbs./1,000 sq. ft. (73 lbs./acre or 0.125 lbs. AI/acre)	2.45 lbs./1,000 sq. ft. (107 lbs./acre or 0.183 lbs. AI/acre)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)
Transition (DE, KS, KY, MD, MO, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY and RI.	2.45 lbs./1,000 sq. ft. (107 lbs./acre or 0.183 lbs. AI/acre)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)
South (AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN and TX)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)	6.67 lbs./1,000 sq. ft. (291 lbs./acre or 0.50 lbs. AI/acre)
Coastal South (HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, and TX)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)	6.67 lbs./1,000 sq. ft.* (291 lbs./acre or 0.50 lbs. AI/acre)	6.67 lbs./1,000 sq. ft.* (291 lbs./acre or 0.50 lbs. AI/acre)
* Split applications may be made 5-10 weeks apart.			

**SPLIT OR SEQUENTIAL APPLICATIONS**

Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with rates in the table above being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in

the spring following a fall application provided the maximum application rate per year is not exceeded. Refer to the "Use Restrictions" section above.

**CONTROL OF OTHER GRASS AND BROADLEAF WEEDS**

**SPRING APPLICATION:** When used as directed for crabgrass control in spring, Dithiopyr® 0.172% Plus Fertilizer will also control or suppress the following weeds when applied prior to their emergence:

<b>GRASSES</b>	
Common Name	Botanical Name
Barley	<i>Hordeum</i> spp.
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bluegrass (annual roughstalk)	<i>Poa annua</i>
Brome	<i>Bromus</i> spp.
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Crowfootgrass*	<i>Dactyloctenium aegyptium</i>
Dallisgrass (seedling)	<i>Paspalum dilatatum</i>
Foxtail (yellow & green)	<i>Setaria faberi, Setaria Verdi</i>
Goosegrass	<i>Eleusine indica</i>
Kikuyugrass*	<i>Pennisetum clandestinum</i>
Oats, wild	<i>Avena fatua</i>
Ryegrass (annual & perennial)	<i>Lolium</i> spp.
Sandbur	<i>Cenchrus</i> spp.
Smutgrass	<i>Sporobolus indicus</i>
<b>BROADLEAF WEEDS</b>	
Bittercress*	<i>Cardamine</i> spp.
Carpetweed	<i>Mullugo verticillata</i>
Chickweed*	<i>Stellaria</i> spp.
Geranium, Carolina*	<i>Geranium carolinianum</i>
Henbit	<i>Lamium</i> spp.
Knotweed, prostrate	<i>Polygonum aviculare</i>
Lespedeza, common*	<i>Lespedeza striata</i>
Marestail	<i>Conyza Canadensis</i>
Medic, black	<i>Medicago lupulina</i>
Mustard	<i>Brassica</i> spp.
Oxalis, buttercup	<i>Oxalis pes-caprae</i>
Parsley-piert*	<i>Alchemilla arvensis</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pineappleweed*	<i>Matricaria matricarioides</i>
Purslane, common	<i>Portulaca oleracea</i>
Rocket, London	<i>Sisymbrium irio</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Speedwell, corn*	<i>Veronica arvensis</i>
Spurge, garden	<i>Euphorbia hirta</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculate</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>
* Suppression only.	

**FALL APPLICATION:** When used as directed for late summer or early fall use, for crabgrass control through the early part of next spring, Dithiopyr 0.172% Plus Fertilizer will also control or suppress the following weeds when applied prior to their emergence:

<b>WEEDS CONTROLLED</b>	
Common Name	Botanical Name
Bluegrass (annual)	<i>Poa annua</i>
Bittercress	<i>Cardamine</i> spp.
Chickweed	<i>Stellaria</i> spp.
Geranium (Carolina)	<i>Geranium carolinianum</i>
Henbit	<i>Lamium</i> spp.
Parsley-piert	<i>Alchemilla arvensis</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>

**APPLICATION EQUIPMENT AND INSTRUCTIONS**

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer’s directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference.

Apply this product uniformly over the treatment area. More uniformity can be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Check equipment frequently to verify calibration and proper functioning. Clean and lubricate spreader after use.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**CONDITIONS OF SALE AND WARRANTY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying and using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Lawn and turf injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of

Sunniland Corporation. or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Sunniland Corporation warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

TO THE EXTENT PERMITTED BY LAW, Sunniland Corporation. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Sunniland Corporation. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

45 lb Bag