

SARS-CoV-2 Spike S1 protein

Receptor binding domain (RBD)

Cat. no. P2020-001


Product Information

Protein:	SARS-CoV-2-S1(RBD)-His (~ 27.5 kDa)
Sequence:	MRVQPTESIVRFPNITNLCPFGEVFNATRFASVYAWNRKRISNADVADYSVLYNSASFSTFKCYGVSPSTKLNLDLCTFNYYADSFVIRGDEVQRQIAPGQGTGKIADYNYKLPDDFTGCVIAWN SNNLDSKVGGNYYLYRFRKSNLKPFRDISTEIQAGSTPCNGVEGFNCYFPLQSYGF QPTNGVGYQPYRVVVLSEFELLHAPATVCGPKKSTNLVKNKCVNF
	Methionine at pos. 1 present due to cloning constraints, C-terminal His-tag not shown in sequence.
Source:	Recombinantly expressed in HEK293 cells.
Tag(s):	His-tag, C-terminal
Purification:	Purified by affinity chromatography and subsequent buffer exchange.
Formulation:	PBS; pH 7.4
	Liquid, stored and shipped at -80 °C.
Purity:	> 85 % (will be determined by densitometry of Coomassie stained gel, example next page)
Concentration:	Will be determined by BCA-Assay.
Long-term storage:	No recommendations.
Comment:	Protein migrates at higher molecular weight during SDS-PAGE due to posttranslational modifications.

Background Information:

The spike (S) glycoprotein of coronaviruses is essential for binding of the virus to the host cell at the beginning of the infection process. The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein is responsible for membrane fusion and is therefore required for virus entry and cell fusion. The target protein is also a major immunogen and a possible target for entry inhibitors.

The SARS-CoV-2 spike (S) protein is a large type I transmembrane protein composed of two subunits, S1 and S2. The S1 subunit contains a receptor-binding domain (RBD) responsible for binding to the host cell receptor angiotensin-converting enzyme 2 (ACE2). The S2 subunit mediates fusion between the viral and host cell membranes. The S1 RBD protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.



Structural model of the spike protein of SARS-CoV-2 (monomer) with its receptor binding domain (RBD) highlighted (red).


SARS-CoV-2 Spike S1 protein

Receptor binding domain (RBD)


Cat. no. P2020-001

Product Information

Quality Information (provided for each lot):




SDS-PAGE/Coll.Coomassie




	Area	Percent
1	325.991	1.436
2	589.627	2.598
3	1045.355	4.606
4	20733.217	91.359

Histogram (of marked lane in gel picture)


Activity Information (general information, not lot specific):




Analytic size exclusion chromatography (SEC) of the purified protein confirms a monomeric running behavior (arrow).



Bi-layer interferometry binding analysis, binding of hsACE2 (ECD), Cat# P2020-016.



Identification of major glycans by MS analysis after HILIC separation.



Stability during storage at 4°C determined by functional ELISA (binding to hsACE2 (ECD) - Cat# P2020-016; detection by mAb CR3022).