pro-TGF-beta 1, GFP/His-tag

Cat. no. P2020-118



Product Information

Protein: pro-TGF-beta 1, GFP/His-tag (~ 71.4 kDa)

Uniprot#: P01137

Sequence: MLSTSKTIDMELVKRKRIEAIRGQILSKLRLASPPSQGEVPPG

PLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVLMVETHNEIYDKFKQSTHSI YMFFNTSELREAVPEPVLLSRAELRLLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSD SPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSCDSRDNTLQVDINGFTTGRRGDLATIH GMNRPFLLLMATPLERAQHLQSSRHRRALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWI HEPKGYHANFCLGPCPYIWSLDTQYSKVLALYNQHNPGASAAPCCVPQALEPLPIVYYVG

RKPKVEQLSNMIVRSCKCS

Methionine at pos. 1 present due to cloning constraints, N-terminal GFP-fusion and His-tag

not shown in sequence.

Source: Recombinantly expressed in HEK293 cells.

Tag(s): GFP/His-tag, N-terminal

Purification: Purified by affinity chromatography and subsequent buffer exchange.

Formulation: PBS; pH 7.4.

Liquid, stored and shipped at -80 °C.

Purity: > 95 % (will be determined by densitometry of Coomassie stained gel, example next page)

Concentration: Will be determined by BCA-Assay.

Long-term storage: No recommendations.

Comment: Pro-TGF-beta 1 consists of the latency associated peptide (LAP) and the mature


transforming growth factor (TGF)-beta 1 together forming the small latent complex (SLC).

This results in three distinct protein bands in SDS-PAGE as illustrated on page 2.

Background Information:

Transforming growth factor beta 1 (TGF- β 1) belongs to the transforming growth factor beta superfamily of cytokines and is involved in a myriad of cellular functions including the control of cell growth, cell proliferation, cell differentiation, and apoptosis.

The transforming growth factor beta-1 is synthesized as precursor consisting of the latency-associated peptide (LAP, 249 aa) and the transforming growth factor beta 1 (112 aa). The precursor is also named small latend complex (SLC) or latent TGF-ß1. The precursor proprotein is cleaved in the Golgi apparatus by Furin, but the disulfide-linked homodimers of LAP and TGF-beta 1 remain non-covalently associated after secretion.




Structural model of pro-TGF-beta 1, His-tag




Product Information

Quality Information (provided for each lot):



SDS-PAGE/Coll.Coomassie



Histogram (of marked lane in gel picture)

SDS-PAGE running behavior of SLC:

