




As a precaution, gloves and eye protection should be worn when handling drug substances and using test units. Do not puncture any part of unit. Do not remove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.

Test Preparation

1 Remove the Smart-Tip[™] probe from test unit.



2 Sample suspected substance.

Powder

Touch the Smart-Tip[™] probe to the sample.




Tablet

Firmly press Smart-Tip[™] probe on the tablet while scrubbing in a circular motion until the tip is completely covered with substance.



Test Preparation


3 Lock Smart-Tip[™] probe into test unit. Press firmly.



STOP
Do not continue unless probe is locked below the Flex Pins and cannot be removed from the test unit.

Identification

4 With thumb and forefinger, squeeze chamber 1 until capsule breaks.



5 Gently shake test unit for 2-3 seconds.



Results



The appearance of a reddish-orange color within one minute indicates the presence of **Amphetamine** or **Methamphetamine**.



The appearance of a purple color within one minute indicates the presence of an opiate such as **Heroin**, **Morphine** or **Codeine**.



The appearance of a dark purple to black color indicates the presence of **Ecstasy/MDMA** or its analogues.



If **none** of the above colors develop within one minute, the above listed drugs are **not** present.




Proceed to Dilution (steps 6-9), regardless of whether or not a color appeared.



Dilution

6 With thumb and forefinger, squeeze chamber 2 until capsule breaks.




7 Gently shake test unit for 2-3 seconds.



Wait 5 minutes.
The test window will warm up.
This is normal.

Dilution

8 With thumb and forefinger, squeeze chamber 3 until capsule breaks.



9 Gently shake test unit for 2-3 seconds.



The test window will warm up.
This is normal.



Disposal After Dilution, the test kit is safe to dispose of in the trash.

NET 01-21

IMPORTANT INFORMATION

- IDenta Touch&Know[™] Discreet Drug Test Kits** are used to determine the presence of certain drugs in substances such as powders, crystals, granules, flakes, pills, tablets, or liquids.
- All results are presumptive.** If necessary, further testing by a competent drug testing lab can provide additional verification.
- A color change indicates a positive result for the listed drugs only if it occurs within the first minute.**

If there is **NO** color change within the time indicated, the listed drugs are **NOT** present in the tested sample. After a minute, the sampler and the liquid may begin changing color as a normal part of the testing process and this color change does **NOT** indicate a positive result.