



ferti·lome®

Weed-Out® Lawn Weed Killer

Contains TRIMEC® Herbicide

Kills troublesome broadleaf weeds:

Dandelion, Chickweed, Clover, Henbit & many others as listed

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For Additional
Precautionary Statements
and First Aid

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-
(2-methyl-4-chlorophenoxy) propionic acid... 5.45%
Dimethylamine salt of
2,4-dichlorophenoxyacetic acid..... 5.88%
Dimethylamine salt of dicamba
(3,6-dichloro-o-anisic acid)..... 1.21%

OTHER INGREDIENTS..... 87.46%
TOTAL:..... 100.00%

See back panel for substatement of ingredient statement

NET CONTENTS ONE PINT (16 FL. OZS. / 473 ML)

fertiome® Weed-Out Lawn Weed Killer

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person.
If On Skin Or On Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a Poison Control Center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

In the event of a medical or chemical emergency contact
North America 1-800- 255-3924



fertiome® is a registered trademark of Voluntary Purchasing Groups, Inc.

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 2217-597-7401
EPA Est. 7401-TX-01
Product Questions?
855-270-4776



PEEL FROM CORNER OF BOOKLET

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

10510-1015-CL



7 32221 10510 2

**KEEP FROM FREEZING.
STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL
PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets. Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not exceed specified dosages for any area.
- Do not spray carpetgrass, dichondra, or lawn where desirable clovers are present. Do not apply this product to "Floritam" St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact or spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Failure to follow above precautions may result in injury.

PRODUCT INFORMATION

Ferti•lome® Weed Out Lawn Weed Killer controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

WHEN TO USE

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

BROADCAST TREATMENTS

Limitations on broadcast applications to residential turfgrass:

The maximum application rate is 4 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 fl. oz. of product per 1,000 sq. ft.

BROADCAST TREATMENTS WITH HOSE-END SPRAYERS WITH DILUTION: Spray preparation for hose-end dilution sprayers:

Add the required amount of **ferti•lome® Weed-Out Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Table 1 - RATES OF ferti•lome® Weed-Out Lawn Weed Killer FOR VARIOUS LAWN GRASSES

Gallons of Water Spray - Hose End Sprayer with Dilution	Area to Treat (Sq. Ft.)	St. Augustinegrass, Centipedegrass, Bahia grass (fl. oz.)	Bermudagrass, Buffalograss, Bentgrass (fl. oz.)	Bluegrass Zoysiagrass, Fescue (fl. oz.)
1	250	½	⅔	⅔ to 1
4	1,000	2	2 ⅔	2 ⅔ to 4
8	2,000	4	5 ⅓	5 ⅓ to 8
20	5,000	10	13 ⅓	13 ⅓ to 20
32	8,000	16	21 ⅓	21 ⅓ to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
 2 fluid ounces = 4 tablespoons 4 fluid ounces = ½ cup
 8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
 32 fluid ounces = 1 quart (qt.)

SPOT TREATMENT WITH PRESSURE SPRAYERS

Limitations on spot treatments to residential turfgrass:
 Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4 fl. oz. per 1,000 sq. ft. per application. The maximum

number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND BAHIAGRASS:

Mix 2/3 fl. oz. (4 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. Do not apply to St. Augustinegrass, centipedegrass, or bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on "Floritam" St. Augustinegrass in Florida.

BERMUDAGRASS, BUFFALOGRASS, AND BENTGRASS:

Mix 1 fl. oz. (2 Tbs) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

KENTUCKY BLUEGRASS, FESCUE, PERENNIAL RYEGRASS, AND ZOYSIAGRASS:

Mix 1-1/3 fl. oz. (8 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

- BROADLEAF WEEDS -

Annual Yellow Sweetclover	Bristly Oxtongue	Chickweed, Common
Aster	Broadleaf Dock	Chickweed, Mouseear
Austrian Fieldcress	Broadleaf Plantain	Chicory
Bedstraw	Broomweed	Cinquefoil
Beggarticks	Buckhorn	Clover, Crimson
Betony, Florida	Buckhorn	Clover, Hop
Bindweed, Field	Plantain	Clover, Red
Bird Vetch	Bulbous	Clover
Bitter	Buttercup	Strawberry
Wintercress	Bull Thistle	Clover, Sweet
Bittercress, Hairy	Bullnettle	Clover, White
Bitterweed	Burclover	Cockle
Black-Eyed Susan	Burdock	Cocklebur
Black Medic	Burning Nettle	Common
Black Mustard	Burweed	Mullein
Blackseed	Buttercup	Creeping Jenny
Plantain	Buttonweed	Cudweed
Blessed Thistle	Canada Thistle	Curly Dock
Bloodflower	Carolina	Daisy, English
Milkweed	Geranium	Daisy Fleabane
Blue Lettuce	Carpetweed	Daisy, Oxeye
Blue Vervain	Catchweed	Dandelion
Bracted Plantain	Bedstraw	Dichondra
	Catnip	Dogfennel
	Catsear	Dollarweed

- BROADLEAF WEEDS (Continued) -

False Dandelion	Lambsquarters	Pepperweed
Falseflax	Lespedeza	Pigweed
False Sunflower	Mallow	Pineywoods
Fiddleneck	Matchweed	Bedstraw
Florida Pusley	Mexicanweed	Plains Coreopsis
Frenchweed	Milk Vetch	Plantain
Galinsoga	Morningglory	Poison Ivy
Goldenrod	Mouseear	Poison Oak
Ground Ivy	Hawkweed	Pokeweed
Gumweed	Mugwort	Poorjoe
Hairy Fleabane	Musk Thistle	Prairie
Hawkweed	Mustard	Sunflower
Healall	Narrowleaf	Prickly Lettuce
Heartleaf	Plantain	Prickly Sida
Drymary	Narrowleaf	Prostrate
Heathaster	Vetch	Knotweed
Hedge Bindweed	Nettle	Prostrate
Hedge Mustard	Orange	Pigweed
Henbit	Hawkweed	Prostrate Spurge
Hoary Cress	Oriental	Prostrate
Hoary Plantain	Cocklebur	Vervain
Hoary Vervain	Oxalis	Puncturevine
Horsenettle	Parsley-Piert	Purslane,
Jimsonweed	Parsnip	Common
Knawel	Pearlwort	Ragweed
Knotweed	Pennycress	Red Sorrel
Kochia	Pennywort	

- BROADLEAF WEEDS (Continued) -

Redroot Pigweed	Spurweed Stinging Nettle	Wild Buckwheat Wild Carrot
Redstem Filaree	Strawberry India Mock	Wild Four-O'Clock Wild Garlic
Rough Cinquefoil	Tall Nettle Tall Vervain	Wild Geranium Wild Lettuce
Rough Fleabane	Tansy Ragwort Tansy Mustard	Wild Marigold Wild Mustard
Russian Pigweed Russian Thistle	Tanweed Thistle	Wild Onion Wild Parsnip
Scarlet Pimpernel	Trailing Crownvetch	Wild Radish Wild Rape
Scotch Thistle	Tumble Mustard	Wild Strawberry Wild Sweet Potato
Sheep Sorrel	Tumble Pigweed Velvetleaf	Wild Vetch Woodsorrel
Sherherdspurse	Venice Mallow Virginia	Woolly Croton Woolly
Slender Plantain	Virginia Buttonweed	Morningglory Woolly Plantain
Smallflower Galinsoga	Virginia Creeper Virginia	Wormseed Yarrow
Smooth Dock	Pepperweed Wavyleaf	Yellow Rocket Yellowflower
Smooth Pigweed	Bullthistle Western Clematis	Pepperweed
Sorrel	Western Salsify White Mustard	
Sowthistle	Wild Aster	
Spanishneedles		
Speedwell		
Spiny Amaranth		
Spiny Cocklebur		
Spiny Sowthistle		
Spotted Catsear		
Spotted Spurge		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.
Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

This page left blank intentionally.