

BLACKBURN DRUG

PHONE: 307-201-1159 L FAX: 307-201-1553
BLACKBURNDRUG.COM
545 N CACHE - UNIT 10-S JACKSON WY 83001

PATIENT NAME _____ DOB _____
ADDRESS _____ PHONE _____
ALLERGIES _____

SEMAGLUTIDE SUBLINGUAL SUSPENSION @ 1 MG / ML

PLACE 0.25 ML UNDER THE TONGUE EVERY MORNING FOR 8 DAYS. THEN INCREASE TO 0.5 ML UNDER THE TONGUE EVERY MORNING. DO NOT EAT OR DRINK ANYTHING AFTER FOR 30 MINUTES.
15 ML - 5 REFILLS

QSYMIA COMBINATION COMPOUND

PHASE 1

PHENTERMINE 10 MG / TOPIRAMATE 12.5 MG / CHROMIUM 100 MCG SR CAPSULES
TAKE 1 CAPSULE BY MOUTH DAILY FOR 14 DAYS THEN 2 CAPSULES DAILY EVERY MORNING
#46 CAPSULES

PHASE 2

PHENTERMINE 25 MG / TOPIRAMATE 40 MG / CHROMIUM 100 MCG SR CAPSULES
TAKE 1 CAPSULE BY MOUTH EVERY MORNING.
#30 CAPSULES

PHASE 3

PHENTERMINE 35 MG / TOPIRAMATE 60 MG / CHROMIUM 100 MCG SR CAPSULES
TAKE 1 CAPSULE BY MOUTH EVERY MORNING.
#30 CAPSULES

CONTRAVE COMBINATION COMPOUND

STARTING DOSE: NALTREXONE 8 MG CAPS #70 AND BUPROPION ER 100 MG #63
WEEK 1 - TAKE 1 NALTREXONE AND 1 BUPROPION IN THE MORNING
WEEK 2 - TAKE 1 NALTREXONE AND 1 BUPROPION TWICE A DAY
WEEK 3 - TAKE 2 NALTREXONE AND BUPROPION IN THE MORNING AND 1 OF EACH IN PM
WEEK 4 - TAKE 2 NALTREXONE TWICE A DAY, 2 BUPROPION IN AM AND 1 IN PM

CONTINUING DOSE:

NALTREXONE 16 MG CAPS #60 AND BUPROPION ER 100MG #90
TAKE 1 NALTREXONE TWICE A DAY, 2 BUPROPION IN AM AND 1 IN PM

PHYSICIAN NAME _____ DATE _____
SIGNATURE _____ DEA _____