

It is important to read this leaflet carefully before using Xailin Gel. The information provided summarises the important points about the product but if you have any further questions then please ask your doctor, pharmacist or optician.

What is Xailin Gel?

Xailin Gel is a multi-dose eye gel for the long lasting relief of dry eye sensations. The pack contains 10 g of sterile eye gel.

Composition:

Xailin Gel contains Carbomer 0.2% w/w, Sodium perborate, and excipients.

Xailin Gel contains a disappearing preservative that turns into water and oxygen upon contact with your eye, thereby minimizing the irritation that may be caused by traditional preservatives while still providing the convenience of a multidose form. All you feel is the cooling and soothing relief of the gel.

When should it be used?

Xailin Gel is used to lubricate and protect the eye in certain eye conditions and provide long lasting relief of dry eye sensations. Xailin Gel provides cooling and soothing relief from dry eye sensations, such as soreness, irritation or a gritty sensation. Dry eye symptoms occur as a result of ocular surface disorders that can be caused by a number of different reasons where the normal tears are either of poor quality or produced in too little quantity to keep the eye sufficiently moist. Xailin Gel can be used by contact lens wearers, but lenses are recommended to be removed prior to use.

How to use:

The usual dosage is one drop to be applied two to four times a day as needed, or as directed by your healthcare practitioner who may advise you to use this product in a different way. No dosage adjustment is required in children. Before opening a new tube of Xailin Gel, please check that the safety ring on the cap of the tube is unbroken.

1. Rotate the cap anticlockwise to break the seal
2. Wash and dry your hands well before use
3. Remove the outer cap
4. If you are a contact lens wearer remove your lenses
5. Tilt your head back and look up
6. Gently pull down your lower eyelid and squeeze one drop of Xailin Gel into the gap between your eye and eyelid
7. Blink your eye a few times
8. If you think that you have missed the eye, then insert another drop
9. Carefully wipe away any excess Xailin Gel around the eyelid with a tissue
10. Repeat the process for the other eye, if required
11. Replace the outer cap on the container, trying not to touch the applicator tip with anything, including your eye or fingers
12. Leave at least 30 minutes between administrations of any other ophthalmic drugs

If you are unsure about how to use Xailin Gel or if your eye conditions persist, consult your doctor, pharmacist or optician.

Precautions and warnings:

- Do not use if the safety ring on the cap of the tube is broken
- Do not use if you are allergic to the product
- Do not touch the tip of the tube with your eye or any surface
- Do not share Xailin Gel with any other person
- Xailin Gel can cause transient mild burning sensation and or blurring of vision after application. Wait until your vision is clear before driving or using machinery
- Do not use after the expiry date shown on the carton or printed on the crimp of the tube
- Store between 1°C and 25°C
- Discard the contents 28 days after first opening
- Keep out of the reach and sight of children
- External ocular use only, do not swallow
- Consult your doctor, pharmacist or optician before you use Xailin Gel if you are pregnant or breast-feeding
- Dispose of the used tube according to your local waste management regulations

S-CAREPHD-14-PLH-PV2

nicox Distributed by Nicox Pharma, CS10313, Sophia Antipolis, 06560 Valbonne, France - nicoxpharma@nicox.com

Medicom Healthcare Ltd., 235 Hunts Pond Road, Fareham, Hampshire, PO14 4PJ, U.K.

Store between 1°C and 25°C
Discard the contents 28 days after first opening
Caution, see instructions for use
Consult Instructions for use

STERILE A Sterilized using aseptic processing techniques

Xailin is a trademark of Nicox
Last revision of the text: October-2013

