



Advance 500 mg Tablets

Paracetamol

Please read right through this leaflet before you start using this medicine.

This medicine is available without prescription, but you still need to use Panadol Advance 500 mg Tablets carefully to get the best results from them.

- Keep this leaflet, you may need to read it again.
- If you have any questions, or if there is anything you do not understand, ask your pharmacist.
- Contains paracetamol.

In this leaflet:

1. What Panadol Advance 500 mg Tablets do
2. Check before you take Panadol Advance 500 mg Tablets
3. How to take Panadol Advance 500 mg Tablets
4. Possible side effects
5. How to store Panadol Advance 500 mg Tablets
6. Further information

1. What Panadol Advance 500 mg Tablets do

Panadol Advance 500 mg Tablets are faster absorbed than standard paracetamol tablets. They are suitable for the relief of headache, tension headache, migraine, backache, rheumatic and muscle pain, toothache and period pain.

They also relieve sore throat and the fever, aches and pains of colds and flu and are recommended for the relief of pain due to mild arthritis.

The active ingredient is paracetamol which is a painkiller and also reduces your temperature when you have a fever.



2. Check before you take Panadol Advance 500 mg Tablets



Do not take Panadol Advance 500 mg Tablets:

- if you have ever had an **allergic reaction** to paracetamol or to any of the other ingredients (listed in Section 6)
- if you are **taking any other prescription or non-prescription medicines containing paracetamol** to treat pain, fever, symptoms of cold and flu, or to aid sleep.



Ask your doctor before you take this medicine:

- if you suffer from **mild arthritis** and need to take painkillers every day.
- if you have **liver** or **kidney** problems
- if you are **underweight** or **malnourished**
- if you regularly drink alcohol
- if you have a **severe infection** as this may increase the risk of **metabolic acidosis**.

Signs of metabolic acidosis include:

- deep, rapid, difficult breathing
- feeling sick (nausea), being sick (vomiting)
- loss of appetite

Contact a doctor immediately if you get a combination of these symptoms.

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.



If you are taking other medicines

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly **metoclopramide** or **domperidone** (for **nausea** [feeling sick] or **vomiting** [being sick]) or **colestyramine** (to lower **blood cholesterol**). If you take **blood thinning drugs (anticoagulants** e.g. **warfarin**) and you need to take a pain reliever on a daily basis, **talk to your doctor** because of the risk of bleeding. But you can still take **occasional** doses of Panadol Advance 500 mg Tablets at the same time as anticoagulants.

NPA005270





Pregnancy and breast feeding

Talk to your **healthcare professional** before taking Panadol Advance 500 mg Tablets if you are **pregnant**. You can take this product whilst **breast feeding**.

3. How to take Panadol Advance 500 mg Tablets



Adults (including the elderly) and children aged 16 years and over:
Swallow 1-2 tablets every 4 to 6 hours as needed.
Do not take more than 8 tablets in 24 hours.

Children aged 10-15 years:
Give 1 tablet every 4 to 6 hours as needed. Do not give more than 4 tablets in 24 hours.

Do not give for more than 3 days at a time unless your doctor tells you to.



- **Do not take more frequently than every 4 hours.**
- **Do not take more than the recommended dose.**
- **Do not give to children under 10 years.**

If you take too many tablets

Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

If your symptoms continue or your headache becomes persistent, **see your doctor**.

4. Possible side effects

Like all medicines, Panadol Advance 500 mg Tablets can have side effects but not everybody gets them. A small number of people have had side effects. Very rare cases of serious skin reactions have been reported.

Stop taking the medicine and **tell your doctor immediately** if you experience:

- **Allergic reactions** which may be severe such as skin rash and itching sometimes with swelling of the mouth or face or shortness of breath
- **Skin rash or peeling, or mouth ulcers**
- **Breathing problems.** These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin
- Unexplained **bruising or bleeding**
- **Nausea**, sudden **weight loss, loss of appetite** and **yellowing of the eyes and skin.**

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Panadol Advance 500 mg Tablets

Keep out of the sight and reach of children.

Do not use this medicine after the 'EXP' date shown on the pack. Store below 30°C.

6. Further information

Active ingredient: Each tablet contains Paracetamol 500 mg.

Other ingredients: Pregelatinised starch, calcium carbonate, alginate acid, crospovidone, povidone, magnesium stearate and colloidal anhydrous silica.

The tablet coating contains opadry white and carnauba wax.

Packs of Panadol Advance 500 mg Tablets contain 16 or 30 tablets.


The marketing authorisation holder is GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

The manufacturer is GlaxoSmithKline Dungarvan Ltd., Co. Waterford, Ireland.

This leaflet was last revised in July 2021.

Trade marks are owned by or licensed to the GSK group of companies.

NPA005270



PHARMA CODE REF. IS: XX

