

Rapilose

Glucose Gel 40%

- Easy-to-swallow fast acting gel providing rapidly absorbed carbohydrates to increase the body's blood sugar levels quickly
- Available in great tasting 'fruits of the forest' flavour in packs of three convenient easy-to-use 25g tubes
- Each tube delivers 10g of anhydrous glucose
- Suitable for dispensing against prescriptions written for glucose or dextrose gel 40% w/w

Rapilose Gel

Fast acting glucose gel 40% w/w
Forest fruits flavour

Each 25g tube contains 10g of glucose gel



3 x 25g Easy to use tubes



EAN 5060258140014

PIP 365-7707

NHS Price £7.16

www.rapilose.com



Why use RapiLOSE Gel?

With no measuring or weighing of food necessary, RapiLOSE Gel provides rapidly absorbed carbohydrates to increase the body's blood sugar levels quickly.

Available in great tasting forest fruits flavour and easy-to-swallow, fast-acting RapiLOSE Gel contains no colour additives and is gluten, lactose, fat, caffeine and alcohol free.

RapiLOSE Gel is particularly useful if you are eating less (e.g. trying to lose weight, upset stomach), or exercising, or in any situation where meals are delayed, skipped or forgotten. Having RapiLOSE fast acting carbohydrate available provides you with the confidence to get-on with your life.

Our pocket-sized tubes are highly portable, water resistant and fit discretely into a bag or a pocket, so you can take them with you everywhere and are always available when required; at home, at school or at work.

What ingredients does RapiLOSE Gel contain?

Each 25g tube delivers 10g 40% w/w glucose. RapiLOSE Gel contains water, glucose syrup, glucose monohydrate, flavouring, citric acid (acidity regulator), potassium sorbate, sodium benzoate (preservatives), ascorbic acid (antioxidant).

For further information contact:

Penlan Healthcare Limited

Calverley House
55 Calverley Road
Royal Tunbridge Wells
TN1 2TU
United Kingdom

T 01892 704 077
info@rapilose.com

For ordering information contact:

Aspire Pharma Limited

Antrobus House Business Centre
18 College Street
Petersfield
GU31 4AD
United Kingdom

T 01730 234 527
sales@aspirepharma.co.uk

What are the Directions for Use for RapiLOSE Gel?

The product is for oral consumption only and is not recommended for children under the age of two years.

Simply twist-off the cap, squeeze the gel into your mouth and swallow. Alternatively, it can be squeezed inside the cheek and the outside of the cheek rubbed gently to aid absorption. Repeat, if necessary. Speak to your healthcare adviser regarding your individual carbohydrate requirements.

Our packaging is functional and easy-to-use, even for patients where dexterity is affected or eyesight is poor.

RapiLOSE Gel must never be given to people who are unconscious. Do not use after the 'Expiry Date' printed on the carton or if the product has been previously opened.

Where should I keep RapiLOSE Gel?

RapiLOSE Gel should be kept with you or a responsible person at all times. It has an 18-month shelf-life kept below 25°C and can be safely carried in your pocket, handbag, school bag or sports kit, stored beside your bed or in a shaded area in the car. Note the gel may become runny in the heat and more difficult to squeeze from the tube if it becomes very cold.

EAN 5060258140014
PIP 365-7707

