

HiScript® III RT SuperMix for qPCR(+gDNA wiper)

R323



Product Description

HiScript III RT SuperMix for qPCR (+gDNA wiper) is an upgraded version of HiScript II Q RT SuperMix for qPCR (+gDNA wiper), including HiScript III Reverse Transcriptase, a new generation of reverse transcriptase with optimized Buffer. This kit further improves the efficiency of cDNA synthesis, and is suitable for two-step qRT-PCR detection. The 4 × gDNA wiper Mix in the kit completely removes residual genomic DNA from the RNA template, ensuring more reliable quantitative results. It simplifies qPCR primer design without the need to design primers across introns; 5 × HiScript III qRT SuperMix contains all components required for the reverse transcription reaction and it can be started rapidly after the addition of template RNA and RNase-free ddH₂O, and the gDNA wiper is terminated to ensure the integrity of the cDNA. It is compatible with dye-based and probe-based qPCR, enabling high-performance gene expression analysis.

Components

Components	R323-01 100 rxns (20 µl/rxn)
<input type="checkbox"/> RNase-free ddH ₂ O	2 × 1 ml
<input checked="" type="checkbox"/> 4 × gDNA wiper Mix	400 µl
<input checked="" type="checkbox"/> 5 × HiScript III qRT SuperMix ^a	400 µl
<input checked="" type="checkbox"/> 5 × No RT Control Mix ^b	40 µl

a. It contains Buffer, dNTP, HiScript III Reverse Transcriptase, RNase inhibitor, Random primers/Oligo (dT)₂₀VN primer Mix.

b. Except for HiScript III Reverse Transcriptase, other components are the same as 5 × HiScript III qRT SuperMix, using for No RT control preparation.

Storage

Store at -30 ~ -15°C and transport at ≤0°C.

Applications

This product is suitable for reverse transcription of animal, plant and microbial RNA, and the products are compatible with dye-based and probe-based qPCR.

Self-prepared Materials

Materials:

- RNase-free centrifuge tube (1.5 ml), RNase-free PCR tube (0.2 ml), RNase-free tips
- Pipette, PCR instrument, ice or ice box

RNA

- High quality RNA is essential for obtaining high quality cDNA. Please verify the RNA integrity by electrophoresis before the experiment.

qPCR Reagent Selection Guide:

- The 1st strand cDNA product can be used as the template for qPCR directly. For PCR, it is recommended that the volume of the template cDNA product should not exceed 1/10 of the total volume of qPCR reaction. AceQ Universal U⁺ Probe Master Mix V2 (Vazyme #Q513) or ChamQ Universal SYBR qPCR Master Mix (Vazyme #Q711) can be selected as the qPCR reagent.

Notes

1. The 4 × gDNA wiper Mix, 5 × HiScript III qRT SuperMix and 5 × No RT Control Mix contain high concentration of glycerol. Please centrifuge briefly and pipette up and down to mix thoroughly before use.
2. It is recommended to add no more than 1 µg of total RNA to the 20 µl reverse transcription reaction system. If target genes with low expression levels, the amount of total RNA can be up to 5 µg. Otherwise, the amount of RNA added is too high, which may exceed the linear range of subsequent qPCR.
3. The cDNA products are only suitable for qPCR reactions and not suitable for long-fragment PCR amplification of downstream experiments such as cloning. If necessary, HiScript III 1st Strand cDNA Synthesis Kit (+ gDNA wiper) (Vazyme #R312) is recommended.
4. Reverse transcription can be performed directly with 5 × HiScript III qRT SuperMix without the genome removal step, but do not use 4 × gDNA wiper Mix with reagents without genome removal module, as the Mix in the kits without genome removal module does not contain the components to terminate the gDNA wiper reaction, which may affect subsequent qPCR results.

Experiment Process

1. Removal of genomic DNA

Mix the following components in a RNase-free centrifuge tube:

RNase-free ddH ₂ O	to 16 µl	□
4 × gDNA wiper Mix	4 µl	■
Template RNA	Total RNA: 1 µg - 1 µg	

Mix gently with a pipette and incubate at 42°C for 2 min.

2. Preparation of Reverse Transcription Reaction Mixture

Add 5 × HiScript III qRT SuperMix to the mixture of previous step:

5 × HiScript III qRT SuperMix	4 µl	■
Mixture from Step 1.	16 µl	

Mix gently with a pipette.

No RT Control Reaction (Optional)

No RT Control Reaction is a negative control which contains no Reverse Transcriptase and is used to indicate whether there is residual genomic DNA in RNA template.

Mix the following components in a RNase-free centrifuge tube:

5 × No RT Control Mix	4 µl	■
Mixture from Step 1.	16 µl	

Mix gently with a pipette.

3. Perform the reverse transcription reaction under the following conditions

37°C*	15 min
85°C	5 sec

* For template with complex secondary structures or high GC content, the temperature can be increased to 50°C, which will benefit the yield.

The product can be directly used in qPCR reactions or be stored at -20°C for 6 months. It is recommended to store in aliquots at -70°C for long term storage, and cDNA should be avoided repeated freezing and thawing.