

VanaLice[®] Lice Killing Gel

Drug Facts

Active ingredients	Purpose
Piperonyl butoxide 3.5%.....	Lice treatment
Pyrethrum extract (equivalent to 0.3% pyrethrins).....	Lice treatment

Uses treats head, pubic (crab), and body lice

Warnings

For external use only.

Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **Important: Read warnings before use**
- **adults and children 2 years and over:**

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

Treat

- apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for 10 minutes, but no longer
- wash area thoroughly with warm water and soap or shampoo
- for head lice, towel dry hair and comb out tangles

Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours. ▶

Drug Facts (continued)

- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments
- **children under 2 years:** ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat
- keep carton for important product information
- see package insert for additional information

Inactive ingredients

carbomer, citronellyl acetate, coconut oil, DMDM hydantoin, PEG-40 hydrogenated castor oil, isopropanol, isopropyl myristate, lemongrass oil, purified water, sodium hydroxide, tea tree oil, tocopheryl acetate (vitamin E)

Questions or comments?

Call 1-888-535-0305 9 a.m. - 5 p.m. CST

GM Pharmaceuticals, Inc.

Distributed by: GM Pharmaceuticals, Inc. Arlington, TX 76015