

FIFRA Section 2(ee)
Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.



PESTS

For Control of Light Brown Apple Moth (LBAM) (*Epiphyas postvittana*), Banana Moth (*Opogona sacchari*), European Grapevine Moth (*Lobesia botrana*) and European Pepper Moth (*Duponchelia fovealis*).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For Light Brown Apple Moth: Apply when newly hatched larvae appear and before leaves are rolled or webbing is significant.

For Banana Moth: Drench bark to newly emergent shoots following pruning or apply to susceptible plant tissues when Banana Moth larvae are active.

For European Grapevine Moth: Apply at blackhead egg stage or when larvae are newly hatched before leaves are rolled, or larvae have entered fruit.

For European Pepper Moth: Begin applications at egg lay and continue at 3-5 day intervals throughout larval feeding period.

For All Pests: Continue applying as part of a normal spray program until pest is adequately controlled. Apply when caterpillars are actively

EPA Reg. No. 70051-47

feeding. To be effective, Agree WG® spray must be deposited at the larval feeding site. Agree WG can be applied by ground or air in water sufficient to insure thorough and even coverage. Thorough and uniform crop coverage is required for adequate insect control. Applications at higher water volumes have demonstrated improved control of targeted pests. Early morning or evening applications, when air is calm, are generally best for aerial applications.

All applicable directions, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. This recommendation must be in the possession of the user at the time of pesticide application.

Continued on reverse side





CROPS	RATES
Cole Crops: Terrestrial Chinese Vegetables, Terrestrial Green Leafy Vegetables, Celery	0.5 to 2.0 pounds per acre
Beans, Peas	0.5 to 2.0 pounds per acre
Tomatoes, Peppers	0.5 to 2.0 pounds per acre
Tobacco	1.0 to 2.0 pounds per acre
Cotton	0.25 to 2.0 pounds per acre
Terrestrial Ornamentals and Flowers	0.5 to 2.0 pounds per acre
Stone Fruit: Peaches, Nectarines, Plums, Prunes, Cherries	1.0 to 2.0 pounds per acre
Tree Nuts: Almonds, Filberts, Walnuts, Pecans	1.0 to 2.0 pounds per acre
Pistachios	1.0 to 2.0 pounds per acre
Greenhouse Vegetables: Tomatoes, Cole Crops, Peppers	0.5 to 2.0 pounds per acre
Grapes	1.0 to 2.0 pounds per acre
Corn	1.0 to 2.0 pounds per acre
Terrestrial Small Fruits and Berries	0.5 to 2.0 pounds per acre
Cranberries	1.0 to 2.0 pounds per acre
Cucurbits	0.5 to 2.0 pounds per acre
Potatoes	1.0 to 2.0 pounds per acre
Soybeans	0.25 to 2.0 pounds per acre
Pome Fruits: Apples, Pears	1.0 to 2.0 pounds per acre
Alfalfa and Other Forage Crops	0.5 to 2.0 pounds per acre
Citrus: Oranges, Grapefruit, Tangerine	1.0 to 2.0 pounds per acre
Sugarbeets, Radish and Other Root Crops, Leaves of Root and Tuber Vegetables	0.5 to 2.0 pounds per acre
Herbs and Spices	0.5 to 2.0 pounds per acre
Peanuts	1.0 to 2.0 pounds per acre
Peppermint, Spearmint	1.0 to 2.0 pounds per acre
Artichokes	1.0 to 2.0 pounds per acre
Bulb Vegetables: Onions, Garlic	1.0 to 2.0 pounds per acre