


# BIOMIN<sup>®</sup> COPPER

1-0-0

Amino Acid Chelated Mineral for Soil and Foliar Application



## Product Information

BIOMIN COPPER contains amino acid chelated copper for soil and/or foliar application. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct copper deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

### GUARANTEED ANALYSIS

Total Nitrogen (N)	1.0%
1.0% Water Soluble Nitrogen	
Copper (Cu)	4.0%
4.0% Chelated Copper	

Derived from copper sulfate and hydrolyzed vegetable protein.

**KEEP OUT OF REACH OF CHILDREN**

**SHAKE WELL BEFORE USE**

**PREVENTS AND CORRECTS  
COPPER DEFICIENCIES**

NET CONTENTS: 1 Gallon

Weight per Gallon: 9.5 lb (4.3 kg) @ 68°F

U.S. Pat.# 5,504,055

Lot #: 301183



## Directions and General Recommendations

### SOIL APPLICATION

BIOMIN COPPER can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre depending on soil conditions and application methods. For maintenance, 1 quart per acre is recommended. BIOMIN COPPER should be diluted with enough water to cover the entire acre.

### FOLIAR APPLICATION

**FIELD CROPS:** Apply 1-4 pints per acre (1.25-5 liters/hectare) of BIOMIN COPPER when crops are about 4-6 inches (10-20 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

**FRUIT TREES AND GRAPES:** Apply 1-4 pints per acre (1.25-5 liters/hectare) starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

**VEGETABLE CROPS:** Apply 1-4 pints per acre (1.25-5 liters/hectare) of BIOMIN COPPER at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1 - 3 weeks if necessary.

### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

**ALMONDS, WALNUTS, PISTACHIOS AND OTHER NUT CROPS:** Apply 1-4 pints per acre (1.25-5 liters/hectare) before bloom. Repeat the application if necessary.

For best results, apply BIOMIN COPPER when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water. BIOMIN COPPER will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN COPPER can be applied to: Strawberry, Lettuce, Tomato, Celery, Stone Fruits, Vegetables, Grapes, Commercial Turf, And Ornamental Gardens.

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006 U.S.A. Phone: 805-650-8933 Website: [www.jhbiotech.com](http://www.jhbiotech.com)

**JH Biotech Inc.**

Copyright © 2013 All rights reserved

0302-D-B-2