# Clonex Rooting Gel

# Rooting Compound

# Keep Out of Reach of Children CAUTION

# 72


See Side Panel for Additional Precautionary Statements

ACTIVE INGREDIENT	
Indole-3-Butyric Acid	0.31%
OTHER INGREDIENTS	<u>99.69%</u>
TOTAL	100.00%

EPA Registration No. 79664-1 EPA Establishment No. 79463-MI-001

Net Content: 50 ml / 1.7 fl oz Net Contents: 100 ml / 3.4 fl oz Net Contents: 250 ml / 8.5 fl oz

Manufactured by:
Growth Technology Ltd.
66, Taunton Trading Estate
Taunton TA2 6RX United Kingdom
Taunton, U.K.
www.growthtechnology.com



#### **DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CLONEX Rooting Gel is used to ensure root production on cuttings of plants by supplying essential hormones needed for accelerated root development. CLONEX Rooting Get may be used on all types of cuttings, including woody, herbaceous, and flowering ornamental species, vegetables, fruit trees and small fruits.

Dip cutting in Clonex Rooting Gel to desired depth. Insert cutting into rooting medium. Mist cuttings and place in propagator or a warm, clean, moist and humid environment. Look for root development in 1 to 2 weeks.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS** 

**CAUTION** 

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

	FIRST AID
If on skin or clothing	<ul> <li>Take off contaminated clothing</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed	<ul> <li>Call poison control center or doctor immediately for treatment advice</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product contained	er or label with you when calling a poison control center or

Have the product container or label with you when calling a poison control center of doctor, or going for treatment. For emergency medical assistance, call 1-800-222-1222.

## **ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

					•			•						
•	•		٠		•			•			•	•	٠	
	ŧ			•		•				•				
	•		4								•	٠	•	2 1
			٠			٠						•		
•		•	,			ŧ	•		,	•				

# STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store only in original container, in a cool dry place out of the reach of children and out of direct sunlight.

# Disposal:

- If empty: Do not reuse this container. Place in trash or offer for recycling if available.
- If partially filled: Your local government may forbid liquids and pesticides in their landfills. Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

11-11-2004