CRESCENDO

8-2-2

ORGANIC WITH 5% CALCIUM

APPLY ONLY AS DIRECTED

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.



Distributed by Seven Springs Farming Supplies LLC 426 Jerry Lane, Check, VA 24072 800-540-9181 www7springsfarm.com order@7springsfarm.com

NET WEIGHT 50 LB. (22.7 KG)

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	. 8.0%
3.2% WATER SOLUBLE NITROGEN	
4.8% WATER INSOLUBLE NITROGEN ³	k
AVAILABLE PHOSPHATE (P2O5)	2.0%
SOLUBLE POTASH (K2O)	2.0%
CALCIUM (Ca)	5.0%

PLANT NUTRIENTS DERIVED FROM: Composted Poultry Layer Manure, Feather Meal, Sulfate of Potash

COVERAGE:

FOR TURF, SHRUBS & TREES -- APPLY AT THE FIRST SIGN OF SPRING GROWTH.
REAPPLY, AS NEEDED, EVERY 6-8 WEEKS THROUGHOUT THE GROWING SEASON.

TURF:

FOR 1.0 LB. OF NITROGEN, APPLY AT A FULL RATE OF 12.5 LBS. PER 1,000 SQ. FT. ONE 50 LB. BAG OF 8-2-2 WILL COVER 4,000 SQ. FT. AT THE FULL RATE. FOR 0.5 LB. OF NITROGEN, APPLY AT A FULL RATE OF 6.25 LBS. PER 1,000 SQ. FT. ONE 50 LB. BAG OF 8-2-2 WILL COVER 8,000 SQ. FT. AT THE HALF RATE.

LANDSCAPE APPLICATION:

SHRUBS - TOPICALLY APPLY OR LIGHTLY SCRATCH INTO THE SOIL SURFACE AT 3.5 LBS. PER 1,000 SQ. FT.

FLOWERS - APPLY 2 LBS. PER 100 SQ. FT.. LATER IN THE GROWING SEASON, A SECOND APPLICATION CAN BE MADE.

TREES - APPLY 6.5 CUPS OF 8-2-2 FOR EACH 1- INCH OF TRUNK DIAMETER.

VEGETABLES:

APPLY AT THE RATE OF 22 POUNDS PER 1000 SQ.FT. OR 1.0 CUPS PER 10 FT. REAPPLY AS NEEDED. SIDE DRESS AT HALF THE RATE 4 TO 6 WEEKS AFTER PLANTING.

MANUFACTURED/GUARANTEED BY BIOSYSTEMS, LLC 2103 REGAN ROAD. BLACKSBURG. VA 24060 / 540-239-5884

^{*4.8%} slowly available nitrogen from poultry manure, feather meal.