

LALSTOP G46 WG



Biofungicide – Controls/prevents soil-borne and foliar pathogens
For use in organic production

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

CAUTION – KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: *Gliocladium catenulatum*

Strain J1446* 93.0%
Other Ingredients: 7%
Total: 100%

* Contains a minimum of 1×10^9 cfu/g of active ingredient.

EPA Reg. No.: 64137-13

EPA Est. No.:

Lot No:

Use Before:

FIRST AID

	Call a poison control center or doctor immediately for treatment advice.
If Swallowed	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
If on Skin or Clothing	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, waterproof gloves and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixer/loaders and applicators must wear a NIOSH-approved particulate respirator with any R or P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS: When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

NET WEIGHT: 2 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

Do not enter or allow workers to enter the treated greenhouse or enclosed space until the ventilation requirements in 40 CFR 170.405(b)(3) have been met and the REI of 4 hours has expired. Until then, only handlers wearing the appropriate personal protective equipment can enter the greenhouse or enclosed space.

EXCEPTION: If the product is soil incorporated or soil injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

PRODUCT INFORMATION

LALSTOP® G46 WG controls seed-borne and soil-borne plant diseases such as damping-off, root and stem rot, charcoal rot and wilt caused by *Alternaria**, *Bipolaris**, *Botrytis*, *Cladosporium**, *Colletotrichum**, *Fusarium*, *Macrophomina**, *Mycosphaerella**, *Penicillium**, *Phytophthora*, *Plasmodiophora**, *Plicaria**, *Pyrenochaeta**, *Pythium*, *Rhizoctonia**, *Sclerotinia** and *Verticillium**; certain storage diseases caused by *Helminthosporium** and *Rhizoctonia**; and foliar diseases like grapevine trunk diseases* (several pathogens) as well as diseases caused by *Alternaria**, *Anthraxnose (Colletotrichum)**, *Bipolaris**, *Botrytis*, *Cladosporium**, *Didymella*, *Fusarium*, *Monilinia**, *Mycosphaerella**, *Penicillium** and *Sclerotinia** on (field-grown) vegetables, herbs, ornamentals, flowering plants, bedding plants, berries, cereals, legumes, pome and stone fruits, nut trees, tree and forest seedlings, turf and other labeled crops. LALSTOP® G46 WG also suppresses powdery mildew symptoms on these (field-grown) crops*. For use on green and flowering plants.

***NOT FOR USE IN CALIFORNIA.**

USE INSTRUCTIONS

Apply LALSTOP® G46 WG by spraying or drenching the growth substrate, by incorporation into the growth substrate, as a foliar spray, by fogging, or by hydroponic or chemigation application in the field as an aqueous suspension.

LALSTOP® G46 WG has been evaluated for phytotoxicity on a variety of crops under various normal growing conditions. However, testing all crop varieties, in all mixtures and combinations, is not feasible. Prior to treating entire crop, test a small portion of the crop for sensitivity.

Tank Mixes: Do not combine LALSTOP® G46 WG in the spray tank with other pesticides, surfactants, adjuvants, or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, effective, and non-injurious under your use conditions. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

To ensure compatibility of tank-mix combinations, they must be evaluated prior to use. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to one quart of water with agitation. Add dry formulations first, then flowables, and then emulsifiable concentrates last. After thoroughly mixing, let this mixture stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Surfactants are not generally needed. A non-ionic surfactant may be used in combination with LALSTOP® G46 WG for crops with waxy/hard-to-wet leaf surfaces.

Integrated Pest Control Programs: LALSTOP® G46 WG can be used in integrated pest control programs. Always check the compatibility of LALSTOP® G46 WG with other biological fungicides or chemical pesticides with the local distributor.

APPLICATION RATES

PRE-HARVEST INTERVAL (PHI) = 0 DAYS

Beans, Lentils, and Peas

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
In-Furrow	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Apply in 5 – 25 gallons of water per acre as a directed spray into the seed furrow just before seeds are covered. Use high rate for high disease pressure and/or high plant density.

Beans, Lentils, and Peas continued.

Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.
---------------------------------------	--

Berries and Small Fruits, including: Blackberry, Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Kiwifruit, Loganberry, Raspberry, Strawberry, and Grape (e.g., Table and Wine)

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Cuttings, Bare Root or Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Cucurbit Vegetables, including: Cucumber, Cantaloupe, Melon, Gourd, Pumpkin, Squash (e.g., Winter), and Watermelon

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).

Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Flowers, Bedding Plants, Foliage and Potted Plants, and Ornamentals, including: Achillea, African Violet, Ageratum, Alyssum, Amaryllis, Anemone, Anthurium, Aster, Azalea, Begonia, Calceolaria, Campanula, Carnation, Centaurea, Cerastium, Chrysanthemum, Cineraria, Coleus, Cyclamen, Daffodil, Dahlia, Daisy, Delphinium, Dianthus, Dieffenbachia, Freesia, Fuchsia, Gaillardia, Gazania, Geranium, Gerbera, Gladiolus, Gloxinia, Gypsophila, Hedera, Hyacinth, Impatiens, Iris, Kalanchoe, Liatris, Lily, Lobelia, Marigold, Matthiola, Monarda, Nigella, Pansy, Pelargonium, Petunia, Phlox, Poinsettia, Poppy, Primrose, Rhododendron, Rose, Rudbeckia, Salvia, Senecio, Sinningia, Statice, Sweet Pea, Tulip, Verbena, Vinca, Zinnia

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Cuttings, Bare Root or Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Fruiting Vegetables, including: Eggplant, Pepper (e.g., Bell, Sweet, and Hot), Okra, Tomatillo, and Tomato

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Herbs, Spices, and Mints, including: Aniseed, Basil, Caraway, Chive, Dill, Fennel, Lavender, Marjoram, Oregano, Parsley, Rosemary, Sage, Savory, and Thyme

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.

Herbs, Spices, and Mints continued.

Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.
---------------------------------------	--

Hydroponic Crops: Cucumber, Lettuce and Other Leafy Greens, Herbs and Spices, Tomato, and Strawberry

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
	Apply LALSTOP® G46 WG by injection into hydroponic NFT, ebb-and-flow and rockwool applications with each water change or every 5 – 7 days depending on disease pressure. Presoak transplants in the same solution mix.
Injection into Hydroponic Systems	Rate for NFT and ebb-and-flow applications: 0.09 oz. LALSTOP® G46 WG per 1000 plants, 0.002 oz. LALSTOP® G46 WG/100 sq. ft. Rate for rockwool applications: 0.7 – 0.9 oz. LALSTOP® G46 WG per 1000 plants, 0.013 – 0.016 oz. LALSTOP® G46 WG/100 sq. ft.

Leafy Vegetables, Brassica (Cole) Leafy Vegetables, and Other Vegetables, including:

LEAFY VEGETABLES - Arugula, Celery, Endive, Fennel, Lettuce (e.g., Head and Leaf), Parsley, Radicchio, Rhubarb, Spinach, Swiss Chard, and Watercress

BRASSICA (COLE) LEAFY VEGETABLES - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage, Collards, Kale, Kohlrabi, and Mustard Greens

OTHER VEGETABLES - Asparagus

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area

Leafy Vegetables, Brassica (Cole) Leafy Vegetables, and Other Vegetables continued.

Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.
---------------------------------------	--

Peanut

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Pome Fruits, including: Apple, Pear, and Quince

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in one gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Stone Fruits, including: Apricot, Cherry (e.g., Sweet and Tart), Nectarine, Peach, Plum, Prune (Fresh)

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Tobacco

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Trees, Shrub Seedlings, and Shadehouse and Outdoor Nursery Crops, including: Arborvitae, Birch, Deciduous Trees, (e.g., Maple and Oak), Fir, Forest Trees, Fruit Trees, Hemlock, Juniper, Larch, Ornamentals, Pine, Shrubs, Spruce, Vine Crops, and Woody Ornamentals

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Tree Nuts, including: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Coconut, Filbert, Hickory Nut, Macadamia Nut, Pecan, Pistachio, and Walnut

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/ Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.

Tree Nuts continued.

In-Furrow 0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Root, Tuber, and Bulb Vegetables, including: Beet (e.g., Garden and Sugar), Carrot, Cassava, Celeriac, Chicory, Chinese Artichoke, Chive, Dasheen, Garlic, Ginger, Ginseng, Horseradish, Jerusalem Artichoke, Leek, Onion, Parsnip, Potato, Radish, Rutabaga, Salsify, Shallot, Sweet Potato, Turmeric, Turnip, Turnip-Rooted, Chervil, and Yam

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension). Dip bulbs and tubers before planting or storage.
Foliar Spray	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre. For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.
Field Chemigation (Drip or Sprinkler)	0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

Miscellaneous Crops: Avocado, Banana, Coffee, Globe Artichoke, Hops, Mushrooms, Papaya, Plantain, Other Flowering Plants, and Bedding Plants

APPLICATION METHOD	APPLICATION RATE, TIMING, FREQUENCY AND OTHER APPLICATION INSTRUCTIONS
Nursery Soil/Media Drench	0.33 oz. LALSTOP® G46 WG per 2.5 gallons of water (0.1% suspension). Repeat applications every 3 weeks depending on disease pressure.
Growing Media Incorporation	0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Store treated growing media below 77°F and use within 3 weeks after preparation.
Transplant Dip or Drench	0.1 oz. LALSTOP® G46 WG in 1 gallon of water (0.075% suspension).

Miscellaneous Crops continued.

Foliar Spray 0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). Repeat applications every 3 – 6 weeks depending on disease pressure. See FOLIAR APPLICATION INSTRUCTIONS CHART for the amount of LALSTOP® G46 WG/acre.

For fogging (cold fogging only) – Make a 0.1% suspension (0.13 oz. LALSTOP® G46 WG per gallon of water) and apply as a fog at a rate of 0.6 – 1 gallon of suspension per every 10,000 sq. ft. of crop area.

Field Chemigation (Drip or Sprinkler) 0.45 – 0.90 lbs. LALSTOP® G46 WG/acre. Make first application immediately after seeding/transplanting and repeat at a 3- to 6- week interval. Use high rate for high disease pressure and/or high plant density. See CHEMIGATION USE DIRECTIONS.

PREPARING AN AQUEOUS SUSPENSION OF LALSTOP® G46 WG

To prepare an aqueous suspension of LALSTOP® G46 WG, mix product in a small volume of water such as 1/2 gallon (e.g., 0.35 oz. of LALSTOP® G46 WG per 1/2 gallon of water). For best results, let the suspension stand 30 minutes. Agitate sufficiently to evenly disperse product before diluting to final volume. Apply the aqueous suspension as a dip, drench or spray or via drip or sprinkler chemigation or incorporation in potting media. Use the aqueous suspension within 1 day of preparation. Examples of mixing ratios of LALSTOP® G46 WG with water are as follows:

Concentration of LALSTOP® G46 WG Suspension	Ounces (oz.) of LALSTOP® G46 WG	Gallons of Water	Grams (g) of LALSTOP® G46 WG	Liters (L) of Water
0.01%	0.33	25	10	100
0.05%		5		20
0.1%		2.5		10
0.25%		1		4

FOLIAR APPLICATION INSTRUCTIONS CHART

Foliar Spray: 0.33 oz. LALSTOP® G46 WG per 5 gallons of water (0.05% suspension). When using ultra-low volume sprayers, use a maximum pressure of 0.05 psi and a minimum nozzle size of 80 microns.

Foliar Spray Solution Volume	LALSTOP® G46 WG (0.05% suspension)		
(Gallons/Acre)	Ounces (oz.)/Acre	Pounds (lbs.) /Acre	Grams per Acre
10	0.66	0.041	18.71
30	1.98	0.123	56.13
50	3.30	0.206	93.55
75	4.95	0.309	140.33
100	6.60	0.413	187.11
200	13.20	0.825	374.21
400	26.40	1.650	748.43

Foliar Spray Solution Volume	LALSTOP® G46 WG (0.05% suspension)	
(Gallons/1000 Square Feet)	Ounces (oz.)/1000 Square Feet (sq. ft.)	Grams per 1000 Square Feet (sq. ft.)
0.5	0.033	0.94
1.0	0.066	1.87
2.0	0.132	3.74
3.0	0.198	5.61
4.0	0.264	7.48
5.0	0.330	9.36

FOGGING APPLICATION INSTRUCTIONS

Use cold fogging systems only. Use a pressure of 0.036 psi and nozzle size of 100 – 150 microns.

CHEMIGATION USE DIRECTIONS GENERAL REQUIREMENTS

1. Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system

unless the pesticide label-prescribed safety devices for public water systems are in place.

5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPECIFIC REQUIREMENTS FOR CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC REQUIREMENTS FOR SPRINKLER AND DRIP (TRICKLE) CHEMIGATION

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

APPLICATION INSTRUCTIONS FOR ALL TYPES OF CHEMIGATION

1. Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues, may cause product to lose effectiveness or strength.
2. Determine the treatment rates as indicated in the directions for use and make proper dilutions.
3. Make sufficient stock solution of LALSTOP® G46 WG to allow for even injection into irrigation systems during the irrigation cycle. For preparing a stock solution, use clean water with a neutral pH and devoid of salts and chemicals.
4. Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required. Utilize agitation to keep solution in suspension. Apply continuously for the duration of the water application.
5. Apply in sufficient amount of water to move into root zone. Repeat every 3 – 6 weeks for disease control, depending on disease pressure and new plant growth.

6. When applying through sprinkler irrigation systems for soil-borne disease control, use the high rate and time the application for the early stages of crop growth when crop canopy is sparse, which will allow LALSTOP® G46 WG injected water to reach the ground more efficiently.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place. Avoid overheating.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. (For instances where state and local ordinances do allow burning: If burned, stay out of smoke.)

NOTICE TO USER

Lallemand Specialties Inc. / LALLEMAND PLANT CARE warrants only that this product conforms to the product description on this label and is reasonably fit for the purposes set forth in the Directions for Use when used in accordance with them. However, ineffectiveness or other unintended consequences may result because of such factors as the use, storage or handling of the product contrary to the label instructions, all of which are beyond the control of Lallemand Specialties Inc. / LALLEMAND PLANT CARE. To the extent consistent with applicable law, Lallemand Specialties Inc. / LALLEMAND PLANT CARE shall not be liable for indirect or consequential damages resulting from the use, storage or handling of this product. LALLEMAND SPECIALTIES INC. / LALLEMAND PLANT CARE MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Manufactured for:

Danstar Ferment AG / LALLEMAND PLANT CARE
Poststrasse 30
CH-6300 Zug, Switzerland

Marketed by and distributed by:

Lallemand Inc. / LALLEMAND PLANT CARE
1620 rue Prefontaine
Montreal, Quebec
Canada H1W 2N8
1-844-590-7781