



Grosafe®

DANGER

KEEP OUT OF REACH OF CHILDREN

ECOTOXIC

READ LABEL BEFORE USE



FreeFlo™ Copper

FRUIT & VEGIES

**A protectant fungicide
for the control of a range of fungal and bacterial
diseases on fruit and vegetables.**

ACTIVE INGREDIENT: Contains 300 g / kg copper as copper hydroxide
in the form of a water dispersible granule.

200g

GSFFCF&V/420F01

FREEFLO™ COPPER FRUIT & VEGIES 200g

PEEL HERE

MIXING INSTRUCTIONS: Use the measuring cap supplied. The cap supplied has marked lines and if filled to the top line is equal to 11g. Half fill the spray tank with water, slowly add the required amount of FreeFlo Copper while keeping well stirred. Complete top up with water.

WITHHOLDING PERIOD: 1 day

DIRECTIONS FOR USE: Ensure good coverage of foliage.

CROP	DISEASE	RATE g/10 litres	APPLICATION RATE, TIMING
Apples	Black Spot	11	Full cover spray. Apply between silver tip and green tip (max 1cm green tip)
Citrus	Melanose Verrucosia,	4.5 - 9	Apply before and after blossom
	Brown Rot	9 - 11	Apply in Autumn to the skirts of the trees, to a height of 1 metre, to the drip line and 50 cm beyond. Use higher rate if conditions favour disease.
Feijoas	Leaf Spot	7 - 9	Repeat as necessary
Passionfruit	Septoria Spot, Brown Rot, Grease Spot	4.5	Apply at 2 - 4 week intervals from November to March, alternating with maneb/mancozeb. For grease spot 2 week intervals may be required.
Stone Fruit	Leaf Curl, Shot Hole, Bladder Plum	11	Apply when buds are swelling, within one week before buds open, with a second application of 11.5 g/ 10 litres water, 10 - 14 days later.
	Bacterial Blast	31	Apply at leaf fall and repeat every 4 weeks until bud movement, then treat as for Leaf Curl.

HSNO CLASSIFICATIONS:

Acute toxicity, Category 4; Serious eye damage, Category 1; Skin sensitisation, Category 1; Specific target organ toxicity (repeated exposure), Category 2; Aquatic toxicity, acute and chronic, Category 1; Hazardous to terrestrial vertebrates.



Read full booklet for complete list of application rates and food crops.

Registered and distributed by:

GROSAFE CHEMICALS LTD

20 Jean Batten Drive, Mt Maunganui, 3116

PO Box 14 450 Tauranga Central 3141

0800 220002

info@grosafe.co.nz | www.grosafe.co.nz

See container for Batch code

& Date of Manufacture



GSFFCF&V/420/B01



9 421902 986420 >

TOXICITY

DANGER: Causes serious eye damage.

WARNING: Harmful if swallowed. May cause sensitisation from prolonged contact. May cause lung damage from repeated oral exposure at high doses.

ECOTOXICITY

WARNING: Very toxic to aquatic organisms with long-lasting effects. Harmful to terrestrial vertebrates. Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Avoid contamination of any water supply with chemical or empty container.

PERSONAL PROTECTION: Use only outdoors or in well-ventilated area. Avoid breathing dust, mist or spray. When using wear overalls, a face mask, safety glasses and rubber gloves. Do not eat, drink or smoke while using. Wash hands and face before rest breaks and after using.

STORAGE: Store product in original container, tightly closed, in a cool dry place. Keep out of reach of children, and away from foodstuffs.

Shelf life: When stored appropriately this product should show no significant degradation for two years from date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

SPILLS: If spray mixture is spilled absorb, with soil, sand or sawdust, and bury, or collect into bag that can be closed and disposed on in rubbish collection.

PRODUCT DISPOSAL: If possible dispose of by using according to the label, otherwise dispose of in an approved landfill or bury below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

CONTAINER DISPOSAL: Triple rinse and add to the spray tank. Dispose of rinsed container in rubbish collection. Do not use for any other purpose.

SAFETY DATA SHEET: A safety data sheet is available from Grosafe Chemicals Ltd.

FIRST AID: For advice call a doctor or the National Poisons Centre (0800 764 766). Have product label or container at hand. If swallowed, rinse mouth but do not induce vomiting. Get advice if you feel unwell. If on skin wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice. If in eyes flush with plenty of water. Remove contact lenses if present after first few minutes and keep rinsing. Immediately get medical advice. If inhaled remove to fresh air and keep at rest in a position comfortable for breathing. Get medical advice if person feels unwell.

EMERGENCY RESPONSE 0800 CHEMCALL - (0800 243 622)

It is an offence to use this product on animals.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

DIRECTIONS FOR USE: Ensure good coverage of foliage.

CROP	DISEASE	RATE g/10 litres	APPLICATION RATE, TIMING
Asparagus	Stemphylium Leaf Spot	25 - 37.5	Spray at first appearance of disease, after closing up. Repeat at 14 day intervals. Up to six sprays may be required.
Avocado	Anthracnose	7 - 9	Apply monthly from flowering to harvest. Use higher rate and shorten the interval when conditions favour the disease.
Beans	Bacterial Blight	25 - 31	Apply at 7 - 14 day intervals when plants reach 15 cm high.
Bushfruit, Canefruit	Cane Wilt, Cane Spot, Spur Blight, Leaf Spot	7 - 9	Apply from bud burst at 10 - 14 day intervals. Use the higher rate early season or in conditions that favour the disease.
Brassica crops: Broccoli, Brussel Sprouts, Cabbage, Cauliflower	Downy Mildew	7 - 9	Apply at 10 - 14 day intervals following emergence, or after setting out transplants.
Celery	Septoria Leaf Spot	9	After setting out transplants apply at 5 - 7 day intervals, depending on disease severity. EnSpray 99® spray oil may be added at 1 litre per 100 litres of spray mix.
Curcubits	Angular Leaf Spot, Downy Mildew	7 - 13	Apply at 7 day intervals when plants begin to vine.
Grapes	Black Spot, Downy Mildew	6.5 - 13	Apply at 10 - 14 day intervals from bud break to harvest. Adjust rate or stop if injury is observed
Kiwifruit	Leaf Spot, Phomopsis sp. Botryosphaeria	7 - 9	Apply after harvest before leaf fall, after pruning, and two sprays four weeks apart prior to bud burst.
Lettuce	Downy Mildew	7 - 11	Apply at 7 - 10 day intervals following emergence, or after setting out transplants.
Onions	Downy Mildew Bacterial Blight	9	Apply at the first sign of infection and repeat every 10 - 14 days as required.

CROP	DISEASE	RATE g/10 litres	APPLICATION RATE, TIMING
Peas	Downy Mildew Leaf Spot	7 - 9	Apply at 7 - 14 day intervals following emergence.
Pear (including Nashi)	Black Spot	11	Apply one spray between silver tip and green tip (max. 1 cm green tip). Non-bearing trees: Apply cover sprays at 14 - 21 day intervals until vegetative growth ceases. Post Harvest: Apply 19 g/10 litres at 21 days from harvest to leaf fall. Reduce interval in wet weather.
	Fire Blight (<i>Erwinia amylovora</i>)	2.5 - 4.5	Smooth skin varieties: Do not apply between green tip and harvest or russetting may occur. from harvest to leaf fall. Reduce interval in wet weather. Smooth skin varieties: do not apply between green tip and harvest or russetting may occur.
<i>Pinus radiata</i>	Pine Needle Blight (<i>Dothistroma pini</i>)	12.5 - 31	Apply in October. Repeat annually. Repeat applications may be required at 4 weekly intervals on seedlings over growing season.
Potato	Early Blight, Late Blight	13 - 20	Apply every 7 days when plant reaches 15 cm high, until 14 days before harvest.
Strawberries	Leaf Spot	7 - 9	Begin spraying when plants are established and repeat at 7 - 10 day intervals through the season. Stop if signs of crop injury appear.
Tamarillo	Leaf Spot	7	Apply when infection is seen.
Tomato	Early Blight Late Blight Bacterial Speck Bacterial Spot	7 - 13	Apply at 7 - 10 day intervals depending on weather conditions. Increase volume as crop matures: ensure coverage of upper and lower leaf surfaces. Repeat spray after 13 mm rain, or irrigation.

Registered pursuant to the ACVM Act 1979 No. P9918
See www.foodsafety.govt.nz for registration conditions

HSNO Act 1996, Approval code: HSR000739
See www.epa.govt.nz for approval conditions

WARRANTY: This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

FreeFlo™ is a trademark of Grosafe Chemicals Ltd.