



Biocidin® Throat Spray – Scientific Validation of Botanical Ingredients

Bilberry (*Vaccinium myrtillus*)

Medicinal Actions:

Antiinflammatory, antimicrobial, antioxidant, urinary antiseptic.¹⁻³

Scientific Evidence:

Bilberry is rich in phenolic compounds which possess bacteriostatic and antimicrobial properties.⁴ This herb is renowned for its exceptionally high levels of anthocyanins, which are responsible for bilberry's diverse antioxidant effects.^{5,6} As a natural antiinflammatory agents, bilberry polyphenols help reduce lipopolysaccharide (LPS)-induced nuclear factor kappa-beta (NF-K β) activation.⁷

Based on *in vitro* research, bilberry possesses antiadhesive and antimicrobial properties against the respiratory-associated pathogens *Streptococcus pneumoniae* and *Neisseria meningitidis*.⁸ Bilberry extract may also help protect against dental plaque by inhibiting biofilm formation of common oral bacteria including *Streptococcus mutans*, *Fusobacterium nucleatum* and *Actinomyces naeslundii*.⁹

In vitro trials also show that bilberry is able to inhibit the replication of several respiratory viruses including coxsackie virus B1 (CV-B1), human respiratory syncytial virus A2 (HRSV-A2) and influenza virus A/H3N2.¹⁰

Safety Summary:

Considered safe at the recommended dose.¹ No adverse effects expected during pregnancy and breastfeeding.²

Noni (*Morinda citrifolia*)

Medicinal Actions:

Antiinflammatory, antimicrobial, antioxidant.^{3,11}

Scientific Evidence:

To date, over 160 different phytochemical compounds have been identified in the noni plant. The major secondary metabolites include phenolic compounds, organic acids and alkaloids which give rise to noni's potent antioxidant and antiinflammatory properties.¹¹ *In vitro* research has shown that noni is highly effective at inhibiting hydroxyl radicals which are known to cause oxidative damage to proteins, lipids and deoxyribonucleic acid (DNA).¹¹ Noni has also been shown to decrease nitric oxide production. Nitric oxide is a compound produced by macrophages that plays a key role in inflammation and has also been shown to interfere with DNA repair processes. Based on animal trials, the antinociceptive and antiinflammatory effects of noni appear to work in a dose-dependent manner.¹¹ As a natural antiinflammatory agent, noni inhibits LPS-induced activation of a number of chemical mediators including cyclooxygenase (COX)-1 and COX-2, nitric oxide and prostaglandins E2 (PGE2) in a dose-dependent manner.¹²

Research supports the use of noni as both a cellular and humoral immunostimulant given its ability to stimulate the proliferation of T and B lymphocytes.¹³ Based on *in vitro* and *in vivo* animal studies, noni



may help reduce the adverse effects of allergy-mediated respiratory conditions by down regulating the T-helper (Th)-2 cytokine interleukin (IL)-4 and increasing the production of interferon (IFN)- γ .¹⁴

Other *in vitro* trials have also confirmed noni's immunomodulatory properties. In murine effector cells (thymocytes and splenocytes), noni stimulated the release of a number of mediators including tumour

necrosis factor (TNF)- α , IL-1 β , IL-10, IL-12 p70, IFN- γ and nitric oxide and suppressed IL-4. The opposing effects of noni on IL-12 and IL-4 cytokines suggests the herb may enhance the cytotoxicity of natural killer (NK) cells and T lymphocytes.¹⁵

Traditionally noni was used for tuberculosis infections, which has now been substantiated by *in vitro* studies indicating noni is nearly as effective as Rifampicin (with inhibition rates of 89% and 97% respectively).^{16,17}

The active compounds acubin, L-asperuloside and alizarin isolated from noni have demonstrated antibacterial activity against a number of pathogens including *Pseudomonas aeruginosa*, and *Staphylococcus aureus*.¹⁸

Noni has also demonstrated antifungal activity against *Candida albicans* in a dose-dependent manner.¹⁹ Aqueous extracts of noni may also help protect against the conversion of cellular *Candida albicans* into the hyphenated or filamentous form of the yeast. Germ tube formation or hyphenation from blastoconidia by *Candida* species is thought to be a virulence factor in their pathogenesis. Similarly, noni has been shown to inhibit the germination of spores from the filamentous fungi *Aspergillus nidulans*.²⁰

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended.^{3,21} No adverse effects expected during pregnancy and breastfeeding.^{3,22}

Milk Thistle (*Silybum marianum*)

Medicinal Actions:

Antimicrobial, antioxidant.^{1,2}

Scientific Evidence:

Milk thistle is rich in flavanolignans which comprise of silybin A and silybin B (diastereoisomers), silydianin, silychristin and diastereoisomers isosilybin A and isosilybin B. These polyphenolic molecules are collectively referred to as silymarin.¹ Research has shown that the flavanolignans from milk thistle possess potent antibacterial activity against Gram-positive bacteria, but no antimicrobial activity against Gram-negative bacteria or fungi.²³

Silibinin (an equal extract of silybin A and silybin B) has demonstrated antibacterial activity against methicillin-resistant strains of *Staphylococcus aureus*.^{1,24} When silibinin was combined with the antibiotics oxacilin or ampicillin, there was a more than four-fold reduction in the minimum inhibitory bactericidal concentrations. Based on *in vitro* research, silibinin's antimicrobial properties are due to its ability to inhibit ribonucleic acid (RNA) and protein synthesis of Gram-positive organisms (as opposed to attacking the bacterial membrane).²⁴ Silibinin has also demonstrated antioxidant and antiinflammatory properties on LPS-stimulated human monocytes through an inhibitory effect on hydrogen peroxide release and TNF- α production.²⁵

Silibin derivatives of milk thistle have demonstrated significant antiviral activity. In animal studies, silybin derivatives significantly reduced influenza virus A/PR8 replication and its associated mortality

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



in infected mice.²⁶ Silibin appears to work by inhibiting several components induced by influenza A virus infection including oxidative stress, the activation of extracellular signal-regulated kinase (ERK)/p38 mitogen-activated protein kinase (MAPK) and I κ B kinase (IKK) pathways, as well as the expression of autophagic genes (Atg7 and Atg3).²⁶

In the trial by Song and Choi, silymarin demonstrated greater antiviral activity against influenza A/PR/8/34 virus than the pharmaceutical agent Tamiflu® (oseltamivir) (98% vs. 52% respectively).²⁷

Safety Summary:

Contraindicated in persons allergic to plants from the Compositae family.²⁸ No other known warnings, precautions or contraindications.²⁸ No adverse effects expected during pregnancy and breastfeeding.²⁸

Echinacea (*Echinacea purpurea* & *Echinacea angustifolia*)

Medicinal Actions:

Antiinflammatory, antifungal, antiviral, depurative, immune enhancing, immune modulating, lymphatic.^{2,3,28}

Scientific Evidence:

Echinacea possesses both antiinflammatory and immunostimulating properties.²⁹ Alkylamides, one of the active constituents of echinacea are thought to be responsible for the herb's antiinflammatory activity. Emerging research suggests that bacterial lipoproteins and lipopolysaccharides within echinacea (endophytes) represent the major source of immune enhancing properties of this herb.^{30,31} Human cells of the innate immune system detect Braun-type lipoproteins and LPS through Toll-like receptor 2 and 4 pathways, macrophage activation and upregulation of NK cell activity in the body.^{32,33}

Echinacea has demonstrated antimicrobial activity against a number of respiratory pathogens including *Streptococcus pyogenes*, *Staphylococcus aureus*, *Haemophilus influenzae*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Legionella pneumophila* and *Candida albicans*.^{34,35} Based on *in vitro* research, echinacea has both antiinflammatory and direct killing activity against *Streptococcus pyogenes*, *Haemophilus influenzae* and *Legionella pneumophila*. Echinacea has been shown to inhibit the production of the following cytokines: IL-4, IL-6, IL-8, MCP-1, VEGF, and GRO α .³⁶

Because echinacea supports low-running immune systems, it is an effective herb during times of stress which can be associated with increased susceptibility to common colds and respiratory infections. This herb is also safe to use in persons with well-performing immune systems without any risk of over stimulation.³⁷ Based on clinical trials by Goel et al., therapeutic doses of echinacea can improve the systemic immune response in subjects suffering from the common cold. Supplementation with echinacea reduced daily cold and flu symptoms and resulted in a significant and sustained increase in the total number of circulating white blood cells, monocytes, neutrophils and NK cells.^{38,39}

As a natural antiviral agent, echinacea has demonstrated efficacy against a number of respiratory viruses including influenza viruses (A and B strains), respiratory syncytial virus, and rhinovirus.^{34,40-43} Echinacea has also demonstrated potent antiviral activity against strains of highly pathogenic avian influenza viruses including human Victoria (H3N2) and PR8 (H1N1), avian strains KAN-1 (H5N1) and FPV (H7N7), and the pandemic S-OIV (H1N1).⁴⁴ Based on *in vitro* research, possible antiviral mechanisms of action for echinacea include proinflammatory cytokine inhibition (specifically IL-6 and IL-8) and upregulation of inducible nitric oxide synthase.^{41,45-47}

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.

**Safety Summary:**

Contraindicated in persons allergic to plants from the Compositae family.² Exercise caution with patients taking immunosuppressant medications (short term use only).² No other known warnings, precautions or contraindications.⁴⁸ No adverse effects expected during pregnancy and breastfeeding.^{2,28}

Golden Seal (*Hydrastis canadensis*)**Medicinal Actions:**

Antibacterial, antihistamine, antiinflammatory, antimicrobial, mucous membrane trophorestorative.^{1,3,28,49,50}

Scientific Evidence:

Golden seal root contains a number of alkaloids, the most abundant of which is berberine. Both *in vitro* and *in vivo* studies have revealed that berberine possesses antimicrobial activity against bacteria and fungi.^{2,51}

Golden seal leaves are rich in flavonoids (specifically sideroxylin, 8 desmethyl-sideroxylin and 6 desmethyl-sideroxylin).⁵² While the flavonoids from golden seal have no inherent bactericidal properties, they enhance the antimicrobial activity of berberine by acting as efflux pump inhibitors.⁵² It should be noted that one of the major mechanisms by which bacteria become resistant to antibiotics is by over expression of efflux pumps.⁵³

The combined effects of the active constituents in golden seal make this herb a potent antimicrobial agent for a number of respiratory-associated pathogens including methicillin-resistant *Staphylococcus aureus*, and *Neisseria gonorrhoeae* isolates (including antibiotic-resistant strains).^{51,52,54} Berberine has also demonstrated antifungal activity against non-albicans *Candida* species (specifically *Candida krusei*, *Candida kefyr*, *Candida glabrata*, *Candida tropicalis* and *Candida parapsilosis*). When combined with the antimycotic drugs Miconazole or Fluconazole, berberine was able to reduce biofilm formation of pathogenic *Candida albicans*.⁵⁵

One of the key mechanisms by which golden seal inhibits microbial growth is through quenching of the agr quorum sensing (QS) system.^{52,56} QS is a bacterial cell-to-cell communication that controls genes and influences a number of processes including bioluminescence, sporulation, competence, antibiotic production, biofilm formation and virulence factor secretion.⁵⁷

Berberine has been shown to inhibit the growth of several viruses including cytomegalovirus and human H1N1 strains of influenza A viruses. As an antiinflammatory agent, berberine works by inhibiting influenza A-induced production of TNF- α and PGE2 from infected macrophages.⁵⁸

Berberine has also demonstrated antimicrobial activity against the oral pathogens *Streptococcus mutans* and *Fusobacterium nucleatum*.⁵⁹ When compared with sterile saline irrigation, berberine was found to be more effective at eradicating the endodontic pathogens *Fusobacterium nucleatum*, *Enterococcus faecalis* and *Prevotella intermedia*.⁶⁰

Safety Summary:

Exercise caution in patients with kidney disease.³ No other known warnings, precautions or contraindications at the dose recommended.² Contraindicated during pregnancy in therapeutic doses.³ Discouraged during breastfeeding in therapeutic doses.¹



Shiitake Mushroom (*Lentinula edodes*)

Medicinal Actions:

Antibacterial, antifungal, antioxidant, immune modulating.^{3,61}

Scientific Evidence:

Shiitake mushroom contains activated hexose correlated compound (AHCC), which possesses immune-modulating properties.⁶² Another mechanism by which Shiitake enhances immune function is

by increasing the Th-1 cytokine IL-2.⁶³ In human trials, Shiitake mushroom has also been shown to increase the number of circulating B cells in healthy elderly adults.⁶⁴

Shiitake mushroom also contains the polysaccharides lentinan, LEM, KS-2 and eritadenine, which have demonstrated antimicrobial and antiviral activity.⁶⁵ As an antimicrobial agent, lentinan works by activating macrophages and the cytokines TNF- α and IFN- γ with resultant stimulation of T lymphocytes and enhanced immunity.⁶⁵

Based on *in vitro* research, Shiitake mushroom has demonstrated antibacterial activity against a number of respiratory-associated pathogens including: *Streptococcus pyogenes*, *Klebsiella* sp., *Pseudomonas aeruginosa* and *Staphylococcus* sp. (including methicillin-resistant *Staphylococcus aureus* (MRSA)).⁶⁶⁻⁷⁰ Shiitake has also demonstrated antifungal activity against the following organisms: *Candida albicans*, *Candida glabrata*, *Candida parapsilosis*, *Aspergillus fumigatus*, *Aspergillus niger*, and *Scedosporium apiospermum*.⁶⁸⁻⁷⁰

Shiitake may be effective against oral pathogens associated with dental caries and dental plaque accumulation.^{71,72} *In vitro* studies have demonstrated antigingivitis and anticaries activities of Shiitake against the following oral pathogens: *Streptococcus sanguinis*, *Streptococcus mutans*, *Fusobacterium nucleatum*, *Prevotella intermedia*, *Veillonella dispar*, *Neisseria subflava*, *Actinomyces naeslundii* and *Lactobacillus casei*.^{71,73-75} Shiitake protects against gingivitis and caries through a number of mechanisms including inhibition of cell division, prevention of coaggregation and biofilm formation as well as disruption of preexisting biofilms.^{71,73} These effects are explained in part by Shiitake's alpha-glucan content. Shiitake mushroom is rich in alpha-glucanase, an enzyme that has been shown to inhibit sucrose-induced formation of oral biofilms from *Streptococcus mutans* and *Streptococcus sobrinus* species.^{76,77}

Safety Summary:

Considered safe and well tolerated at doses of up to 2.5mg per day for 6 weeks.⁶⁴ Doses of 9 grams per day of liquid AHCC have also been trailed for two weeks in healthy adults with no changes in blood chemistry markers or significant adverse events.⁷⁸ No adverse effects expected during pregnancy and breastfeeding at the dose recommended.⁷⁹

White Willow Bark (*Salix alba*)

Medicinal Actions:

Analgesic, antiinflammatory.¹

Scientific Evidence:

The key active constituents of white willow bark comprise of phenolic glycosides including the salicylates salicortin and salicin.¹ Other important actives include the flavonoids naringenin and isosalipurposide (also known as eriodictyol) and condensed tannins.⁸⁰⁻⁸²

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



Initially it was thought that salicin (converted to salicylic acid *in vivo*) was responsible for the antiinflammatory effects of this herb.⁸² More recent evidence suggests that the potent antiinflammatory effect of white willow bark is derived the sum total of the medicinal actives given the analgesic effects are much broader acting than non-steroidal antiinflammatory drugs (NSAIDs) which contain acetyl salicylic acid.^{1,81} Unlike NSAIDs, white willow bark is not associated with the unwanted side effects of gastric erosion.⁸²

The synergistic effect of the salicylates, flavonoids and tannins found in white willow bark have been shown to inhibit COX-2 and subsequent generation of free radicals by converting arachidonic acid to prostaglandins.⁸³ Other downstream products of COX activity include nitric oxide release and upregulation of proinflammatory cytokines.⁸¹

In vitro studies assessing LPS activated monocytes show that *Salix alba* is able to block nitric oxide release and reduce IL-6 and TNF- α production.^{81,84} While the underlying mechanisms have not been fully elucidated, white willow bark appears to induce monocyte apoptosis and block transcription factor NF-K β activation.^{81,82} This multifactorial effect is thought to be an innate protective mechanism to control local and systemic inflammatory responses in the body.⁸¹

Safety Summary:

Contraindicated in people with salicylate sensitivity.³ No other known warnings, precautions or contraindications at the dose recommended.³ No adverse effects expected during pregnancy and breastfeeding.³

Garlic (*Allium sativum*)

Medicinal Actions:

Antiinflammatory, antimicrobial, antioxidant.¹

Scientific Evidence:

The main active antimicrobial constituent of garlic is allicin (allyl 2-propene thiosulfinate), which is formed when the herb is crushed and alliinase (an enzyme from the bundle sheath cells) combines with the substrate allin. Crushed garlic contains a number of QS compounds such as ajoene and other organosulfides that are produced as degradation products of allicin.^{85,86}

Garlic is an effective virulence blocking agent in *Streptococcus pyogenes* (group A streptococci) infection. *In vitro* research revealed allicin was able to neutralize the haemolytic activity of streptolysin O, a potent cytolytic toxin produced by nearly all strains of *Streptococcus pyogenes*.⁸⁷

Both *in vitro* and *in vivo* studies have identified ajoene as the major QS component of garlic that is able to inhibit the expression of 11 virulence genes controlled by QS – these genes are considered crucial for *Pseudomonas aeruginosa* pathogenicity.^{85,88} In addition to *Pseudomonas aeruginosa*, ajoene has demonstrated antimicrobial activity against the following respiratory-associated pathogens; *Klebsiella pneumoniae*, *Moraxella catarrhalis*, *Staphylococcus aureus*, *Candida albicans*, *Neisseria gonorrhoea*, *Aspergillus niger* and *Paracoccidioides brasiliensis*.^{85,89,90} Garlic has also been shown to be effective against a number of multi-drug-resistant strains of *Pseudomonas aeruginosa*, and *Staphylococcus aureus*.^{91,92}



QS inhibitors such as garlic have demonstrated a synergistic effect when combined with antibiotics. Based on *in vitro* research, the addition of ajoene to a *Pseudomonas* biofilm plus tobramycin killed more than 90% of the bacteria (compared with no effect when tobramycin was tested in isolation).⁸⁵

The thiosulfates in garlic have demonstrated virucidal activity against a number of respiratory viruses including; influenza B virus, parainfluenza virus type 3, coxsackie virus, vaccinia virus, vesicular stomatitis virus, and human rhinovirus type 2.^{93,94}

Garlic has also demonstrated antibacterial activity against oral microbes associated with dental plaque and caries including *Streptococcus mutans*, *Streptococcus sanguis*, *Streptococcus salivarius*, *Pseudomonas aeruginosa*, and *Lactobacillus* spp..^{95,96} Other periodontal pathogens for which garlic has demonstrated antimicrobial activity include *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*. Garlic appears to inhibit the growth of these organisms through antiproteolytic activity and by inhibiting total protease activity.^{97,98}

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended.²⁸ No adverse effects expected during pregnancy and breastfeeding.²⁸

Grape Seed (*Vitis vinifera*)

Medicinal Actions:

Antibacterial, antiinflammatory, antioxidant.³

Scientific Evidence:

Grape seed contains over 95% flavonols, which are predominately comprised of oligomeric proanthocyanins (~82%) and active monomeric proanthocyanins (~12%).⁹⁹

Grape seed extracts have demonstrated antimicrobial activity against a number of respiratory pathogens including *Moraxella catarrhalis*, *Staphylococcus aureus*, *Streptococcus* sp. Group F, *Streptococcus pneumoniae* and *Pseudomonas aeruginosa*.¹⁰⁰ Grape seed extract has also demonstrated antibacterial activity against MRSA strains when assayed through *in vitro* experiments. While the underlying mechanism has not been fully elucidated, grape seed appears to reduce microbial growth by disrupting or breaking down cell wall surfaces.⁹⁹

Another subclass of flavonols found in grape seed identified as procyanidins have demonstrated strong antiviral properties when compared with other botanical agents. These compounds were able to inhibit the replication of influenza A virus at several stages of the life cycle.¹⁰¹ The mechanism by which grape seed exerts antiviral effects relates to the herb's immunostimulatory properties. Grape seed helps upregulate Th-1 immune responses by inducing IFN- γ in mononuclear cells.¹⁰²

Based on *in vitro* research, grape seed has demonstrated antiplaque and antibiofilm activity against oral microbes associated with periodontitis and other acute periodontal diseases. The antimicrobial activity of grape seed has been documented for the following organisms: *Porphyromonas gingivalis*, *Fusobacterium nucleatum*, *Streptococcus mutans*, *Streptococcus sobrinus*, *Lactobacillus rhamnosus* and *Actinomyces viscosus*.¹⁰³ The proanthocyanidins found in grape seed have also demonstrated a protective effect against endotoxin-induced experimental periodontitis (animal research).¹⁰⁴ Other *in vitro* studies indicate grape seed extract supports healthy remineralization of teeth and may be more effective than oral fluoride or calcium glycerophosphate.^{105,106}

Safety Summary: No known warnings, precautions or contraindications at the dose recommended.³ Exercise caution during pregnancy and breastfeeding as safety has not been scientifically established during these times.³

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



Black Walnut (*Juglans nigra*)

Medicinal Actions:

Antimicrobial, depurative.¹

Scientific Evidence:

The main active constituents of black walnut include naphthoquinones (juglone and plumbagin), tannins (ellagic acid) and flavanoids.^{1,28,107} Tannins comprise ~45% of the medicinal actives and exert an astringent effect on mucosal tissue by dehydrating mucosal secretions and protecting the outer layer of mucosal cells.¹⁰⁸

Antimicrobial, antifungal and antiviral actions have also been attributed to the plumbagin and juglone content of black walnut. *In vitro* studies suggest these quinones exert their cytotoxic effects on microorganisms through redox cycling and the oxidation of glutathione.¹⁰⁹

Based on *in vitro* research, black walnut has demonstrated broad spectrum antimicrobial activity against the following respiratory and oral pathogens: *Staphylococcus aureus*, *Micrococcus luteus* and

Streptococcus mutans.^{107,110} Black walnut has also been shown to inhibit biofilm formation and adherence of methicillin-resistant *Staphylococcus aureus*.¹¹¹

Respiratory-associated yeast and fungal organisms susceptible to black walnut comprise of *Candida albicans*, *Trichophyton rubrum*, *Aspergillus niger* and *Rhizopus nigricans*.¹¹⁰

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended.²⁸ Contraindicated during pregnancy and breastfeeding in therapeutic doses.¹¹²

Raspberry (*Rubus idaeus*)

Medicinal Actions:

Antiinflammatory, antimicrobial, antioxidant.^{1,3}

Scientific Evidence:

Raspberry is rich in anthocyanins (mainly cyanidin-3-sophoroside) and phenolic compounds (primarily ellagitannins and ellagic acid). Raspberry also contains quercetin and kaempferol-based flavanols.¹¹³⁻¹¹⁵ Research shows that antioxidant properties of raspberry are attributed to the polyphenolic compounds (specifically ellagitannins) which are highly effective free radical scavengers.^{113,115} Results of an *in vitro* study indicate that raspberry's phenolic compounds are able to protect DNA and decrease lipid peroxidation of lymphocytes in a concentration-dependent manner.¹¹³

The active ellagitannin constituents (sanguin H-6 and lambertianin C) have also demonstrated antiinflammatory properties. Based on *in vitro* research, they inhibit the increase of NF- κ B driven nuclear transcription and resultant TNF- α production in a dose-dependent manner.¹¹⁶ Raspberry actives also been shown to reduce inflammation by inhibiting the release of the enzyme elastase secreted by neutrophils, which is considered a major component of the inflammatory cascade.¹¹⁷

Phenolic compounds also possess antimicrobial properties and have been shown to inhibit the growth of the following respiratory-associated pathogens: *Pseudomonas aeruginosa*, *Staphylococcus aureus*

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



(including methicillin-resistant strains) and *Candida albicans*.^{4,118-121} The mechanism by which phenolic compounds affect the growth of different bacterial species include destabilization of the cytoplasmic membrane, permeabilization of plasma membranes and inhibition of extracellular microbial enzymes. Phenolic compounds also have direct actions on microbial metabolism by depriving bacterial cells of the substrates necessary for growth.¹²² Adherence of bacteria to epithelial surfaces is a prerequisite for colonization of many pathogens, therefore the antimicrobial activity of raspberry may be related in part to the antiadherence activity suggested by Puupponen et al.⁴

In addition to its antimicrobial effects, raspberry has also demonstrated antiviral activity against a number of respiratory pathogens. Based on *in vitro* research, methanol extracts of raspberry have been shown to inhibit the replication of coxsackie virus B1, influenza A virus and influenza virus A/H3N2.¹⁰

Research also shows that raspberry is effective against oral pathogens associated with dental caries and gingivitis. Raspberry has demonstrated antigingivitis and anticaries activities against the following organisms: *Streptococcus sanguinis*, *Streptococcus mutans*, *Fusobacterium nucleatum*, *Prevotella intermedia*, *Veillonella dispar*, *Neisseria subflava*, *Actinomyces naeslundii* and *Lactobacillus casei*.^{73,75} Mechanisms by which raspberry protects against oral pathogens include: preventing bacterial coaggregation and biofilm formation, disruption of preexisting biofilms and decreasing the expression of genes involved in gingival cellular proliferation and differentiation.⁷⁵

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended. Take away from alkaloid-containing medications, metal ion supplements and vitamin B1 (thiamine).²⁸ No adverse effects expected during pregnancy and breastfeeding.²⁸

Fumitory (*Fumaria officinalis*)

Medicinal Actions:

Antimicrobial, antioxidant.¹²³

Scientific Evidence:

The active constituents of fumitory include alkaloids, flavonoids and organic acids.¹ The biological activities of this herb are mainly associated with the isoquinoline alkaloids, in particular protopine.^{124,125} Protopine, has also demonstrated antihistamine effects.¹²⁶ The antioxidant capacity of fumitory is thought to be due to the synergistic effect of the active constituents.¹²³

To date the scientific evaluation of fumitory as an antimicrobial agent is somewhat limited. Results of an *in vitro* study assessing a methanol extract of fumitory have demonstrated significant antimicrobial activity against the following respiratory-associated microorganisms: *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Cladosporium herbarum*.¹²³ The isoquinoline alkaloids from fumitory have also demonstrated significant antifungal activity against *Candida albicans*.¹²⁷

In vitro research also suggests fumitory possesses antiviral properties. Fumitory isoquinolones and their derivatives have demonstrated selective inhibition against parainfluenza type 3 virus.

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended.¹²⁸ Exercise caution during pregnancy and breastfeeding as safety has not been scientifically established during these times.¹²⁶



Gentian (*Gentiana lutea*)

Medicinal Actions:

Antiinflammatory, antimicrobial, antioxidant.^{1,3,129}

Scientific Evidence:

Gentian contains a number of secoiridoid bitter compounds including; gentiopicrin, amarogentin, gentianine, gentianadine, swerosid and swertiamarin. The medicinal constituents also include a group of xanthenes (isovitexin and isogentisin) as well as phenolic acids and phytosterol flavanoids.^{1,129-131} These active constituents give rise to the herb's potent antioxidant, antiinflammatory and antibacterial properties.^{129,131}

The antioxidant and cytoprotective action of gentian is due to the herb's ability to scavenging reactive oxygen species such as hydroxyl radicals, thereby reducing free radical injury to cells.¹³⁰⁻¹³²

Based on *in vitro* trials, the antiinflammatory activity arises from gentian's ability to inhibit myeloperoxidase enzymes which are released during degranulation of neutrophils and monocytes. Myeloperoxidase upregulation is known to contribute to the development of inflammatory and immune-mediated conditions.¹²⁹

The bitter compounds in gentian include gentiopicrin and xanthone isogentisin. These substances possess antimicrobial properties and have been shown to inhibit the growth of a number of respiratory-

associated pathogens including *Streptococcus pyogenes*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Sarcina lutea*, *Micrococcus flavus*, *Micrococcus luteus* and *Candida albicans*.^{133,134}

Safety Summary:

No known warnings, precautions or contraindications at the dose recommended.² No adverse effects expected during pregnancy and breastfeeding.²⁸

Tea Tree (*Melaleuca alternifolia*) Essential Oil

Medicinal Actions:

Antifungal, antimicrobial.¹³⁵⁻¹⁴¹

Scientific Evidence:

Tea tree oil is composed of a complex mixture of compounds, mainly monoterpene and sesquiterpene hydrocarbons and their associated alcohols such as pinene, sabinene, α -terpinene, limonene; p-cymene, 1,8-cineole, γ -terpinene, terpinolene, terpinen-4-ol, α -terpineol, aromadendrene, ledene, δ -cadinene, globulol and viridiforol.^{138,139,141,142} The diverse active constituents give rise to tea tree's antimicrobial activity against a broad range of Gram-positive and Gram-negative bacteria and fungi.^{139,143}

The main antibacterial constituents of tea tree oil are terpinen-4-ol and γ -terpinene. It has been indicated, that terpene compounds found in tea tree oil act on the phospholipid layer of the microbial cell membrane destroying its normal structure and function.¹⁴⁰ Tea tree oil has demonstrated consistent effective antimicrobial activity against methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococcus*, multi-resistant *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*.¹⁴⁴ The antimicrobial effect of tea tree oil against *Pseudomonas aeruginosa* was found to be superior to commercially used antibacterial agents (0.1 % chlorhexidine and 70% ethanol solutions).¹⁴⁴ Tea tree oil has also been shown to decolonize and eradicate biofilms from

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



Staphylococcus aureus (both coagulase-negative and coagulase-positive strains).¹³⁶ Numerous *in vitro* studies have also documented the potent antifungal activity of tea tree oil against *Candida albicans* in both planktonic and biofilm culture.^{135,137}

The antiviral activity of tea tree oil has been primarily attributed to terpinen-4-ol and its ability to prevent the viruses from entering host cells and by inhibiting the 'uncoating' phase of replication.^{145,146} Based on *in vitro* research, tea tree oil has demonstrated an inhibitory effect on influenza A virus and influenza A/PR8 virus subtype H1N1.¹⁴⁵⁻¹⁴⁷ Tree oil has also demonstrated strong antiviral activity against influenza virus A strain NWS/G70C (H1N9) by inactivating the virus and inducing viral decay both in aerosol form as well as in the vaporized phase.¹⁴⁸

Research shows tea tree oil is capable of killing oral pathogens. In a clinical setting, tea tree oil has demonstrated anti-inflammatory properties when applied topically to inflamed gingival tissue in subjects with severe chronic gingivitis.¹⁴⁹ Topical application of tea tree oil in a gel format has also demonstrated efficacy in controlling microbial biofilms associated with salivary *Streptococcus mutans* in orthodontic patients.¹⁵⁰ Based on *in vitro* research, tea tree oil has demonstrated growth-inhibiting and bactericidal effects as well as adhesion-inhibiting effects on a number of oral organisms including *Porphyromonas gingivalis*, *Actinobacillus actinomycetemcomitans*, *Fusobacterium nucleatum*, *Streptococcus mutans*, and *Streptococcus sobrinus*.¹⁴⁰ Tea tree oil has also demonstrated inhibitory activity against the Gram-positive bacillus *Solobacterium moore*, an oral microbe associated with halitosis.¹⁴³

Safety Summary:

Considered safe and well tolerated at the dose recommended. Tea tree oil is generally regarded as non-toxic, and non-irritating.¹⁴¹ Exercise caution during pregnancy and breastfeeding as safety has not been scientifically established during these times.¹⁵¹

Galbanum (*Ferula galbaniflua*) Essential Oil

Medicinal Actions:

Antiinflammatory, antimicrobial, antiseptic.^{141,152}

Scientific Evidence:

Galbanum is composed of mainly monoterpene and sesquiterpene hydrocarons and their associated alcohols including tricyclene, α -pinene, camphene, β -pinene, myrcene, δ -3-carene, limonene, cis-ocimene, trans-ocimene and terpinolene.^{141,153} It is the high concentrations of monoterpenes and sesquiterpenes that give rise to galbanum's antiinflammatory, antimicrobial and antiseptic properties.^{141,154}

Terpenes have been shown to be active against bacteria, fungi and viruses. The mechanism by which terpenes exert their antimicrobial properties involves disruption of the lipophilic compounds of the cellular membranes of pathogens.¹⁵⁵

To date, few scientific studies have been conducted with galbanum oil. Traditionally the herb has been used in the treatment of inflammatory and skin disorders, in wound healing and for ailments of the respiratory, digestive and nervous systems.^{141,154}

Safety Summary:

Galbanum oil is generally regarded non-toxic, non-irritating and non-sensitizing.¹⁴¹ Exercise caution during pregnancy and breastfeeding as safety has not been scientifically established during these times.¹⁵⁶

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.

**Lavender (*Lavandula officinalis*) Essential Oil****Medicinal Actions:**

Antifungal, antiinflammatory, antimicrobial.^{141,157-160}

Scientific Evidence:

Lavender oil contains a complex mixture of aromatic compounds including terpenes and sesquiterpenes (which include linalyl acetate), linalool, caryophyllene, terpinen-4-ol, 2-myrcene, trans-ocimene, borneol, 1,8-cineole, camphor and limonene.^{141,159,161}

This essential oil of lavender has been found to be active against many species of bacteria and fungi. Based on *in vitro* studies, lavender oil has demonstrated antibacterial activity against both methicillin-sensitive and methicillin-resistant strains of *Staphylococcus aureus*.^{159,160,162}

Lavender oil has demonstrated both fungistatic and fungicidal activity against *Candida albicans*. Research also shows lavender is effective against oropharyngeal strains of *Candida albicans*. In the study by D'Auria et al., lavender oil inhibited both germ tube formation and hyphal elongation of *Candida albicans*.¹⁵⁸

Based on *in vitro* experiments, lavender helps protect against LPS-induced inflammation from Gram-negative bacteria. Exposure to LPS in tissues induces an inflammatory reaction which triggers the release of proinflammatory cytokines and subsequent free radical pathology. Research by Huang et al. verified lavender oil was able to inhibit LPS-dependent superoxide anion generation, NF- κ B activation and IL-1 β production.¹⁵⁷

Inhalations of lavender oil have been shown to be effective against allergy-mediated bronchial asthma. Animal studies show inhaled lavender helps reduce allergic inflammation and mucosal hyperplasia. Lavender oil also reduced the Th-2 cytokines IL-4 and IL-5 in bronchoalveolar lavage fluids and lung tissue.¹⁶³

Safety Summary:

Lavender oil is generally regarded as non-toxic, non-irritant and non-sensitizing.¹⁴¹ No adverse effects are expected during pregnancy and breastfeeding at the dose recommended.¹⁶⁴

Oregano (*Origanum vulgare*) Essential Oil**Medicinal Actions:**

Antibacterial, antifungal, antiinflammatory.^{137,165}

Scientific Evidence:

Active constituents of oregano oil include phenolic monoterpenes and sesquiterpenes such as carvacol, thymol, p-cymene, cis-ocimene, caryophyllene and linalool.¹⁴¹ Research shows that antimicrobial activity of oregano oil is predominantly attributed to carvacrol and thymol.^{166,167}

Oregano oil has demonstrated antibacterial activity against the following respiratory pathogens: *Staphylococcus aureus* (including methicillin-resistant strains), *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Mycoplasma mycoides* subsp. *carpi*, *Micrococcus flavus* and *Mycobacterium terrae*.^{165,167-171} The bacteriostatic and bactericidal properties of oregano oil are thought to be due to its effects on cell membrane and membrane components of microorganisms. Based on *in vitro* trials, oregano oil and its constituents impair cell membrane integrity and damage intracellular nucleic acids by

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



NutritionalMedicine NZ



Biocidin
Botanicals

stimulating potassium and phosphate ion leakage and changes to the internal pH of the cell.¹⁶⁵ In a comparative *in vitro* study by Sokovic et al., oregano oil demonstrated greater antibacterial activity than streptomycin against the following pathogens: *Micrococcus flavus*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*.¹⁶⁷

Oregano oil has also demonstrated antifungal activities against *Candida* species.^{137,172} In the study by Pozzatti et al., oregano inhibited the growth and hyphenation of both *Candida albicans* and *Candida dubliniensis*.¹⁷³ The main mechanism of the antifungal activity is associated with the lipophilicity of oregano oil and consequent interaction with the microbial cell membrane. The lipophilic nature of the oil results in changes and losses of enzymatic and structural components of fungal cells such as adenosine triphosphatase, 1,3- β -D-glucan synthases, chitin and mannans, which are also components involved in germ tube formation.¹⁷³ Oregano oil may also exert its antifungal effects through the inhibition of chain respiration through interactions with mitochondrial membranes with resultant decreased energy production and inhibition of germ tube formation and/or cell growth.¹⁷²⁻¹⁷⁴ The antifungal activity of oregano has also been documented for the following species: *Candida tropicalis*, *Candida krusei*, *Candida guilliermondii*, *Candida parapsilosis*, *Cryptococcus neoformans*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*, *Microsporum canis*, *Epidermophyton floccosum*, *Aspergillus niger*, *Aspergillus fumigatus* and *Aspergillus flavus*.¹⁷⁵

Safety Summary:

Generally considered safe and well tolerated at the dose recommended. Active phenolic compounds such as thymol and cavacrol in oregano oil may in some sensitive individuals cause skin and mucus

membrane irritation.¹⁴¹ Exercise caution during pregnancy and breastfeeding as safety has not been scientifically established during these times.¹⁷⁶

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 /00116 21 941 556.

**References**

1. Fisher C. *Materia Medica of Western Herbs*. Nelson, New Zealand: Vitex Medica; 2009.
2. Bone K. *A Clinical Guide to Blending Liquid Herbs: Herbal Formulations for the Individual Patient*. Edinburgh, Scotland: Churchill Livingstone; 2003.
3. Braun L, Cohen M. *Herbs and Natural Supplements an Evidenced Based Guide*. 3rd ed. Chatswood, NSW: Churchill Livingstone; 2010.
4. Puupponen-Pimia R, Nohynek L, Alakomi HL, Oksman-Caldentey KM. The action of berry phenolics against human intestinal pathogens. *BioFactors (Oxford, England)*. 2005;23(4):243-251.
5. Jakesevic M, Xu J, Aaby K, Jeppsson B, Ahrne S, Molin G. Effects of bilberry (*Vaccinium myrtillus*) in combination with lactic acid bacteria on intestinal oxidative stress induced by ischemia-reperfusion in mouse. *Journal of agricultural and food chemistry*. Apr 10 2013;61(14):3468-3478.
6. Jakesevic M, Aaby K, Borge GI, Jeppsson B, Ahrne S, Molin G. Antioxidative protection of dietary bilberry, chokeberry and *Lactobacillus plantarum* HEAL19 in mice subjected to intestinal oxidative stress by ischemia-reperfusion. *BMC complementary and alternative medicine*. 2011;11:8.
7. Karlsen A, Paur I, Bohn SK, et al. Bilberry juice modulates plasma concentration of NF-kappaB related inflammatory markers in subjects at increased risk of CVD. *European journal of nutrition*. Sep 2010;49(6):345-355.
8. Toivanen M, Ryyanen A, Huttunen S, et al. Binding of *Neisseria meningitidis* pili to berry polyphenolic fractions. *Journal of agricultural and food chemistry*. Apr 22 2009;57(8):3120-3127.
9. Riihinen K, Ryyanen A, Toivanen M, Kononen E, Torronen R, Tikkanen-Kaukanen C. Antiaggregation potential of berry fractions against pairs of *Streptococcus mutans* with *Fusobacterium nucleatum* or *Actinomyces naeslundii*. *Phytother Res*. Jan 2011;25(1):81-87.
10. Nikolaeva-Glomb L, Mukova L, Nikolova N, et al. In vitro antiviral activity of a series of wild berry fruit extracts against representatives of Picorna-, Orthomyxo- and Paramyxoviridae. *Natural product communications*. Jan 2014;9(1):51-54.
11. Serafini MR, Santos RC, Guimaraes AG, et al. *Morinda citrifolia* Linn leaf extract possesses antioxidant activities and reduces nociceptive behavior and leukocyte migration. *Journal of medicinal food*. Oct 2011;14(10):1159-1166.
12. Dussossoy E, Brat P, Bony E, et al. Characterization, anti-oxidative and anti-inflammatory effects of Costa Rican noni juice (*Morinda citrifolia* L.). *Journal of ethnopharmacology*. Jan 7 2011;133(1):108-115.
13. Nayak S, Mengi S. Immunostimulant activity of noni (*Morinda citrifolia*) on T and B lymphocytes. *Pharmaceutical biology*. Jul 2010;48(7):724-731.
14. Palu AK, Kim AH, West BJ, Deng S, Jensen J, White L. The effects of *Morinda citrifolia* L. (noni) on the immune system: its molecular mechanisms of action. *Journal of ethnopharmacology*. Feb 12 2008;115(3):502-506.
15. Hirazumi A, Furusawa E. An immunomodulatory polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (noni) with antitumour activity. *Phytother Res*. Aug 1999;13(5):380-387.
16. American Chemical Society. Noni may yield new drugs to fight tuberculosis. Press Release the 2000 International Chemical Congress of Pacific Basis Societies; 2000.
17. Noni plant may help TB. *AIDS patient care and STDs*. Mar 2001;15(3):175.
18. Wang MY, West BJ, Jensen CJ, et al. *Morinda citrifolia* (Noni): a literature review and recent advances in Noni research. *Acta pharmacologica Sinica*. Dec 2002;23(12):1127-1141.
19. Jainkittivong A, Butsarakamruha T, Langlais RP. Antifungal activity of *Morinda citrifolia* fruit extract against *Candida albicans*. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*. Sep 2009;108(3):394-398.
20. Banerjee S, Johnson AD, Csiszar K, Wansley DL, McGeedy P. An extract of *Morinda citrifolia* interferes with the serum-induced formation of filamentous structures in *Candida albicans* and inhibits germination of *Aspergillus nidulans*. *The American journal of Chinese medicine*. 2006;34(3):503-509.
21. West BJ, White LD, Jensen CJ, Palu AK. A double-blind clinical safety study of noni fruit juice. *Pacific health dialog*. Nov 2009;15(2):21-32.
22. Wang MY, Hurn J, Peng L, Nowicki D, Anderson G. A multigeneration reproductive and developmental safety evaluation of authentic *Morinda citrifolia* (noni) juice. *The Journal of toxicological sciences*. Jan 2011;36(1):81-85.
23. Lee DG, Kim HK, Park Y, et al. Gram-positive bacteria specific properties of silybin derived from *Silybum marianum*. *Archives of pharmaceutical research*. Aug 2003;26(8):597-600.
24. Kang HK, Kim HY, Cha JD. Synergistic effects between silybinin and antibiotics on methicillin-resistant *Staphylococcus aureus* isolated from clinical specimens. *Biotechnology journal*. Nov 2011;6(11):1397-1408.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



25. Bannwart CF, Peracoli JC, Nakaira-Takahagi E, Peracoli MT. Inhibitory effect of silibinin on tumour necrosis factor-alpha and hydrogen peroxide production by human monocytes. *Natural product research*. Nov 2010;24(18):1747-1757.
26. Dai JP, Wu LQ, Li R, et al. Identification of 23-(s)-2-amino-3-phenylpropanoyl-silybin as an antiviral agent for influenza A virus infection in vitro and in vivo. *Antimicrobial agents and chemotherapy*. Sep 2013;57(9):4433-4443.
27. Song JH, Choi HJ. Silymarin efficacy against influenza A virus replication. *Phytomedicine*. Jul 15 2011;18(10):832-835.
28. Mills S, Bone K. *The Essential Guide to Herbal Safety*. Philadelphia, U.S.A.: Churchill Livingstone; 2005.
29. Gan XH, Zhang L, Heber D, Bonavida B. Mechanism of activation of human peripheral blood NK cells at the single cell level by Echinacea water soluble extracts: recruitment of lymphocyte-target conjugates and killer cells and activation of programming for lysis. *International immunopharmacology*. Jun 2003;3(6):811-824.
30. Tamta H, Pugh ND, Balachandran P, Moraes R, Sumiyanto J, Pasco DS. Variability in in vitro macrophage activation by commercially diverse bulk echinacea plant material is predominantly due to bacterial lipoproteins and lipopolysaccharides. *Journal of agricultural and food chemistry*. Nov 26 2008;56(22):10552-10556.
31. Pugh ND, Tamta H, Balachandran P, et al. The majority of in vitro macrophage activation exhibited by extracts of some immune enhancing botanicals is due to bacterial lipoproteins and lipopolysaccharides. *International immunopharmacology*. Jul 2008;8(7):1023-1032.
32. Pugh ND, Jackson CR, Pasco DS. Total bacterial load within Echinacea purpurea, determined using a new PCR-based quantification method, is correlated with LPS levels and in vitro macrophage activity. *Planta medica*. Jan 2013;79(1):9-14.
33. Sullivan AM, Laba JG, Moore JA, Lee TD. Echinacea-induced macrophage activation. *Immunopharmacology and immunotoxicology*. 2008;30(3):553-574.
34. Hudson JB. Applications of the phytomedicine Echinacea purpurea (Purple Coneflower) in infectious diseases. *Journal of biomedicine & biotechnology*. 2012;2012:769896.
35. Bany J, Siwicki AK, Zdanowska D, Sokolnicka I, Skopinska-Rozewska E, Kowalczyk M. Echinacea purpurea stimulates cellular immunity and anti-bacterial defence independently of the strain of mice. *Polish journal of veterinary sciences*. 2003;6(3 Suppl):3-5.
36. Sharma SM, Anderson M, Schoop SR, Hudson JB. Bactericidal and anti-inflammatory properties of a standardized Echinacea extract (Echinaforce): dual actions against respiratory bacteria. *Phytomedicine*. Jul 2010;17(8-9):563-568.
37. Schapowal A. Efficacy and safety of Echinaforce(R) in respiratory tract infections. *Wiener medizinische Wochenschrift (1946)*. Feb 2013;163(3-4):102-105.
38. Goel V, Lovlin R, Barton R, et al. Efficacy of a standardized echinacea preparation (Echinilin) for the treatment of the common cold: a randomized, double-blind, placebo-controlled trial. *J Clin Pharm Ther*. Feb 2004;29(1):75-83.
39. Goel V, Lovlin R, Chang C, et al. A proprietary extract from the echinacea plant (Echinacea purpurea) enhances systemic immune response during a common cold. *Phytother Res*. Aug 2005;19(8):689-694.
40. Cech NB, Kandhi V, Davis JM, Hamilton A, Eads D, Laster SM. Echinacea and its alkylamides: effects on the influenza A-induced secretion of cytokines, chemokines, and PGE(2) from RAW 264.7 macrophage-like cells. *International immunopharmacology*. Oct 2010;10(10):1268-1278.
41. Senchina DS, Martin AE, Buss JE, Kohut ML. Effects of Echinacea extracts on macrophage antiviral activities. *Phytother Res*. Jun 2010;24(6):810-816.
42. Ghaemi A, Soleimanjahi H, Gill P, Arefian E, Soudi S, Hassan Z. Echinacea purpurea polysaccharide reduces the latency rate in herpes simplex virus type-1 infections. *Intervirology*. 2009;52(1):29-34.
43. Binns SE, Hudson J, Merali S, Arnason JT. Antiviral activity of characterized extracts from echinacea spp. (Heliantheae: Asteraceae) against herpes simplex virus (HSV-1). *Planta medica*. Sep 2002;68(9):780-783.
44. Pleschka S, Stein M, Schoop R, Hudson JB. Anti-viral properties and mode of action of standardized Echinacea purpurea extract against highly pathogenic avian influenza virus (H5N1, H7N7) and swine-origin H1N1 (S-OIV). *Virology journal*. 2009;6:197.
45. Sharma M, Schoop R, Hudson JB. The efficacy of Echinacea in a 3-D tissue model of human airway epithelium. *Phytother Res*. Jun 2010;24(6):900-904.
46. Sharma M, Anderson SA, Schoop R, Hudson JB. Induction of multiple pro-inflammatory cytokines by respiratory viruses and reversal by standardized Echinacea, a potent antiviral herbal extract. *Antiviral research*. Aug 2009;83(2):165-170.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 /00116 21 941 556.



47. Sharma M, Schoop R, Hudson JB. Echinacea as an antiinflammatory agent: the influence of physiologically relevant parameters. *Phytother Res*. Jun 2009;23(6):863-867.
48. Brendler T, Grünwald J, Jänicke C. Echinaceae angustifoliae herba. In: Heilpflanzen - Herbal Remedies. Medpharm Scientific Publishers; 2003. Accessed March 1st, 2011.
49. Mills S, Bone K. *Principles and Practice of Phytotherapy*. London, England: Churchill Livingstone; 2000.
50. Natural Standard. Bottom Line Monograph: Goldenseal (*Hydrastis canadensis*). <http://www.naturalstandard.net>. Accessed March 1st, 2011.
51. Scazzocchio F, Cometa MF, Tomassini L, Palmery M. Antibacterial activity of *Hydrastis canadensis* extract and its major isolated alkaloids. *Planta medica*. Aug 2001;67(6):561-564.
52. Cech NB, Junio HA, Ackermann LW, Kavanaugh JS, Horswill AR. Quorum quenching and antimicrobial activity of goldenseal (*Hydrastis canadensis*) against methicillin-resistant *Staphylococcus aureus* (MRSA). *Planta medica*. Sep 2012;78(14):1556-1561.
53. Junio HA, Sy-Cordero AA, Etefagh KA, et al. Synergy-directed fractionation of botanical medicines: a case study with goldenseal (*Hydrastis canadensis*). *Journal of natural products*. Jul 22 2011;74(7):1621-1629.
54. Cybulska P, Thakur SD, Foster BC, et al. Extracts of Canadian first nations medicinal plants, used as natural products, inhibit neisseria gonorrhoeae isolates with different antibiotic resistance profiles. *Sexually transmitted diseases*. Jul 2011;38(7):667-671.
55. Wei GX, Xu X, Wu CD. In vitro synergism between berberine and miconazole against planktonic and biofilm *Candida* cultures. *Archives of oral biology*. Jun 2011;56(6):565-572.
56. Wang X, Yao X, Zhu Z, et al. Effect of berberine on *Staphylococcus epidermidis* biofilm formation. *International journal of antimicrobial agents*. Jul 2009;34(1):60-66.
57. Rutherford ST, Bassler BL. Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harbor perspectives in medicine*. Nov 2012;2(11).
58. Cecil CE, Davis JM, Cech NB, Laster SM. Inhibition of H1N1 influenza A virus growth and induction of inflammatory mediators by the isoquinoline alkaloid berberine and extracts of goldenseal (*Hydrastis canadensis*). *International immunopharmacology*. Nov 2011;11(11):1706-1714.
59. Hwang BY, Roberts SK, Chadwick LR, Wu CD, Kinghorn AD. Antimicrobial constituents from goldenseal (the Rhizomes of *Hydrastis canadensis*) against selected oral pathogens. *Planta medica*. Jul 2003;69(7):623-627.
60. Xie Q, Johnson BR, Wenckus CS, Fayad MI, Wu CD. Efficacy of berberine, an antimicrobial plant alkaloid, as an endodontic irrigant against a mixed-culture biofilm in an in vitro tooth model. *Journal of endodontics*. Aug 2012;38(8):1114-1117.
61. Zhong M, Liu B, Wang X, et al. De novo characterization of *Lentinula edodes* C(91-3) transcriptome by deep Solexa sequencing. *Biochemical and biophysical research communications*. Feb 1 2013;431(1):111-115.
62. Love KM, Barnett RE, Holbrook I, et al. A natural immune modulator attenuates stress hormone and catecholamine concentrations in polymicrobial peritonitis. *The journal of trauma and acute care surgery*. Jun 2013;74(6):1411-1418.
63. Kawanishi T, Ikeda-Dantsuji Y, Nagayama A. Effects of two basidiomycete species on interleukin 1 and interleukin 2 production by macrophage and T cell lines. *Immunobiology*. Jul 2010;215(7):516-520.
64. Gaullier JM, Sleboda J, Ofjord ES, et al. Supplementation with a soluble beta-glucan exported from Shiitake medicinal mushroom, *Lentinus edodes* (Berk.) singer mycelium: a crossover, placebo-controlled study in healthy elderly. *International journal of medicinal mushrooms*. 2011;13(4):319-326.
65. Jung BG, Lee JA, Lee BJ. Immunoprophylactic effects of shiitake mushroom (*Lentinula edodes*) against *Bordetella bronchiseptica* in mice. *Journal of microbiology (Seoul, Korea)*. Dec 2012;50(6):1003-1008.
66. Wang Y, Hong Q, Chen Y, Lian X, Xiong Y. Surface properties of polyurethanes modified by bioactive polysaccharide-based polyelectrolyte multilayers. *Colloids and surfaces*. Dec 1 2012;100:77-83.
67. Hearst R, Nelson D, McCollum G, et al. An examination of antibacterial and antifungal properties of constituents of Shiitake (*Lentinula edodes*) and oyster (*Pleurotus ostreatus*) mushrooms. *Complementary therapies in clinical practice*. Feb 2009;15(1):5-7.
68. Rao JR, Smyth TJ, Millar BC, Moore JE. Antimicrobial properties of shiitake mushrooms (*Lentinula edodes*). *International journal of antimicrobial agents*. Jun 2009;33(6):591-592.
69. Kuznetsov O, Mil'kova EV, Sosnina AE, Sotnikova N. [Antimicrobial action of *Lentinus edodes* juice on human microflora]. *Zhurnal mikrobiologii, epidemiologii, i immunobiologii*. Jan-Feb 2005(1):80-82.
70. Hatvani N. Antibacterial effect of the culture fluid of *Lentinus edodes* mycelium grown in submerged liquid culture. *International journal of antimicrobial agents*. Jan 2001;17(1):71-74.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



71. Signoretto C, Marchi A, Bertoncelli A, et al. Effects of mushroom and chicory extracts on the shape, physiology and proteome of the cariogenic bacterium *Streptococcus mutans*. *BMC complementary and alternative medicine*. 2013;13:117.
72. Signoretto C, Burlacchini G, Marchi A, et al. Testing a low molecular mass fraction of a mushroom (*Lentinus edodes*) extract formulated as an oral rinse in a cohort of volunteers. *Journal of biomedicine & biotechnology*. 2011;2011:857987.
73. Spratt DA, Daglia M, Papetti A, et al. Evaluation of plant and fungal extracts for their potential anti-angiogenic and anticancer activity. *Journal of biomedicine & biotechnology*. 2012;2012:510198.
74. Ciric L, Tymon A, Zaura E, et al. In vitro assessment of shiitake mushroom (*Lentinula edodes*) extract for its anti-angiogenic activity. *Journal of biomedicine & biotechnology*. 2011;2011:507908.
75. Canesi L, Borghi C, Stauder M, et al. Effects of fruit and vegetable low molecular mass fractions on gene expression in gingival cells challenged with *Prevotella intermedia* and *Actinomyces naeslundii*. *Journal of biomedicine & biotechnology*. 2011;2011:230630.
76. Yano A, Kikuchi S, Yamashita Y, Sakamoto Y, Nakagawa Y, Yoshida Y. The inhibitory effects of mushroom extracts on sucrose-dependent oral biofilm formation. *Applied microbiology and biotechnology*. Mar 2010;86(2):615-623.
77. Shouji N, Takada K, Fukushima K, Hirasawa M. Anticaries effect of a component from shiitake (an edible mushroom). *Caries research*. Jan-Feb 2000;34(1):94-98.
78. Spierings EL, Fujii H, Sun B, Walshe T. A Phase I study of the safety of the nutritional supplement, active hexose correlated compound, AHCC, in healthy volunteers. *Journal of nutritional science and vitaminology*. Dec 2007;53(6):536-539.
79. Natural Medicines Comprehensive Database. Shiitake Mushroom Monograph. <http://naturaldatabase.therapeuticresearch.com>. Accessed December 22nd, 2013.
80. Poblacka-Olech L, van Norderkassel AM, Vander Heyden Y, Krauze-Baranowska M, Glod D, Baczek T. Chromatographic analysis of salicylic compounds in different species of the genus *Salix*. *Journal of separation science*. Nov 2007;30(17):2958-2966.
81. Bonaterra GA, Heinrich EU, Kelber O, Weiser D, Metz J, Kinscherf R. Anti-inflammatory effects of the willow bark extract STW 33-I (Proaktiv(R)) in LPS-activated human monocytes and differentiated macrophages. *Phytomedicine*. Dec 1 2010;17(14):1106-1113.
82. Bonaterra GA, Kelber O, Weiser D, Metz J, Kinscherf R. In vitro anti-proliferative effects of the willow bark extract STW 33-I. *Arzneimittel-Forschung*. 2010;60(6):330-335.
83. Fiebich BL, Chrubasik S. Effects of an ethanolic salix extract on the release of selected inflammatory mediators in vitro. *Phytomedicine*. Feb 2004;11(2-3):135-138.
84. Drummond EM, Harbourne N, Marete E, et al. Inhibition of proinflammatory biomarkers in THP1 macrophages by polyphenols derived from chamomile, meadowsweet and willow bark. *Phytother Res*. Apr 2012;27(4):588-594.
85. Jakobsen TH, van Gennip M, Phipps RK, et al. Ajoene, a sulfur-rich molecule from garlic, inhibits genes controlled by quorum sensing. *Antimicrobial agents and chemotherapy*. May 2012;56(5):2314-2325.
86. Bhardwaj AK, Vinothkumar K, Rajpara N. Bacterial quorum sensing inhibitors: attractive alternatives for control of infectious pathogens showing multiple drug resistance. *Recent patents on anti-infective drug discovery*. Apr 2013;8(1):68-83.
87. Arzanlou M, Bohlooli S. Inhibition of streptolysin O by allicin - an active component of garlic. *Journal of medical microbiology*. Sep 2010;59(Pt 9):1044-1049.
88. Givskov M. Beyond nutrition: health-promoting foods by quorum-sensing inhibition. *Future microbiology*. Sep 2012;7(9):1025-1028.
89. Naganawa R, Iwata N, Ishikawa K, Fukuda H, Fujino T, Suzuki A. Inhibition of microbial growth by ajoene, a sulfur-containing compound derived from garlic. *Applied and environmental microbiology*. Nov 1996;62(11):4238-4242.
90. Rasheed MU, Thajuddin N. Effect of medicinal plants on *Moraxella catarhalis*. *Asian Pacific journal of tropical medicine*. Feb 2011;4(2):133-136.
91. Karuppiah P, Rajaram S. Antibacterial effect of *Allium sativum* cloves and *Zingiber officinale* rhizomes against multiple-drug resistant clinical pathogens. *Asian Pacific journal of tropical biomedicine*. Aug 2012;2(8):597-601.
92. Gull I, Saeed M, Shaikat H, Aslam SM, Samra ZQ, Athar AM. Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. *Annals of clinical microbiology and antimicrobials*. 2012;11:8.
93. Weber ND, Andersen DO, North JA, Murray BK, Lawson LD, Hughes BG. In vitro virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta medica*. Oct 1992;58(5):417-423.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



94. Tsai Y, Cole LL, Davis LE, Lockwood SJ, Simmons V, Wild GC. Antiviral Properties of Garlic: In vitro Effects on Influenza B, Herpes Simplex and Coxsackie Viruses. *Planta medica*. Oct 1985;51(5):460-461.
95. Chavan SD, Shetty NL, Kanuri M. Comparative evaluation of garlic extract mouthwash and chlorhexidine mouthwash on salivary *Streptococcus mutans* count - an in vitro study. *Oral health & preventive dentistry*. 2010;8(4):369-374.
96. Houshmand B, Mahjour F, Dianat O. Antibacterial effect of different concentrations of garlic (*Allium sativum*) extract on dental plaque bacteria. *Indian J Dent Res*. Jan-Feb 2013;24(1):71-75.
97. Shetty S, Thomas B, Shetty V, Bhandary R, Shetty RM. An in-vitro evaluation of the efficacy of garlic extract as an antimicrobial agent on periodontal pathogens: A microbiological study. *Ayu*. Oct 2013;34(4):445-451.
98. Bakri IM, Douglas CW. Inhibitory effect of garlic extract on oral bacteria. *Archives of oral biology*. Jul 2005;50(7):645-651.
99. Su X, Howell AB, D'Souza DH. Antibacterial effects of plant-derived extracts on methicillin-resistant *Staphylococcus aureus*. *Foodborne pathogens and disease*. Jun 2012;9(6):573-578.
100. Cueva C, Mingo S, Munoz-Gonzalez I, et al. Antibacterial activity of wine phenolic compounds and oenological extracts against potential respiratory pathogens. *Letters in applied microbiology*. Jun 2012;54(6):557-563.
101. Dai J, Wang G, Li W, et al. High-throughput screening for anti-influenza A virus drugs and study of the mechanism of procyanidin on influenza A virus-induced autophagy. *Journal of biomolecular screening*. Jun 2012;17(5):605-617.
102. Nair N, Mahajan S, Chawda R, Kandaswami C, Shanahan TC, Schwartz SA. Grape seed extract activates Th1 cells in vitro. *Clinical and diagnostic laboratory immunology*. Mar 2002;9(2):470-476.
103. Furiga Chusseau A, Roques C, Badet C. Preventive effects of an original combination of grape seed polyphenols with amine fluoride on dental biofilm formation and oxidative damage by oral bacteria. *Journal of applied microbiology*. Nov 19 2013.
104. Govindaraj J, Emmadi P, Deepalakshmi, Rajaram V, Prakash G, Puvanakrishnan R. Protective effect of proanthocyanidins on endotoxin induced experimental periodontitis in rats. *Indian journal of experimental biology*. Feb 2010;48(2):133-142.
105. Xie Q, Bedran-Russo AK, Wu CD. In vitro remineralization effects of grape seed extract on artificial root caries. *Journal of dentistry*. Nov 2008;36(11):900-906.
106. Benjamin S, Sharma R, Thomas SS, Nainan MT. Grape seed extract as a potential remineralizing agent: a comparative in vitro study. *The journal of contemporary dental practice*. Jul-Aug 2012;13(4):425-430.
107. Amarowicz R, Dykes GA, Pegg RB. Antibacterial activity of tannin constituents from *Phaseolus vulgaris*, *Fagopyrum esculentum*, *Corylus avellana* and *Juglans nigra*. *Fitoterapia*. Apr 2008;79(3):217-219.
108. McGuffin M, Hobbs C, Upton R, Goldberg A eds. *American Herbal Products Association's Botanical Safety Handbook*. Boca Raton, FL: CRC Press, LLC; 1997.
109. Inbaraj JJ, Chignell CF. Cytotoxic action of juglone and plumbagin: a mechanistic study using HaCaT keratinocytes. *Chemical research in toxicology*. Jan 2004;17(1):55-62.
110. Heisey RM, Gorham BK. Antimicrobial effects of plant extracts on *Streptococcus mutans*, *Candida albicans*, *Trichophyton rubrum* and other micro-organisms. *Letters in applied microbiology*. 1992;14:136-139.
111. Quave CL, Plano LR, Pantuso T, Bennett BC. Effects of extracts from Italian medicinal plants on planktonic growth, biofilm formation and adherence of methicillin-resistant *Staphylococcus aureus*. *Journal of ethnopharmacology*. Aug 13 2008;118(3):418-428.
112. Natural Medicines Comprehensive Database. Black Walnut Monograph. <http://naturaldatabase.therapeuticresearch.com>. Accessed July 17th, 2012.
113. Godevac D, Tesevic V, Vajs V, Milosavljevic S, Stankovic M. Antioxidant properties of raspberry seed extracts on micronucleus distribution in peripheral blood lymphocytes. *Food Chem Toxicol*. Nov 2009;47(11):2853-2859.
114. Mullen W, McGinn J, Lean ME, et al. Ellagitannins, flavonoids, and other phenolics in red raspberries and their contribution to antioxidant capacity and vasorelaxation properties. *Journal of agricultural and food chemistry*. Aug 28 2002;50(18):5191-5196.
115. Kahkonen M, Kylli P, Ollilainen V, Salminen JP, Heinonen M. Antioxidant activity of isolated ellagitannins from red raspberries and cloudberries. *Journal of agricultural and food chemistry*. Feb 8 2012;60(5):1167-1174.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 /00116 21 941 556.



116. Sangiovanni E, Vrhovsek U, Rossoni G, et al. Ellagitannins from Rubus berries for the control of gastric inflammation: in vitro and in vivo studies. *PLoS ONE*. 2013;8(8):e71762.
117. Johansson S, Goransson U, Luijendijk T, Backlund A, Claeson P, Bohlin L. A neutrophil multitarget functional bioassay to detect anti-inflammatory natural products. *Journal of natural products*. Jan 2002;65(1):32-41.
118. Puupponen-Pimia R, Nohynek L, Hartmann-Schmidlin S, et al. Berry phenolics selectively inhibit the growth of intestinal pathogens. *Journal of applied microbiology*. 2005;98(4):991-1000.
119. Nohynek LJ, Alakomi HL, Kahkonen MP, et al. Berry phenolics: antimicrobial properties and mechanisms of action against severe human pathogens. *Nutrition and cancer*. 2006;54(1):18-32.
120. Nile SH, Park SW. Edible berries: Review on bioactive components and their effect on human health. *Nutrition (Burbank, Los Angeles County, Calif)*. Sep 3 2013.
121. Cavanagh HM, Hipwell M, Wilkinson JM. Antibacterial activity of berry fruits used for culinary purposes. *Journal of medicinal food*. Spring 2003;6(1):57-61.
122. Puupponen-Pimia R, Nohynek L, Alakomi HL, Oksman-Caldentey KM. Bioactive berry compounds—novel tools against human pathogens. *Applied microbiology and biotechnology*. Apr 2005;67(1):8-18.
123. Sengul M, Yildiz H, Gungor N, Cetin B, Eser Z, Ercisli S. Total phenolic content, antioxidant and antimicrobial activities of some medicinal plants. *Pakistan journal of pharmaceutical sciences*. Jan 2009;22(1):102-106.
124. Hentschel C, Dressler S, Hahn EG. [Fumaria officinalis (fumitory)—clinical applications]. *Fortschritte der Medizin*. Jul 10 1995;113(19):291-292.
125. Rakotondramasy-Rabesiaka L, Havet JL, C. Porte, Faucet H. Solid-liquid extraction of protopine from Fumaria officinalis L.—Kinetic modelling of influential parameters. *Industrial Crops and Products*. 2009;29(2-3):516-523.
126. Newall CA, Anderson LA, Philpson JD. *Herbal Medicine: A Guide for Healthcare Professionals*. London, UK: The Pharmaceutical Press; 1996.
127. Orhana I, Ozcelik B, Karaoglu T, Sener B. Antiviral and antimicrobial profiles of selected isoquinoline alkaloids from Fumaria and Corydalis species. *Zeitschrift fur Naturforschung*. Jan-Feb 2007;62(1-2):19-26.
128. Brinkhaus B, Hentschel C, Von Keudell C, et al. Herbal medicine with curcuma and fumitory in the treatment of irritable bowel syndrome: a randomized, placebo-controlled, double-blind clinical trial. *Scandinavian journal of gastroenterology*. Aug 2005;40(8):936-943.
129. Nastasijevic B, Lazarevic-Pasti T, Dimitrijevic-Brankovic S, et al. Inhibition of myeloperoxidase and antioxidative activity of Gentiana lutea extracts. *Journal of pharmaceutical and biomedical analysis*. Jul 2012;66:191-196.
130. Calliste CA, Trouillas P, Allais DP, Simon A, Duroux JL. Free radical scavenging activities measured by electron spin resonance spectroscopy and B16 cell antiproliferative behaviors of seven plants. *Journal of agricultural and food chemistry*. Jul 2001;49(7):3321-3327.
131. Singh A. Phytochemicals of gentianaceae: a review of pharmacological properties. *nt. J. Pharm. Sci. Nanotechnol*. 2008;1:33-36.
132. Kusar A, Zupancic A, Sentjurc M, Baricevic D. Free radical scavenging activities of yellow gentian (*Gentiana lutea* L.) measured by electron spin resonance. *Human & experimental toxicology*. Oct 2006;25(10):599-604.
133. Weckesser S, Engel K, Simon-Haarhaus B, Wittmer A, Pelz K, Schempp CM. Screening of plant extracts for antimicrobial activity against bacteria and yeasts with dermatological relevance. *Phytomedicine*. Aug 2007;14(7-8):508-516.
134. Savikin K, Menkovic N, Zdunic G, Stevic T, Radanovic D, Jankovic T. Antimicrobial activity of Gentiana lutea L. extracts. *Zeitschrift fur Naturforschung*. May-Jun 2009;64(5-6):339-342.
135. Ramage G, Milligan S, Lappin DF, et al. Antifungal, cytotoxic, and immunomodulatory properties of tea tree oil and its derivative components: potential role in management of oral candidosis in cancer patients. *Frontiers in microbiology*. 2012;3:220.
136. Brady A, Loughlin R, Gilpin D, Kearney P, Tunney M. In vitro activity of tea-tree oil against clinical skin isolates of methicillin-resistant and -sensitive Staphylococcus aureus and coagulase-negative staphylococci growing planktonically and as biofilms. *Journal of medical microbiology*. Oct 2006;55(Pt 10):1375-1380.
137. Irkin R, Korukluoglu M. Growth inhibition of pathogenic bacteria and some yeasts by selected essential oils and survival of *L. monocytogenes* and *C. albicans* in apple-carrot juice. *Foodborne pathogens and disease*. Apr 2009;6(3):387-394.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 / 00116 21 941 556.



138. Kurekci C, Padmanabha J, Bishop-Hurley SL, Hassan E, Al Jassim RA, McSweeney CS. Antimicrobial activity of essential oils and five terpenoid compounds against *Campylobacter jejuni* in pure and mixed culture experiments. *International journal of food microbiology*. Sep 16 2013;166(3):450-457.
139. Flores FC, de Lima JA, Ribeiro RF, et al. Antifungal activity of nanocapsule suspensions containing tea tree oil on the growth of *Trichophyton rubrum*. *Mycopathologia*. Apr 2013;175(3-4):281-286.
140. Takarada K, Kimizuka R, Takahashi N, Honma K, Okuda K, Kato T. A comparison of the antibacterial efficacies of essential oils against oral pathogens. *Oral microbiology and immunology*. Feb 2004;19(1):61-64.
141. Battaglia S. *The Complete Guide to Aromatherapy*. Brisbane, Australia: The perfect potion (Aust) Pty Ltd; 2003.
142. Thomsen NA, Hammer KA, Riley TV, Van Belkum A, Carson CF. Effect of habituation to tea tree (*Melaleuca alternifolia*) oil on the subsequent susceptibility of *Staphylococcus* spp. to antimicrobials, triclosan, tea tree oil, terpinen-4-ol and carvacrol. *International journal of antimicrobial agents*. Apr 2013;41(4):343-351.
143. Forrer M, Kulik EM, Filippi A, Waltimo T. The antimicrobial activity of alpha-bisabolol and tea tree oil against *Solobacterium moorei*, a Gram-positive bacterium associated with halitosis. *Archives of oral biology*. Jan 2013;58(1):10-16.
144. Warnke PH, Lott AJ, Sherry E, Wiltfang J, Podschun R. The ongoing battle against multi-resistant strains: in-vitro inhibition of hospital-acquired MRSA, VRE, *Pseudomonas*, ESBL *E. coli* and *Klebsiella* species in the presence of plant-derived antiseptic oils. *J Craniomaxillofac Surg*. Jun 2013;41(4):321-326.
145. Garozzo A, Timpanaro R, Stivala A, Bisignano G, Castro A. Activity of *Melaleuca alternifolia* (tea tree) oil on Influenza virus A/PR/8: study on the mechanism of action. *Antiviral research*. Jan 2011;89(1):83-88.
146. Li X, Duan S, Chu C, et al. *Melaleuca alternifolia* concentrate inhibits in vitro entry of influenza virus into host cells. *Molecules (Basel, Switzerland)*. 2013;18(8):9550-9566.
147. Garozzo A, Timpanaro R, Bisignano B, Furneri PM, Bisignano G, Castro A. In vitro antiviral activity of *Melaleuca alternifolia* essential oil. *Letters in applied microbiology*. Dec 2009;49(6):806-808.
148. Usachev EV, Pyankov OV, Usachev OU, Agranovski IE. Antiviral activity of tea tree and eucalyptus oil aerosol and vapour. *Journal of Aerosol Science*. 2013;59(2013):22-30.
149. Soukoulis S, Hirsch R. The effects of a tea tree oil-containing gel on plaque and chronic gingivitis. *Australian dental journal*. Jun 2004;49(2):78-83.
150. Santamaria M, Jr., Petermann KD, Vedovello SA, Degan V, Lucato A, Franzini CM. Antimicrobial effect of *Melaleuca alternifolia* dental gel in orthodontic patients. *American journal of orthodontics and dentofacial orthopedics : official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*. Feb 2014;145(2):198-202.
151. Natural Standard. Professional Monograph: Tea tree oil (*Melaleuca alternifolia*). <http://www.naturalstandard.net>. Accessed December 24th, 2013.
152. Brendler T, Gruenwald J, Jaenicke C, Galbanum. In: Heilpflanzen – Herbal Remedies. Medpharm Scientific Publishers; 2003. Accessed December 25th, 2013.
153. Kanani MR, Rahiminejad MR, Sonboli A, Mozaffarian V, Kazempour Osaloo S, Nejad Ebrahimi S. Chemotaxonomic significance of the essential oils of 18 *Ferula* species (Apiaceae) from Iran. *Chemistry & biodiversity*. Mar 2011;8(3):503-517.
154. Nazari ZE, Iranshahi M. Biologically active sesquiterpene coumarins from *Ferula* species. *Phytother Res*. Mar 2011;25(3):315-323.
155. Cowan MM. Plant products as antimicrobial agents. *Clinical microbiology reviews*. Oct 1999;12(4):564-582.
156. Natural Medicines Comprehensive Database. Galbanum Monograph. <http://naturaldatabase.therapeuticresearch.com>. Accessed December 25th, 2013.
157. Huang MY, Liao MH, Wang YK, Huang YS, Wen HC. Effect of lavender essential oil on LPS-stimulated inflammation. *The American journal of Chinese medicine*. 2012;40(4):845-859.
158. D'Auria FD, Tecca M, Strippoli V, Salvatore G, Battinelli L, Mazzanti G. Antifungal activity of *Lavandula angustifolia* essential oil against *Candida albicans* yeast and mycelial form. *Medical mycology*. Aug 2005;43(5):391-396.
159. Roller S, Ernest N, Buckle J. The antimicrobial activity of high-necrodane and other lavender oils on methicillin-sensitive and -resistant *Staphylococcus aureus* (MSSA and MRSA). *Journal of alternative and complementary medicine (New York, N.Y.)*. Mar 2009;15(3):275-279.
160. de Rapper S, Kamatou G, Viljoen A, van Vuuren S. The In Vitro Antimicrobial Activity of *Lavandula angustifolia* Essential Oil in Combination with Other Aroma-Therapeutic Oils. *Evid Based Complement Alternat Med*. 2013;2013:852049.

AUTHORS: Corene Humphreys, ND, BHSc, Dip Med Herb, Dip Hom, QTA (Microbiology). | Anita Kasa, BNat, Dip Herb Med, Dip Nutr, Dip Yoga.

NUTRITIONALMEDICINENZ.COM / BIOCIDIN.NZ.

PHONE: NZ: 09 425 4250 / 021 941 556. AUST: 001169 425 4250 /00116 21 941 556.



161. Evandri MG, Battinelli L, Daniele C, Mastrangelo S, Bolle P, Mazzanti G. The antimutagenic activity of *Lavandula angustifolia* (lavender) essential oil in the bacterial reverse mutation assay. *Food Chem Toxicol.* Sep 2005;43(9):1381-1387.
162. Nelson RR. In-vitro activities of five plant essential oils against methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus faecium*. *The Journal of antimicrobial chemotherapy.* Aug 1997;40(2):305-306.
163. Ueno-lio T, Shibakura M, Yokota K, et al. Lavender essential oil inhalation suppresses allergic airway inflammation and mucous cell hyperplasia in a murine model of asthma. *Life sciences.* Jul 17 2014;108(2):109-115.
164. Brendler T, Gruenwald J, Jaenicke C. Comm. E Monograph: *Lavandulae flos* In: Heilpflanzen - Herbal Remedies. In: Publishers MS, ed2003. Accessed December 25th, 2013.
165. Lambert RJ, Skandamis PN, Coote PJ, Nychas GJ. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of applied microbiology.* Sep 2001;91(3):453-462.
166. Gilling DH, Kitajima M, Torrey JR, Bright KR. Antiviral efficacy and mechanisms of action of oregano essential oil and its primary component carvacrol against murine norovirus. *Journal of applied microbiology.* May 2014;116(5):1149-1163.
167. Sokovic M, Glamoclija J, Marin PD, Brkic D, van Griensven LJ. Antibacterial effects of the essential oils of commonly consumed medicinal herbs using an in vitro model. *Molecules (Basel, Switzerland).* Nov 2010;15(11):7532-7546.
168. Friedman M, Henika PR, Levin CE. Bactericidal activities of health-promoting, food-derived powders against the foodborne pathogens *Escherichia coli*, *Listeria monocytogenes*, *Salmonella enterica*, and *Staphylococcus aureus*. *Journal of food science.* Feb 2013;78(2):M270-275.
169. Arjoon AV, Saylor CV, May M. In Vitro efficacy of antimicrobial extracts against the atypical ruminant pathogen *Mycoplasma mycoides subsp. capri*. *BMC complementary and alternative medicine.* 2012;12:169.
170. Eng W, Norman R. Development of an oregano-based ointment with anti-microbial activity including activity against methicillin-resistant *Staphylococcus aureus*. *J Drugs Dermatol.* Apr 2010;9(4):377-380.
171. Preuss HG, Echard B, Enig M, Brook I, Elliott TB. Minimum inhibitory concentrations of herbal essential oils and monolaurin for gram-positive and gram-negative bacteria. *Molecular and cellular biochemistry.* Apr 2005;272(1-2):29-34.
172. Pozzatti P, Loreto ES, Lopes PG, Athayde ML, Santurio JM, Alves SH. Comparison of the susceptibilities of clinical isolates of *Candida albicans* and *Candida dubliniensis* to essential oils. *Mycoses.* Jan 2010;53(1):12-15.
173. Pozzatti P, Loreto ES, Nunes Mario DA, Rossato L, Santurio JM, Aleves SH. Activities of essential oils in the inhibition of *Candida albicans* and *Candida dubliniensis* germ tube formation. *Journal de Mycologie Médicale.* 2010;20(3):185-189.
174. de Loreto ES, Pozzatti P, Alves Scheid L, Santurio D, Morais Santurio J, Alves SH. Differentiation of *Candida dubliniensis* from *Candida albicans* on rosemary extract agar and oregano extract agar. *Journal of clinical laboratory analysis.* 2008;22(3):172-177.
175. Vale-Silva L, Silva MJ, Oliveira D, et al. Correlation of the chemical composition of essential oils from *Origanum vulgare subsp. virens* with their in vitro activity against pathogenic yeasts and filamentous fungi. *Journal of medical microbiology.* Feb 2012;61(Pt 2):252-260.
176. Natural Medicines Comprehensive Database. Oregano Monograph. 2003; <http://naturaldatabase.therapeuticresearch.com>. Accessed December 25th, 2013.