

FOR ANIMAL USE ONLY

QUANTEL®

Tablets

Reg.no.G2717 (Act 36 /1947)

INDICATIONS

Controls roundworm, hookworm, whipworm and tapeworm in dogs and cats.

COMPOSITION

Praziquantel (isoquinoline) 50 mg/tablet

Fenbendazole (benzimidazole) 500 mg/tablet

STORAGE

Store below 25°C.

WARNINGS

- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Do not exceed the stated dose when treating pregnant bitches and queens.

Safe for use in lactating animals.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED

Quantel tablets are administered orally either directly or mixed with a portion of meat or sausage or mixed with the daily diet. Dietary measures or fasting are not necessary.

Routine Treatment of Adult Dogs

Quantel should be administered as a single treatment at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg body mass (equivalent to 1 tablet/10 kg body mass):

Small dogs and puppies over 6 months of age

0.5 to 2.5 kg body mass	¼ tablet
2.6 to 5 kg body mass	½ tablet
6 to 10 kg body mass	1 tablet

Medium dogs

11 to 15 kg body mass	1½ tablets
16 to 20 kg body mass	2 tablets
21 to 25 kg body mass	2½ tablets
26 to 30 kg body mass	3 tablets

Large dogs

31 to 35 kg body mass	3½ tablets
36 to 40 kg body mass	4 tablets

Routine Treatment of Adult Cats

Quantel should be administered as a single treatment at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg body mass (equivalent to ½ tablet/5 kg):

0.5 to 2.5 kg body mass	¼ tablet
2.6 to 5 kg body mass	½ tablet

For routine control adult dogs and cats should be treated once every 3 months.

Weaned Puppies and Kittens under 6 months of age

Quantel should be administered as a single treatment at a dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg body mass (equivalent to ½ tablet/5 kg). Treatment should be repeated for 3 consecutive days.

For the control of *Toxacara*, it is important to deworm young puppies regularly with **Quantel** at 2 weekly intervals from the age of 2 weeks for puppies less than 12 weeks of age. It is then recommended that **Quantel** be administered at intervals of 3 months.

Nursing bitches should be treated at the same time and as frequently as the puppies up to 12 weeks of age. Thereafter the adult worming regime of once every three months is recommended.

Increased Dosing for Specific Infections

For the treatment of clinical worm infestations in adult dogs administer **Quantel** at the dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg body mass daily for 2 consecutive days (equivalent to 1 tablet/10 kg for 2 days).

For the treatment of clinical worm infestations in adult cats administer **Quantel** at the dose rate of 5 mg praziquantel and 50 mg fenbendazole per kg body mass daily for 3 consecutive days (equivalent to ½ tablet/5 kg for 3 days).

EFFICACY

Highly effective against ascarids (*Toxacara canis*, *Toxascaris leonina*), hookworm (*Ancylostoma caninum*), tapeworms (*Dipylidium caninum*, *Echinococcus* spp, *Taenia hydatagena*). Effective against whipworm (*Trichuris vulpis*).

PRESENTATION

Plastic containers containing 100 tablets.

REGISTRATION HOLDER

The Chanelle Group
IDA Business Park
Dublin Road, Loughrea,
Co. Galway, Ireland.

MARKETED BY

MSD Animal Health
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za