



MATERIAŁ: folia

WYMIARY: 80 x 75 mm

KOLORYSTYKA: CMYK

CUT

SIZE


FRONT




75 mm

80 mm

BACK/ BAZA



BOOKLET AWERS



**GENERAL INSTRUCTIONS:**

**ATTRATHOR TARGETED INSECTICIDE** is a targeted crack and crevice treatment for cockroaches and ants that can also be used for the control of other crawling pests. It can be used in warehouses, restaurants, food manufacturing, processing and storage plants, supermarkets, hotels, hospitals, domestic and other similar situations.

**MIXING**

Invert the ATTRATHOR container several times to ensure thorough mixing of concentrate. Half fill the spray tank with water and then add the required quantity of ATTRATHOR. Stir and then top up the spray tank to the required volume. The use of this product in a tank mix with other insecticides is not recommended as the behaviour and efficacy of the product may be affected. Ensure equipment is clean.

**INSECTICIDE RESISTANCE WARNING:**

**GROUP 2B INSECTICIDE**

For insecticide resistance management ATTRATHOR Targeted Insecticide is a Group 2B Insecticide. Some naturally occurring insect biotypes resistant to ATTRATHOR Targeted Insecticide and other Group 2B Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if ATTRATHOR Targeted Insecticide or other Group 2B insecticides are used repeatedly. The effectiveness of ATTRATHOR Targeted Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Ensysstex Australasia Pty Ltd accepts no liability for any losses that may result from the failure of ATTRATHOR Targeted Insecticide to control resistant insects.

ATTRATHOR Targeted Insecticide may be subject to specific resistance management strategies. For further information contact Ensysstex Australasia Pty Ltd.

**PRECAUTIONS**

Avoid contact with food, food utensils, or places where food is prepared or stored. Before use, remove or cover all exposed foodstuffs. Cover all dishes and utensils, and places where food is prepared or stored. Avoid application around edible plants. Do not apply using equipment carried on the back of the user.

**RE-ENTRY PERIOD**

Do not enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing). Clothing must be laundered after each day's use. Do not allow public re-entry into treated areas until the spray has dried.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Highly toxic to fish and aquatic organisms. Do NOT apply to areas where surface water is present. Rinse waters and runoff from treated areas MUST be prevented from entering drains or waterways. Do NOT apply if heavy rains are expected to occur within 48 hours of application. Do NOT contaminate streams, rivers or waterways with the chemical or used containers. Dangerous to bees.

**PROTECTION OF PETS AND LIVESTOCK**

Before spraying remove animals and pets from the areas to be treated. Cover or remove any open food and water containers. Cover or remove (as applicable) fish ponds, aquariums, etc. before spraying.

**ATTRATHOR TARGETED INSECTICIDE**

BOOKLET AWERS

**DIRECTIONS FOR USE:**

**RESTRAINT. DO NOT** apply to excessively wet soils, immediately after or during heavy rain, to avoid runoff of the chemical.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Internal & external areas & surrounds of Domestic, Commercial, Public and Industrial buildings, facilities & structures (including but not limited to food processing establishments, shops, factories, farm buildings, ships, trains, aeroplanes, offices, schools, storerooms, hospitals, barracks, private houses, and external areas associated with such establishments.)	<b>Cockroaches</b>	50 mL per 5 L water	Inspect premises thoroughly and locate all insect harbourages. Apply directly into harbourages such as cracks and crevices away from where it is likely to be removed by cleaning programs. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control. Apply behind sinks, stoves, and pipes paying particular attention to all cracks and crevices which might contain insects. Repeat treatments when pests begin to reappear.
	<b>Ants</b>		Apply prepared spray to indoor and outdoor surfaces where ants are tracking. Internal harbourage sites include (but are not restricted to) areas such as walls, behind and under sinks, under furniture and indoor plants. External harbourage sites include (but are not restricted to) areas such as building exterior, eaves, walls, fences, garages, sheds, gazebos, ornamental plants, bushes, shrubs, hedges, shady or damp areas around buildings. Reapply as necessary. For indoor use, pay particular attention to dark protected areas such as cracks and crevices.
	<b>Crawling insects and pests including:</b> <b>Earwigs</b> ( <i>Forficula auricularia</i> ) <b>House crickets</b> ( <i>Acheta domestica</i> ) <b>Silverfish</b> ( <i>Lepismatidae</i> ) <b>Slaters</b> ( <i>Porcellionidae</i> ) <b>Springtails</b> ( <i>Collembola</i> )		Apply directly onto pests as a spot treatment.
	<b>Houseflies</b> ( <i>Musca domestica</i> )		Apply to surfaces where flies are likely to rest. On non-porous surfaces apply as a coarse spray at the rate of 1 L of emulsion per 20 m <sup>2</sup> . On porous surfaces or use through power equipment, spray at the rate of 750 mL to 1 L of emulsion per 10 m <sup>2</sup> depending on porosity of surface. When treating non-porous and porous surfaces do not exceed the point of run-off.
	<b>Millipedes</b> ( <i>Ommatolulus moreletii</i> )		Outdoor use only. Treat external surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 L of prepared suspension per 25 linear metres. Pay particular attention to potential entry points, such as weep holes, cracks and crevices. Also apply to where pests may be active away from the building including retaining walls, fences, garden beds, sheds etc. Apply at the rate of 1 L of prepared suspension per 16 m <sup>2</sup> . May be applied directly to pests as a spot treatment.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**