NAZWA PLIKU: ID13 Ensystex AUSTRALIA Attrathor 500ml BOOKLET 80x75mm REV-13-202305-1-3-4

MATERIAŁ: folia

WYMIARY: 80 x 75 mm

KOLORYSTYKA: CMYK


CUT —

SIZE -


FRONT



BACK/ BAZA



BOOKLET AWERS



GENERAL INSTRUCTIONS:

ATTRATHOR TARGETED INSECTICIDE is a targeted crack and crevice treatment All trail flow larger to insect its be is a argretic crack and crewe treatment for cockroaches and ants that can also be used for the control of other crawling pests. It can be used in warehouses, restaurants, food manufacturing, processing and storage plants, supermarkets, hotels, hospitals, domestic and other similar situations.

Invert the ATTRATHOR container several times to ensure thorough mixing of concentrate. Half fill the spray tank with water and then add the required quantit of ATTRATHOR. Stir and then top up the spray tank to the required volume. The use of this product in a tank mix with other insecticides is not recommended as the behaviour and efficacy of the product may be affected. Ensure equipment is clean.

INSECTICIDE RESISTANCE WARNING:

GROUP 2B INSECTICIDE

For insecticide resistance management ATTRATHOR Targeted Insecticide is a Group 2B Insecticide. Some naturally occurring insect biotypes resistant to ATTRATHOR Targeted Insecticide and other Group 2B Insecticides may exist ATTRATHOR largeted insections and other droup 26 insectiones may exist individuals can eventually dominate the insect population. The resistant individuals can eventually dominate the insect population if ATTRATHOR Targeted Insecticide or other Group 28 insecticides are used repeatedly. The effectiveness of ATTRATHOR Targeted Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Ensystex Australasia Pty Ltd accepts no liability for any losses that may result from the failure of ATTRATHOR Targeted Insecticide to control resistant insects.

ATTRATHOR Targeted Insecticide may be subject to specific resistance management strategies. For further information contact Ensystex Australasia Pty Ltd.

PRECAUTIONS
Avoid contact with food, food utensils, or places where food is prepared or stored. Before use, remove or cover all exposed foodstuffs. Cover all dishes and utensils, and places where food is prepared or stored. Avoid application around edible plants. Do not apply using equipment carried on the back of the user

RE-ENTRY PERIOD

Do not enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing). Clothing must be laundered after each day's use. Do not allow public re-etry into treated areas until the

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Highly toxic to fish and aquatic organisms. Do NOT apply to areas where surface water is present. Rinse waters and runoff from treated areas MUST be prevented from entering drains or waterways. Do NOT apply if heavy rains are expected to occur within 48 hours of application. Do NOT contaminate streams, rivers or waterways with the chemical or used

containers. Dangerous to bees. PROTECTION OF PETS AND LIVESTOCK

Before spraying remove animals and pets from the areas to be treated.

Cover or remove any open food and water containers. Cover or remove (as applicable) fish ponds, aquariums, etc. before spraying.



BOOKLET AWERS

| SITUATION | PEST | RATE | CRITICAL COMMENTS |
|---|---|-------------------------------|---|
| Internal & external areas & surrounds of Domestic, Commercial, Public and Industrial buildings, facilities & structures (including but not limited, facilities & structures (including but not limited, facilities, farm buildings, ships, trains, acroplanes, facilities, farm buildings, ships, trains, acroplanes, paraneks, private bouses, and external areas associated with such establishments.) | Cockroaches | 50 mL per 5 L water | Inspect premises thoroughly and locate all insect harbourages. Apply directly into harbourages such as cracks and crevices away from where it is likely to be removed by cleaning programs. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control. Apply behind sinks, stoves, and pipes paying particular attention to all cracks and crevices which might contain insects. Repeat treatments when pests begin to reappear. |
| | Ants | | Apply prepared spray to indoor and outdoor surfaces where anits are tracking. Internal harbourage sites include (but are not restricted to) areas such as walls, behind and under sinks, under furniture and indoor plants. External harbourage sites include (but are not restricted to) areas such as walls, behind and under sinks, under furniture and indoor plants. External harbourage sites include (but are not restricted to) areas such as walls, fences, garages, sheds, gazebos, omamental plants, bushes, shrubs, hedges, shady or damp areas around buildings. Reapply as necessary. For indoor use, pay particular attention to dark protected areas such as cracks and crevices. |
| | Crawling insects and pests including: Earwigs (Forticula suricularia) House crickets (Acheta domesticus) Silverfish (Lepismatidae) Slaters (Porcellionidae) Springtails (Collembola) | | Apply directly onto pests as a spot treatment. |
| | Houseflies (Musca domestica) | | Apply to surfaces where files are likely to rest. On non-porous surfaces apply as a coarse spray at the rate of 1 L of emulsion per 20 m². On porous surfaces or use through power equipment, spray at the rate of 750 mL to 1 L of emulsion per 10 m² depending on porosity of surface. When treating non-porous and porous surfaces do not exceed the point of run-off. |
| | Millipedes (Ommatoiulus moreletii) | 115 mL per 5 L of water | Outdoor use only. Treat external surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 L of prepared suspension per 25 linear metres. Pay particular attention to potential entry points, such as weep hotes, cracks and crevices. Also apply to where pests may be active away from the building including retaining walls, fences, garden beds, sheds etc. Apply at the rate of 1 L of prepared suspension per 1 on "May be applied indeerly to pests as a pot treatment." |