

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

RODENTHOR[®]

SOFT BAIT RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM

For effective control of rodents around industrial, commercial, agricultural and residential buildings. Controls rodents resistant to other anticoagulants, including warfarin, coumatetralyl and bromadiolone.

ONE FEED KILLS RATS AND MICE*

ENSYSTEX AUSTRALASIA PTY LTD

ABN 53 102 221 965
Warehouse D, Building 6, The Switchyard
161 Manchester Road
Auburn NSW 2144

www.ensystex.com.au
CUSTOMER SERVICE FREECALL 13 35 36
APVMA APPROVAL NO: 66331/55716




Net Contents:

10 kg

Contains 667 x 15 g bait Sachets
which it is illegal to sell separately

ENSYSTEX[®]



DIRECTIONS FOR USE

Restraints

- DO NOT apply bait to crops.
- DO NOT place bait in the open.
- DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments.
- DO NOT remove bait from paper sachet.

Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits under cover.
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used.
- Ensure baits are removed when rodent activity has ceased.

Clean-Out Treatments (Pulse Baiting)

- Clean-Out treatments are used to remove existing infestations of rats and mice.
- Always record the number and locations of bait sachets for the purpose of bait replacement and later removal.
- For rats, place 1 to 3 baits at intervals of 5 to 9 m in infested areas.
- For mice, place single baits at 2 to 3 m intervals in infested areas.
- The use of secured tamper resistant bait stations is preferred.
- Place baits so as to prevent access to non-target animals. Where access to non-target animals cannot otherwise be prevented, secured tamper resistant bait stations must be used.
- Use the greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken.

Monitoring and/or Maintenance Treatments

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatment may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- For rats and mice place one bait at intervals of between 15 and 30 m in areas prone to rodent infestation.
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.
- Bait stations should be checked at regular intervals according to an agreed and documented schedule.
- Use placebo- or unpoisoned baits if the placement of the bait away from buildings presents a hazard to non-target animals.
- When a rodent infestation is detected the operator should revert to the directions for Clean-Out treatments.

Application

Situations	Pests	Rate	Critical Comments
Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	Rats (<i>Rattus rattus</i> , <i>R. norvegicus</i>), Mice (<i>Mus musculus</i>)	1 to 3 baits/ bait station	Set out bait in places frequented by mice and rats in and around buildings (within 2 m) or enclosed spaces, e.g. drains. Clean-Out Treatment: For rats: Place 1 to 3 baits at intervals of 5 to 9 m in infested areas. For mice: Place single baits at 2 to 3 m intervals in infested areas. See also Monitoring and/or Maintenance Treatments.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Ensure baits are inaccessible to children, pets, domestic animals and wildlife. The baits are ready to use - do not unwrap individual baits. Rats and mice will chew through the wrapper. Place bait along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail in place if relevant. Record the number and location of bait placements for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Look for dead rats or mice, burn or bury them. DO NOT place in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

“ Note: One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimize this.

PRECAUTION Avoid contact with food, food utensils, or places where food is prepared or stored.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. DO NOT place baits in locations which are accessible to pets, domestic animals, livestock or birds. DO NOT contaminate dams, waterways and drains with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 m) or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If no landfill is available, bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS Poisonous if swallowed. DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS Brodifacoum, the active constituent of RODENTHOR Soft Bait Rodenticide, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

SAFETY DATA SHEET Additional information is listed on the Safety Data Sheet and is available from Ensysstex Australasia Pty Ltd on request. Call Customer Service Toll Free on 13 35 36 or visit our web site at www.Ensysstex.com.au

NOTICE Ensysstex Australasia Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Ensysstex Australasia Pty Ltd, or under abnormal conditions.



Batch No.:

Date of Manufacture: