

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

MUSKIL[®]

SOFT BAIT

WITH TWO ACTIVES

FOR FASTER KILL* OF RATS & MICE

FLUO-NP[™]

UV TRACKING TECHNOLOGY

ACTIVE CONSTITUENTS:

0.025 g/kg BROMADIOLONE

0.025 g/kg DIFENACOU

Ready to use bait for the control of rats and mice in and around industrial, commercial, public services, agricultural and residential buildings. Controls rodents resistant to other anticoagulants, including warfarin.

Net Contents:

10 kg

Contains 667 x 15 g bait Sachets which it is illegal to sell separately.

ENSYSTEX AUSTRALASIA PTY LTD

Warehouse D, Building 6, The Switchyard
161 Manchester Road Auburn NSW 2144

www.ensystex.com.au

CUSTOMER SERVICE AND EMERGENCY
RESPONSE (ALL HOURS) 13 35 36

ENSYSTEX[®]

APVMA Approval No. 82450/113141



DIRECTIONS FOR USE

Restraints

DO NOT apply to crops.

DO NOT place baits in the open.

DO NOT use in places accessible to children.

DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments.

DO NOT remove bait from paper sachet.

Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits under cover.
- This product is restricted to use in buildings or enclosed spaces, or within a 2 metre area around buildings.

Clean-Out Treatments (Pulse Baiting)

- Clean-Out treatments are used to remove existing infestations of rats and mice.
- Always map location and number of baits to assist in retrieval and ongoing surveillance.
- For rats, place 1 to 3 baits at intervals of 5 to 9 m in infested areas.
- For mice, place single baits at 2 to 3 m intervals in infested areas.
- The use of tamper resistant bait stations is preferred. Alternatively, nails or wire may be used to fix baits to prevent them being removed by rodents.
- Securely fasten bait and bait stations to minimise risk of consumption by other animals or of being dragged away. Place baits so as to prevent access to non-target animals. Tamper-resistant bait stations must be used at locations where access of non-target animals cannot otherwise be prevented.
- Use the greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. UV (black light) can be used to increase visibility of the fluorescent baits. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication. Most deaths occur 4 to 7 days after consumption of bait.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken.

Monitoring and/or Maintenance Treatments

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatment may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always map location and number of baits to assist in retrieval and ongoing surveillance.
- For rats and mice place one bait at intervals of between 15 and 30 m in areas prone to rodent infestation.
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.
- Bait stations should be checked at regular intervals according to an agreed and documented schedule.

- Use placebo or unpoisoned baits if the placement of the bait away from buildings presents a hazard to non-target animals.
- When a rodent infestation is detected the operator should revert to the directions for Clean-Out treatments.

Application

Situation	Pests	Rate	Critical Comments
Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	Rats (<i>Rattus rattus</i> , <i>R. norvegicus</i>), Mice (<i>Mus musculus</i>)	1 to 3 baits/ bait station	Set out bait in places frequented by rats and mice, in and around buildings (within 2 m) or enclosed spaces, eg drains. Clean-Out Treatment: For rats: Place 1 to 3 baits at intervals of 5 to 9 m in infested areas. For mice: Place single baits at 2 to 3 m intervals in infested areas. See also Monitoring and/or Maintenance Treatments.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS AND RODENT LIKE ANIMALS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

General Instructions

Place bait along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail in place if relevant. The baits are ready to use – do not unwrap individual baits.

Rats and mice will chew through the wrapper.

MUSKIL contains FLUO-NP, a special additive that makes bait and rodent droppings glow bright red under UV light. By shining a handheld UV lamp (black light) on the rodent droppings it is easy to identify and track rodent movement. When used as a perimeter bait, one can more accurately determine if rodents are gaining access to a building's interior and, if so, where the access points are located.

Identifying rodent's movement gives valuable insight for establishing additional bait placements or determining where exclusionary or sanitation measures are needed. FLUO-NP also helps in tracking rodent activity in drop ceilings, sub-floors, wall voids and other dark, hard-to-access areas where the droppings might not normally be visible. The addition of FLUO-NP also helps confirm security of bait placements.

*When compared to brodifacoum soft baits.

PRECAUTION Do not allow contact with food. Avoid contact with food utensils, or places where food is prepared or stored.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. Place bait stations in areas which are inaccessible to non-target animals, wildlife or birds. Map location and number of baits to assist in retrieval and ongoing surveillance. Ensure all baits are removed after treatment. Regularly search for, remove and destroy carcasses

and moribund rodents so that cats, dogs or non-target animals don't eat poisoned rats and mice. Dispose of carcasses by burning or burying. DO NOT contaminate dams, waterways and drains with the bait or its used container.

STORAGE AND DISPOSAL Store in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. DO NOT use containers which have held bait for any other purpose.

SAFETY DIRECTIONS Poisonous if swallowed. Repeated minor exposure may have a cumulative poisoning effect. When using the product wear elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves. Vitamin K (Phytomenadione) is antidotal.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

NOTE TO PHYSICIANS AND VETERINARIANS Bromadiolone and difenacoum, the active constituents of MUSKIL Soft Bait with Two Actives for Faster Kill of Rats and Mice, are long lasting anticoagulant chemicals, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

SAFETY DATA SHEET Additional information is listed on the Safety Data Sheet and is available from Ensysyex on request. Call Customer Service 133536 or visit our web site at www.Ensysyex.com.au

NOTICE Ensysyex Australasia Pty Ltd (Ensysyex) warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Ensysyex, or under abnormal conditions.



Batch No.:

Date of Manufacture: