

GENERAL INSTRUCTIONS

This product is intended for use only in rodent bait stations. DO NOT apply the gel out in the open. Ensure bait is inaccessible to children, pets, domestic animals and wildlife. Place bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and location of bait stations for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Collect and dispose of all dead animals and rodent faeces by burning or burial, and left over bait as per disposal instructions. DO NOT place in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased. #Note: One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimise this. Insert the tube inside the caulking gun; cut a hole in the conical tip of the tube and insert the nozzle. Cut the nozzle above the screw thread taking care not to damage the thread itself, then dispense the product. Once finished, screw on the protecting cap and release the pressure of the caulking gun in order to prevent loss of the product. Keep the tube upside down.

PRECAUTIONS

DO NOT place bait in places accessible to children, domestic animals, livestock and non-target wildlife, unless in tamper resistant bait stations.

DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, or surfaces that come into direct contact with food.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Depending on the amount eaten, this bait may kill or injure. Bait, and dead or affected rodents are hazardous to birds, dogs, cats, pigs, poultry and non-target wildlife. Regularly search for, remove and destroy dead or affected rodents. Place bait stations in areas which are inaccessible to non-target animals, wildlife or birds. Map location and number of baits to assist in retrieval and ongoing surveillance. Ensure all baits are removed after treatment. Regularly search for, remove and destroy carcasses and moribund rodents so that cats, dogs or non-target animals don't eat poisoned rats and mice. Dispose of carcasses by burning or burying. DO NOT contaminate dams, waterways and drains with the bait or its used container.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. DO NOT use containers which have held bait for any other purpose. To the extent possible, unopened baits should be destroyed by burning or burial according to state or territory requirements.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS

Brodifacoum, the active constituent of RODENTHOR GEL Rodenticide, is a long-lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet and is available from Ensynstex Australasia Pty Ltd on request. Call Customer Service 13 35 36 or visit our web site at www.Ensynstex.com.au.

NOTICE

Ensynstex Australasia Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Ensynstex Australasia Pty Ltd, or under abnormal conditions.

Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

RODENTHOR
GEL RODENTICIDE

ACTIVE CONSTITUENT:

0.05 g/kg BRODIFACOU

For effective control of RATS
and MICE around industrial,
commercial, agricultural and
residential buildings.

Controls rodents resistant to other
anticoagulants, including warfarin,
coumatetralyl and bromadiolone

ONE FEED KILLS RATS AND MICE[#]

NET CONTENTS:

300 g

APVMA Approval No: 87706/119034

ENSYNTEX AUSTRALASIA PTY LTD

ABN 53 102 221 965
Warehouse D, Building 6, The Switchyard
161 Manchester Road Auburn NSW 2144

www.ensynstex.com.au

CUSTOMER SERVICE / EMERGENCY RESPONSE
(ALL HOURS) 13 35 36



[#] Registered Trademarks of Ensynstex, Inc. used by licence to Ensynstex Australasia Pty. Ltd.

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply bait to crops.
DO NOT place bait in the open.
DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments.

Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits only within a tamper proof bait station.
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used.
- Ensure baits are removed when rodent activity has ceased.

Clean-Out Treatments (Pulse Baiting)

- Clean-Out treatments are used to remove existing infestations of rats and mice.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- For rats, apply 15 g - 30 g bait gel to the base of bait stations at intervals of 5 to 9 m in infested areas
- For mice, apply 3 to 4 g bait gel to the base of bait stations at 2 to 3 m intervals in infested areas.
- Place baits so as to prevent access to non-target animals. Where access to non-target animals cannot otherwise be prevented, secured tamper-resistant bait stations must be used.
- Use the greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken.

Monitoring and/or Maintenance Treatments

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatment may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- For rats and mice apply up to 15 g bait gel to bait stations at intervals of between 15 and 30 m in areas prone to rodent infestation.
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.
- Bait stations should be checked at regular intervals according to an agreed and documented schedule.
- Use placebo or unpoisoned baits if the placement of the bait away from buildings presents a hazard to non-target animals.
- When a rodent infestation is detected the operator should revert to the directions for Clean-Out treatments.

Application

SITUATION	PESTS	RATE	CRITICAL COMMENTS
Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	Rats (<i>Rattus rattus</i> , <i>R. norvegicus</i>)	15 g to 30 g (15 to 30 mL) bait gel per bait station	Set out bait in places frequented by mice and rats in and around buildings (within 2 m) or enclosed spaces, e.g. drains.
	Mice (<i>Mus musculus</i>)	3 g to 4 g (3 to 4 mL) bait gel per bait station	Clean-Out Treatment: For rats: Place bait stations at intervals of 5 to 9 m in infested areas. For mice: Place bait stations at 2 to 3 m intervals in infested areas. See also Monitoring and/or Maintenance Treatments.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

L-AU-RDTG1000 1.03.05.23

ENSYNTEX