## **Product Data Sheet**



Product Name 3-D Life Scrambled RGD Peptide

Catalog Number 09-P-003

Description RGD sequences are motifs on extracellular matrix proteins promoting

adhesion to cellular integrin receptors. The 3-D Life Scrambled RGD Peptide contains a scrambled RGD motif and a thiol group on the N-terminus. The thiol group is used to immobilize the peptide on maleimide polymers in 3-D Life Hydrogels. The immobilized peptide is used in control experiments to determine the specificity of cell adhesion caused by a genuine RGD peptide (e.g. 3-D Life RGD Peptide).

For instructions, please consult General Protocols GP-1 "Preparation of 3-D Life Fast Gelling Hydrogels" or GP-2 "Preparation of 3-D Life Slow Gelling Hydrogels" and the 3-D Life Hydrogels User Guide on www.cellendes.com.

Quantity 1 µmol

Components

Material	Quantity	Concentration of reactive groups	Storage
Scrambled RGD peptide, lyophilized	1 µmol	20 mmol/L*	Lyophilisate and after reconstitution: -20°C or lower
Water	600 µl	n/a	Room temperature or lower

All materials are filter-sterilized.

## Reconstitution Scrambled RGD Peptide:

- 1. Briefly centrifuge vial containing the Scrambled RGD Peptide lyophilisate to make sure that entire product is at the bottom of the reaction tube.
- 2. Add 48 µl 3-D Life Water per tube of Scrambled RGD Pepetide to dissolve lyophilisate at a concentration of 20 mmol/L thiol groups.
- 3. Close tube and briefly vortex.
- 4. Incubate for 5 min.
- 5. Briefly vortex and centrifuge again.
- 6. The Scrambled RGD Peptide is now ready for use.

Pept. Sequence Acetyl-Cys-Doa\*-Doa-Gly-Arg-Asp-Gly-Ser-Pro-NH2

\*: Doa:8-amino-3,6-dioxaoctanoic acid

Molar Mass 1022 g/mol

>90% (HPLC) Purity

> Intended for research use only. Not for use in human therapeutic or diagnostic applications.



www.cellendes.com