

Product Data Sheet



Product Name *3-D Life* Dextran-CD Hydrogel SG

Catalog Number G93-1

Description The *3-D Life* Dextran-CD Hydrogel SG Kit provides reagents for setting up slow gelling, cell-compatible hydrogels. Its major components are SG-Dextran and the crosslinker CD-Link. When the two reagents are combined, thiol groups on CD-Link form stable thioether bonds with thiol-reactive groups on SG-Dextran, which results in the formation of the gel. The components are mixed at physiological pH (pH 7.2) for optimal cell compatibility. The slow gelation kinetics allows enough time to conveniently manipulate the solution before the onset of gel formation.

CD-Link is composed of polyethylene glycol and a matrix metalloprotease (MMP)-cleavable peptide (Pro-Leu-Gly-Leu-Trp-Ala). The MMP-cleavable peptide is designed for a broad range of MMP cleavage including MMPs MMP1, MMP3, MMP7 and MMP9 [1]. It allows cells to spread and migrate within the hydrogel if they express the indicated MMPs. In most cases cell spreading and migration also requires the presence of adhesion peptides.

Prior to the crosslinking step, cell adhesion peptides (e.g. *3-D Life* RGD Peptide, Cat. No. 09-P-001) can be covalently attached to a portion of the SH-reactive groups on SG-Dextran to provide a cell-adhesive matrix.

SG-Dextran hydrogels can be dissolved by the addition of dextranase (*3-D Life* Dextranase, Cat. No. D10-1), which allows the recovery of chemically fixed or live cells for post-culture analyses (e.g. RT-PCR) or for further cultivation.

For instructions, please consult the General Protocol GP-2 "Preparation of *3-D Life* Slow Gelling Hydrogels" and the *3-D Life* Hydrogels User Guide on www.cellendes.com.

Quantity Allows formation of up to 2 ml *3-D Life* Hydrogel depending on the stiffness of the gel.

Components

Material	Quantity	Concentration of reactive groups	Storage
SG-Dextran	170 µl	30 mmol/L	Short term (≤2 months): 4°C Long Term: -80°C
CD-Link, lyophilized	200 µl*	20 mmol/L	Lyophilisate and after reconstitution: -20°C to -80°C
10 x CB (pH 7.2)	200 µl	n.a.	Short term (≤2 months): 4°C Long term: -20°C to -80°C
Water	2x 1500 µl	n.a	RT to -80°C

All materials are filter-sterilized.

*Volume/concentration after reconstitution of lyophilisate.

Continued on next page.

Reconstitution CD-Link:

1. Briefly centrifuge vial containing CD-Link lyophilisate to make sure that the entire material is at the bottom of the reaction tube.
2. Add 188 µl *3-D Life Water* per tube for a concentration of 20 mmol/L thiol groups.
3. Close tube and briefly vortex.
4. Incubate for 5 min.
5. Briefly vortex and centrifuge again.
6. CD-Link is now ready for use.

References [1] Knight, C. G. et al. FEBS 296:263-66 (1992)

Intended for research use only. Not for use in human therapeutic or diagnostic applications.

Distributed by:
Ilex Life Sciences LLC
Tel: (828) 531-9949
Email: info@ilexlife.com
Web: <https://ilexlife.com/>

