Product Data Sheet



Product Name 3-D Life PEG-Link

Catalog Number L50-1

Description

3-D Life PEG-Link is a component of the 3-D Life Hydrogel system. The PEG-Link molecule consists of polyethylene glycol carrying a thiol at each end. When combined with polymers of the 3-D Life Hydrogel system, thiol groups on PEG-Link form stable thioether bonds with the thiol-reactive groups on the polymers, which results in the formation of the hydrogel. PEG-Link can be used with 3-D Life functionalised polymers Mal-PVA, Mal-Dextran, SG-PVA and SG-Dextran. For using this product please consult General Protocols GP-1 "Preparation of 3-D Life Fast Gelling Hydrogels" or GP-2 "Preparation of 3-D Life Slow Gelling Hydrogels" and the 3-D Life Hydrogels User Guide on www.cellendes.com.

Quantity

When used with polymers of the 3-D Life Hydrogel system up to 2 ml 3-D Life Hydrogel can be generated, depending on the stiffness of the gel.

Components

Material	Quantity	Concentration of reactive groups	Storage
PEG-Link, lyophilized	200 μΙ*	20 mmol/L*	Lyophilisate and after reconstitution: -20°C to -80°C
Water	600 µl	n/a	Room temperature or lower

All materials are filter-sterilized.

Reconstitution PEG-Link:

- 1. Briefly centrifuge vial containing the PEG-Link lyophilisate to make sure that the entire material is at the bottom of the reaction tube.
- 2. Add 188 µl 3-D Life Water per tube for a concentration of 20 mmol/L thiol groups. This results in a 200 µl PEG-Link solution.
- 3. Close tube and briefly vortex.
- 4. Incubate for 5 min.
- 5. Briefly vortex and centrifuge again.
- 6. PEG-Link is now ready for use.

Intended for research use only. Not for use in human therapeutic or diagnostic applications.



Distributed by: Ilex Life Sciences LLC Tel: (828) 531-9949 Email: info@ilexlife.com Web: https://ilexlife.com/

^{*}Volume/concentration after reconstitution of lyophilisate.