Product Data Sheet



Product Name 3-D Life HyLink

Catalog Number L70-3

Description 3-D Life HyLink is a component of the 3-D Life Hydrogel system. The HyLink

molecule consists of hyaluronic acid carrying thiol groups along its backbone. When combined with SG-Polymers of the *3-D Life* Hydrogel system, thiol groups on HyLink form stable thioether bonds with the thiol-reactive groups on the polymers, which results in the formation of the hydrogel. HyLink can be used with *3-D Life* functionalised polymers SG-PVA and SG-Dextran. For using this product please consult General Protocols GP-4 "Preparation of *3-D Life* Hyaluronic Acid (HA) Hydrogels" and the *3-D Life* Hydrogels User

Guide on www.cellendes.com.

Quantity Sufficient for 5 ml hydrogel of a crosslinking strength of 1.2 mmol/l when

used with SG-Polymers of the 3-D Life Hydrogel system.

Components

Material	Quantity	Concentration of reactive groups	Storage
HyLink, lyophilized	3x 200 μl*	10 mmol/L*	Lyophilisate and after reconstitution: -70°C
Water	600 µl	n/a	Room temperature or lower

All materials are filter-sterilized.

Reconstitution HyLink:

- 1. Briefly centrifuge vial containing HyLink lyophilisate to make sure that the entire material is at the bottom of the reaction tube.
- 2. Add 188 μ l *3-D Life* Water per tube for a concentration of 10 mmol/L thiol groups.
- 3. Close tube and briefly vortex.
- 4. Incubate for 60 min at room temperature.
- 5. Briefly vortex and centrifuge again.
- 6. HyLink is now ready for use.

Notes

After reconstitution, HyLink is a viscous solution. Low binding pipet tips are recommended for accurate pipetting up to 20 μ l. For volumes above 20 μ l use wide orifice pipet tips.

Intended for research use only. Not for use in human therapeutic or diagnostic applications.



Distributed by: Ilex Life Sciences LLC Tel: (828) 531-9949 Email: info@ilexlife.com Web: https://ilexlife.com/

^{*}Volume/concentration after reconstitution of lyophilisate.