Maximum Marks: 200 Time: 45 Minutes

## **General Instructions:**

- (i) This paper consists of 50 MCQs, attempt any 40 out of 50.
- (ii) Correct answer or the most appropriate answer: Five marks (+5).
- (iii) Any incorrect option marked will be given minus one mark (-1).
- (iv) Unanswered/Marked for Review will be given no mark (0).
- (v) If more than one option is found to be correct then Five marks (+5) will be awarded to only those who have marked any of the correct options.
- (vi) If all options are found to be correct then Five marks (+5) will be awarded to all those who have attempted the question.
- (vii) If none of the options is found correct or a Question is found to be wrong or a Question is dropped then all candidates who have appeared will be given five marks (+5).

(viii) Calculator / any electronic gadgets are not permitted.

- **1.** Which of the following conditions favours the existence of a substance in the solid state?
  - (1) High temperature
  - (2) Low temperature
  - (3) High thermal energy
  - (4) Weak cohesive forces
- **2.** Calculate the emf of the following cell at 298 K:  $Mg(s) | Mg^{2+} (0.1 M) | | Cu^{2+} (1.0 \times 10^{-3} M) | Cu(s)$  $[Given = E^{\circ}_{Cell} = 2.71 V]$ 
  - (1) 1.426 V
- (2) 2.503 V
- (3) 2.651 V
- (4) 1.8 V
- **3.** The lattice site in a pure crystal cannot be occupied
  - (1) molecule
- (2) ion
- (3) electron
- (4) atom
- **4.** There are 14 elements in actinoid series. Which of the following element does not belong to this series?
  - (1) U
- (2) Np
- (3) Tm
- (4) Fm
- **5.** The IUPAC name of the compound shown below is:



- (1) 2-bromo-6-chlorocyclohex-1-ene
- (2) 6-bromo-2-chlorocyclohexene
- (3) 3-bromo-1-chlorocyclohexene
- (4) 1-bromo-3-chlorocyclohexene

- 6. Which stoichiometric defect does not change the density of the crystal?
  - (1) Frenkel defect
- (2) Schottky defect
- (3) Interstitial defect
- (4) F-centres
- 7. Which of the following species can act as the strongest base?
  - (1) <sup>Θ</sup>OH
- (2) <sup>Θ</sup>OR
- (3) <sup>Θ</sup>OC<sub>6</sub>H<sub>5</sub>
- **8.** The pair  $[Co(NH_3)_4Cl_2]Br_2$  and  $[Co(NH_3)_4Br_2]Cl_2$ will show
  - (1) Linkage isomerism
  - (2) Hydrate isomerism
  - (3) Ionization isomerism
  - (4) Coordinate isomerism
- **9.** Following reactions occur at cathode during the electrolysis of aqueous silver chloride solution:

$$Ag^{+}(aq) + e^{-} \rightarrow Ag(s)$$
 ;  $E^{\circ} = +0.80 \text{ V}$   
 $H^{+}(aq) + e^{-} \rightarrow \frac{1}{2}H_{2}(g)$  ;  $E^{\circ} = 0.00 \text{ V}$ 

On the basis of their standard reduction electrode potential (E°) values, which reaction is feasible at the cathode?

- (1)  $Ag^{+}(aq) + e^{-} \rightarrow Ag(s)$
- $E^{\circ} = +0.80 \text{ V}$
- (2)  $H^+(aq) + e^- \rightarrow \frac{1}{2}H_2(g)$   $E^\circ = 0.00 \text{ V}$
- (3) Both reactions are feasible
- (4) None of the above

## 2

- **10.** Electronic configuration of a transition element X in +3 oxidation state is [Ar] $3d^5$ . What is its atomic number?
  - **(1)** 25
- **(2)** 26
- (3) 27
- **(4)** 24
- **11.** IUPAC name of *m*-cresol is
  - (1) 3-methylphenol
- (2) 3-chlorophenol
- (3) 3-methoxyphenol
- (4) benzene-1,3-diol
- **12.** In which mode of expression, the concentration of a solution remains independent of temperature?
  - (1) Molarity
- (2) Normality
- (3) Formality
- (4) Molality
- **13.** Write the product(s) in the following reactions:

- (1) No product formed (2)
- (3)
- (4) OH

OH.


- **14.** What type of isomerism is shown by the pair  $[Cr(H_2O)_6]Cl_3$  and  $[Cr(H_2O)_5Cl]Cl_2 \cdot H_2O$ ?
  - (1) Ionisation isomerism
  - (2) Coordination isomerism
  - (3) Solvate isomerism
  - (4) Linkage isomerism
- **15.** In the presence of a catalyst, heat evolved or absorbed during reaction
  - (1) increases.
  - (2) decreases.
  - (3) remains unchanged.
  - (4) may increase or decrease.
- **16.** Identify the compound Y in the following reaction.

$$(a) \qquad \begin{array}{c} NH_2 \\ \hline NaNO_2 + HCI \\ \hline 273 - 278 \text{ K} \end{array} \qquad \begin{array}{c} \mathring{N}_2 \bar{C}I \\ \hline Cu_2 Cl_2 \\ \hline \end{array} \qquad Y + N$$


$$(a) \qquad \begin{array}{c} CI \\ \hline \end{array} \qquad (4) \qquad \begin{array}{c} CI \\ \hline \end{array}$$

- **17.** The oxidation state of Fe in  $K_3$  [Fe( $C_2O_4$ )<sub>3</sub>] is
  - **(1)** 0
- **(2)** 2
- **(3)** 3
- **(4)** 4
- **18.** Bond angle in ethers is slightly more than
  - (1) Square planar angle
  - (2) Trigonal bipyramidal angle
  - (3) Tetrahedral angle
  - (4) None of the above


- **19.** Which one of the following is not applicable to the phenomenon of adsorption?
  - **(1)**  $\Delta H > 0$
- (2)  $\Delta G < 0$
- (3)  $\Delta S < 0$
- **(4)**  $\Delta H < 0$
- **20.** Arrange the following compounds in increasing order of acid strength.





- (1) (i) > (ii) > (iii)
- (2) (ii) < (i) < (iii)
- (3) (iii) < (i) < (ii)
- (4) (iii) > (i) > (ii)
- **21.** Which of the following structures is enantiomeric with the molecule (a) given below?




- **72.** Amongst the following, the most stable complex is
  - (1)  $[Fe(H_2O)_6]^{3+}$
- (2)  $[Fe(NH_3)_6]^{3+}$
- (3)  $[Fe(C_2O_4)_3]^{3-}$
- (4) [FeCl<sub>6</sub>]<sup>3-</sup>
- **23.** K<sub>H</sub> value for Ar(g), CO<sub>2</sub>(g), HCHO(g) and CH<sub>4</sub>(g) are 4.039, 1.67,  $1.83 \times 10^{-5}$ , and 0.143, respectively. Arrange these gases in the order of their increasing solubility.
  - (1)  $HCHO < CH_4 < CO_2 < Ar$
  - (2)  $HCHO < CO_2 < CH_4 < Ar$
  - (3)  $Ar < CO_2 < CH_4 < HCHO$
  - (4)  $Ar < CH_4 < CO_2 < HCHO$
- **24.**  $CH_3CONH_2$  on reaction with NaOH and  $Br_2$  in alcoholic medium gives:
  - (1) CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>
- (2) CH<sub>3</sub>CH<sub>2</sub>Br
- (3) CH<sub>3</sub>NH<sub>2</sub>
- (4) CH<sub>3</sub>COONa
- **25.** Which of the following pairs represents anomers?










- **26.** Which of the following is not a favourable condition for physical adsorption?
  - (1) High pressure
  - (2) Negative δH
  - (3) Higher critical temperature of adsorbate
  - (4) High temperature
- **27.** Which of the following will give a white precipitate upon reacting with AgNO<sub>3</sub>?
  - (1)  $K_2 [Pt(en)_2Cl_2]$
- (2)  $[Co(NH_3)_3Cl_3]$
- (3)  $[Cr(H_2O)_6]Cl_3$
- (4) [Fe(H<sub>2</sub>O)<sub>3</sub>Cl<sub>3</sub>]
- **28.** Which one of the following is not a condensation polymer?
  - (1) Melamine
- (2) Glyptal
- (3) Dacron
- (4) Neoprene
- **29. Assertion (A):** Rate of reaction doubles when concentration of reactant is doubled if it is a first order reaction.

Reason (R): Rate constant also doubles.

- (1) Both A and R are true and R is the correct explanation of A
- (2) Both A and R are true but R is NOT the correct explanation of A
- (3) A is true but R is false
- (4) A is false and R is true
- **30.** The oxidation of Ni in  $[Ni(CO)_4]$  is
  - **(1)** 0
- **(2)** 2
- **(3)** 3
- **(4)** 4
- **31.** Which of the following reagents would not be a good choice for reducing an aryl nitro compound to an amine?
  - (1)  $H_2(excess)/Pt$
- (2) LiAlH<sub>4</sub> in ether
- (3) Fe and HCl
- (4) Sn and HCl

- **32.** Which of the following statement is correct?
  - (1) The rate of a reaction decreases with passage of time as the concentration of reactants decreases.
  - (2) The rate of a reaction is same at any time during the reaction.
  - (3) The rate of a reaction is independent of temperature change.
  - (4) The rate of a reaction decreases with increase in concentration of reactant(s).
- **33.** Which of the following should be most volatile?
  - i) CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>
- (ii) (CH<sub>3</sub>)<sub>3</sub>N



- (1) (ii)
- (2) (iv)
- (3) (i)
- (4) (iii)
- **34.** A colloidal system having a solid substance as a dispersed phase and a liquid as a dispersion medium is classified as
  - (1) solid sol
- (2) gel
- (3) emulsion
- (3) sol
- **35.** Which of the following statements is not correct?
  - (1) Copper liberates hydrogen from acids.
  - (2) In its higher oxidation states, manganese forms stable compounds with oxygen and fluorine.
  - (3) Mn<sup>3+</sup> and Co<sup>3+</sup> are oxidising agents in aqueous solution.
  - (4)  $Ti^{2+}$  and  $Cr^{2+}$  are reducing agents in aqueous solution.
- **36.** Which of the following compounds will give butanone on oxidation with alkaline  ${\rm KMnO_4}$  solution?
  - (1) Butan-1-ol
- (2) Butan-2-ol
- (3) Both of these
- (4) None of these
- **37.** Value of Henry's constant K<sub>H</sub>\_\_\_\_\_\_
  - (1) increases with increase in temperature.
  - (2) decreases with increase in temperature
  - (3) remains constant
  - (4) first increases then decreases.
- **38.** Which one of the following elements constitutes a major impurity in pig iron?
  - (1) Silicon
- (2) Oxygen
- (3) Sulphur
- (4) Graphite
- **39.** Which of the following help in food preservation by retarding the action of oxygen on food.
  - (1) Food colours
- (2) Antioxidants
- (3) Preservatives
- (4) Fat emulsifiers
- **40.** Debye-Huckel Onsager equation for strong electrolytes:

$$\wedge = \wedge_{o} - A\sqrt{C}$$

Which of the following equality holds?

- (1)  $\wedge = \wedge_0 \text{ as } C \longrightarrow \sqrt{A}$  (2)  $\wedge = \wedge_0 \text{ as } C \longrightarrow \infty$
- (3)  $\wedge = \wedge_0 \text{ as } C \longrightarrow 0$  (4)  $\wedge = \wedge_0 \text{ as } C \longrightarrow 1$
- **41.** For Lysine,  $H_2N-(CH_2)_4-CH-COOH$  , which of NH,

the following is incorrect?

- (1) α-Amino acid.
- (2) Basic amino acid.
- (3) Amino acid synthesised in body.
- (4) β-Amino acid.
- **42.** Which one of the following sets forms the biodegradable polymer?
  - (1)  $CH_2$ =CH-CN and  $CH_2$ =CN-CH= $CH_2$
  - (2)  $H_2N$ — $CH_2$ —COOH and  $H_2N$ — $(CH_2)_5$ —COOH
  - (3)  $HO CH_2 CH_2 OH$  and HOOC COOH

(4) 
$$\bigcirc$$
 —CH=CH<sub>2</sub> and CH<sub>2</sub>—CH—CH—CH<sub>2</sub>

- **43.** Which of the following can possibly be used as an analgesic without causing addiction and mood modification?
  - (1) Diazepam
  - (2) Tetrahydrocatenol
  - (3) Morphine
  - (4) N-Acetyl-para-aminophenol
- **44.** Rate law cannot be determined from balanced chemical equation if
  - (1) reverse reaction is involved.
  - (2) it is an elementary reaction.
  - (3) it is a sequence of elementary reactions.
  - (4) any of the reactants is in excess.
- **45.** Glucose and fructose are
  - (1) isomers of each other
  - (2) homologous of each other

- (3) anomers of each other
- (4) enantiomers of each other
- **46.** Cassiterite is an ore of:
  - (1) Mn
- (2) Ni
- (3) Sb
- (4) Sn

## **Case Based**

## Read the passage given below and answer the following questions:

In spite of the predictions of stable noble gas compounds since at least 1902, unsuccessful attempts at their synthesis gave rise to the widely held opinion that noble gases are not only noble but also inert. It was not until 1962 that this dogma was shattered when Bartlett in Canada published the first stable noble gas compound XePtF<sub>6</sub>. This discovery triggered a worldwide frenzy in this area, and within a short time span many new xenon, radon, and krypton compounds were prepared and characterized. The recent discoveries show the ability of xenon to act as a ligand. The discovery by Seppelt's group that more than one xenon atom can attach itself to a metal center which in the case of gold leads to surprisingly stable Au- Xe bonds. The bonding in [AuXe<sub>4</sub>]<sup>2+</sup> involves 4 Xe ligands attached by relatively strong bonds to a single Au(II) center in a square planar arrangement with a Xe-Au bond length of about 274 pm This discovery provides not only the first example of multiple xenon ligands but also represents the first strong metal - xenon bond.

- **47.** In the complex ion  $[AuXe_4]^{2+}$ , Xe acts as:
  - (1) central atom
- (2) ligand
- (3) chelating agent
- (4) electrophile
- **48.** Hybridisation shown by Au in  $[AuXe_4]^{2+}$  is:
  - (1)  $sp^3$
- (2)  $sp^3d$
- (3)  $sp^3d^2$
- (4)  $sp^2$
- **49.** Compounds of noble gases except \_\_\_\_\_ are known
  - (1) Krypton
- (2) Radon
- (3) Helium
- (4) Xenon
- **50.** Xe is a \_\_\_\_\_ ligand.
  - (1) ambidentate
- (2) bidentate
- (3) unidentate
- (4) hexadentate