





Trace the Mind Map


- First Level
- Second Level
- Third Level












ALDEHYDES AND KETONES:

(i) Physical:


Boiling points are higher than hydrocarbons and ethers of comparable molecular masses and lower than alcohols of similar molecular masses due to absence of intermolecular hydrogen bonding.

(ii) Chemical :Nucleophilic addition reactions :

Aldehydes are more reactive than ketones due to steric and electronic reasons.



Reduction : (a) To alcohols – aldehydes and ketones reduce to primary and secondary alcohols respectively by NaBH_4 or LiAlH_4 .




Oxidation: $\text{RCHO} \xrightarrow{[\text{O}]} \text{RCOOH}$ Silver mirror




Tollen's test: $\text{RCHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH}^- \rightarrow \text{RCOO}^- + 2\text{Ag} \downarrow + 2\text{H}_2\text{O} + 4\text{NH}_3$

Fehling's test: $\text{RCHO} + 2\text{Cu}^{2+} + 5\text{OH}^- \rightarrow \text{RCOO}^- + \text{Cu}_2\text{O} \downarrow + 3\text{H}_2\text{O}$ Red brown ppt

Haloform reaction:



Reactions due to α -hydrogen:



Cannizzaro reaction : $2\text{HCHO} + \text{conc KOH} \xrightarrow{\Delta} \text{CH}_3\text{OH} + \text{HCOOK}$

Electrophilic substitution reaction:





Carboxylic acids:

(i) Physical: Higher boiling points than aldehydes, ketones or alcohols. Solubility decreases with increasing number of C atoms.

(ii) Chemical : $2\text{RCOOH} + 2\text{Na} \rightarrow 2\text{RCOONa} + \text{H}_2$

Forms corresponding anhydride on heating with mineral acids



1. Aldehydes and Ketones


Common names :

- Replacing -e with -al and -one as required.
- Alkyl phenyl ketones by adding acyl group as prefix to phenone.

IUPAC names :

- Replacing -e with -al and -one as required.

Structure of Carbonyl group




2. Carboxylic Acids

Common names : end with -ic with -oic acid.

- IUPAC names : replace -e in the corresponding alkane with -oic acid.

Structure of Carboxyl Group



3. USES

- (a) Carboxylic acids**
 - Methanoic acid in rubber, textile, dyeing, leather industries.
 - Ethanoic acid as solvent.
 - Higher fatty acids in manufacture of soaps and detergents.
- (b) Aldehydes of ketones**
 - As solvents.
 - Starting materials and reagents for synthesis of products.

ALDEHYDES:


1. From acyl chloride




2. From nitriles and esters : Stephen reaction



3. From hydrocarbons : Eiland reaction



Gattermann - Koch reaction




KETONES:


1. From acyl chloride



2. From nitriles




3. From benzene or substituted benzenes




Carboxylic Acids:


1. From primary alcohols and aldehydes




2. From alkylbenzenes




3. From nitriles and amides



4. From Grignard reagents



5. From acyl halides and anhydrides



6. From esters



Nomenclature, Structure and Uses

Preparation


Properties

Aldehydes, Ketones and Carboxylic Acids



Trace the Mind Map

- First Level
- Second Level
- Third Level




Optically active polyhydroxy aldehydes or ketones or compounds which produce units with specific functional groups on hydrolysis.

Classification :

- (i) **Monosaccharides** : (Aldehyde group – aldose, keto group –ketose)
Glucose : Preparation : $C_6H_{12}O_6 + H_2O \xrightarrow{H^+} C_6H_{12}O_6 + C_6H_{12}O_6$ (Glucose Fructose)
- (a) **From starch** : $(C_6H_{10}O_5)_n + nH_2O \xrightarrow{393K; 2-3 \text{ atm}} nC_6H_{12}O_6$

Structure :



- (II) **Disaccharides** : (Sucrose, maltose) Linkage between 2 monosaccharides– Glycosidic linkage
- (III) **Polysaccharides** : Large number of monosaccharides units joined by glycosidic linkages. e.g., (a) Starch : Polymer of α -glucose with two components amylose and amylopectin (c) Glycogen

Importance :

- Form a major portion of food.
- As storage molecules.
- Cellulose forms cell wall of bacteria and plants.
- Raw materials for industries like textiles, paper, lacquers and breweries.

- (i) A sequence of bases on DNA is unique for a person and information regarding this is called DNA fingerprinting. It is same for every cell and cannot be altered by any known treatment. DNA fingerprinting is now used
- (ii) in forensic laboratories for identification of criminals.
- (iii) to determine paternity of an individual.
- (iv) to identify the dead bodies in any accident by comparing the DNA's of parents or children.
- (v) to identify racial groups to rewrite biological evolution.

Enzymes
 Globular proteins specific for particular reaction and for particular substrate.
 Mechanism : Reduces the magnitude of activation energy

Carbohydrates

Proteins

(Polymers of amino acids)
 -Amino acids contain -NH₂ and -COOH group.

Classification :


- **On the basis of relative number of -NH₂ and -COOH group**:
 (i) Neutral : Equal number of -NH₂ and -COOH group.
 (ii) Basic : More number of -NH₂ than -COOH group.
 (iii) Acidic : More number of -COOH than -NH₂ group.

• **On the basis of place of synthesis :**

- (i) Essential – cannot be synthesized in the body.
- (ii) Non-essential – synthesized in the body.

• **On the basis of shape :**

- (i) Fibrous : Fibre-like structure
- (ii) Globular : Spherical



Denaturation of protein :

When a protein in its native form is subjected to physical change, globules unfold, helix get uncoiled and protein loses its biological activity.

Organic compounds required in diet in small amounts to perform specific biological functions for maintenance and growth.

Classification :

- (i) **Fat soluble** : Soluble in fats and oils but insoluble in water. (Vitamins A, D, E and K)
- (ii) **Water soluble** : B group and vitamin C are soluble in water.

Vitamin

Chromosomes : Particles in nucleus responsible for heredity. Chromosomes are made up of proteins and nucleic acid.


Two types : Deoxyribonucleic acid (DNA), ribonucleic acid (RNA)

Composition : In DNA, sugar is β -D-2-deoxyribose whereas in RNA is β -D-ribose. DNA contains A,G,C,T whereas RNA has A,G,C,U.

Structure :

Nucleoside : Formed by attachment of a base to 1' of sugar

Nucleotide : Formed by link to phosphoric acid at 5' of sugar.



Types of RNA : m-RNA, t-RNA, r-RNA

Biological Functions :

- Chemical basis of heredity.
- Responsible for identity of different species of organisms.
- Nucleic acids are responsible for protein synthesis in cell.

Nucleic Acids

DNA Fingerprinting :
 Unique sequence of bases on DNA

Biomolecules

Trace the Mind Map
 • First Level • Second Level • Third Level