

# Solved Paper, 2021-2022

## MATHEMATICS

### Term-I, Set-4

Series : SSJ/2

Question Paper

Code No. 065/2/4

Time allowed : 90 Minutes

Max. Marks : 40

#### General Instructions :

- This question paper comprises of 50 questions out of which 40 questions are to be attempted as per instructions. All questions carry equal marks.
- The question paper consists of three Sections – Section A, B and C.
- Section – A contains 20 questions. Attempt any 16 questions from Q. No. 1 to 20.
- Section – B also contains 20 questions. Attempt any 16 questions from Q. No. 21 to 40.
- Section – C contains 10 questions including one Case Study. Attempt any 8 questions from Q. No. 41 to 50.
- There is only one correct option for every Multiple Choice Question (MCQ). Marks will not be awarded for answering more than one option.
- There is no negative marking.

#### SECTION - A

In this section, attempt any 16 questions out of Questions 1-20. Each question is of one mark.

- Differential of  $\log [\log (\log x^5)]$  w.r.t  $x$  is
  - $\frac{5}{x \log(x^5) \log(\log x^5)}$
  - $\frac{5}{x \log(\log x^5)}$
  - $\frac{5x^4}{\log(x^5) \log(\log x^5)}$
  - $\frac{5x^4}{\log x^5 \log(\log x^5)}$
- The number of all possible matrices of order  $2 \times 3$  with each entry 1 or 2 is
  - 16
  - 6
  - 64
  - 24
- A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is defined as  $f(x) = x^3 + 1$ . Then the function has
  - no minimum value
  - no maximum value
  - both maximum and minimum values
  - neither maximum value nor minimum value
- If  $\sin y = x \cos(a + y)$ , then  $\frac{dx}{dy}$  is
  - $\frac{\cos a}{\cos^2(a + y)}$
  - $\frac{-\cos a}{\cos^2(a + y)}$
  - $\frac{\cos a}{\sin^2 y}$
  - $\frac{-\cos a}{\sin^2 y}$
- The points on the curve  $\frac{x^2}{9} + \frac{y^2}{25} = 1$ , where tangent is parallel to X-axis are
  - $(\pm 5, 0)$
  - $(0, \pm 5)$
  - $(0, \pm 3)$
  - $(\pm 3, 0)$
- Three points  $P(2x, x + 3)$ ,  $Q(0, x)$  and  $R(x + 3, x + 6)$  are collinear, then  $x$  is equal to
  - 0
  - 2
  - 3
  - 1
- The principal value of  $\cos^{-1}\left(\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right)$  is
  - $\frac{\pi}{12}$
  - $\pi$
  - $\frac{\pi}{3}$
  - $\frac{\pi}{6}$
- If  $(x^2 + y^2)^2 = xy$ , then  $\frac{dy}{dx}$  is
  - $\frac{y + 4x(x^2 + y^2)}{4y(x^2 + y^2) - x}$
  - $\frac{y - 4x(x^2 + y^2)}{x + 4(x^2 + y^2)}$
  - $\frac{y - 4x(x^2 + y^2)}{4y(x^2 + y^2) - x}$
  - $\frac{4y(x^2 + y^2) - x}{y - 4x(x^2 + y^2)}$
- If a matrix A is both symmetric and skew symmetric, then A is necessarily a
  - Diagonal matrix
  - Zero square matrix

- (c) Square matrix  
(d) Identity matrix
10. Let set  $X = \{1, 2, 3\}$  and a relation  $R$  is defined in  $X$  as:  
 $R = \{(1, 3), (2, 2), (3, 2)\}$ , then minimum ordered pairs which should be added in relation  $R$  to make it reflexive and symmetric are
- (a)  $\{(1, 1), (2, 3), (1, 2)\}$   
(b)  $\{(3, 3), (3, 1), (1, 2)\}$   
(c)  $\{(1, 1), (3, 3), (3, 1), (2, 3)\}$   
(d)  $\{(1, 1), (3, 3), (3, 1), (1, 2)\}$
11. A Linear Programming Problem is as follows:  
Minimise  $z = 2x + y$   
subject to the constraints  $x \geq 3, x \leq 9, y \geq 0$   
 $x - y \geq 0, x + y \leq 14$
- The feasible region has
- (a) 5 corner points including  $(0, 0)$  and  $(9, 5)$   
(b) 5 corner points including  $(7, 7)$  and  $(3, 3)$   
(c) 5 corner points including  $(14, 0)$  and  $(9, 0)$   
(d) 5 corner points including  $(3, 6)$  and  $(9, 5)$
12. The function  $f(x) = \begin{cases} e^{3x} - e^{-5x} & \text{if } x \neq 0 \\ k & \text{if } x = 0 \end{cases}$  is continuous at  $x = 0$  for the value of  $k$ , as
- (a) 3 (b) 5  
(c) 2 (d) 8
13. If  $C_{ij}$  denotes the cofactor of element  $P_{ij}$  of the matrix  $P = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & 2 & 4 \end{bmatrix}$ , then the value of  $C_{31} \cdot C_{23}$  is
- (a) 5 (b) 24  
(c) -24 (d) -5
14. The function  $y = x^2 e^{-x}$  is decreasing in the interval
- (a)  $(0, 2)$  (b)  $(2, \infty)$   
(c)  $(-\infty, 0)$  (d)  $(-\infty, 0) \cup (2, \infty)$
15. If  $R = \{(x, y); x, y \in \mathbb{Z}, x^2 + y^2 \leq 4\}$  is a relation in set  $\mathbb{Z}$ , then domain of  $R$  is
- (a)  $\{0, 1, 2\}$   
(b)  $\{-2, -1, 0, 1, 2\}$   
(c)  $\{0, -1, -2\}$   
(d)  $\{-1, 0, 1\}$
16. The system of linear equations  
 $5x + ky = 5,$   
 $3x + 3y = 5;$   
will be consistent if
- (a)  $k \neq -3$  (b)  $k = -5$   
(c)  $k = 5$  (d)  $k \neq 5$

17. The equation of the tangent to the curve  $y(1 + x^2) = 2 - x$ , where it crosses the  $X$ -axis is
- (a)  $x - 5y = 2$  (b)  $5x - y = 2$   
(c)  $x + 5y = 2$  (d)  $5x + y = 2$
18.  $\begin{bmatrix} 3c+6 & a-d \\ a+d & 2-3b \end{bmatrix} = \begin{bmatrix} 12 & 2 \\ -8 & -4 \end{bmatrix}$  are equal, then value of  $\frac{ab-cd}{ab-cd}$  is
- (a) 4 (b) 16  
(c) -4 (d) -16
19. The principal value of  $\tan^{-1}\left(\tan \frac{9\pi}{8}\right)$  is
- (a)  $\frac{\pi}{8}$  (b)  $\frac{3\pi}{8}$   
(c)  $-\frac{\pi}{8}$  (d)  $-\frac{3\pi}{8}$
20. For two matrices  $P = \begin{bmatrix} 3 & 4 \\ -1 & 2 \\ 0 & 1 \end{bmatrix}$  and  $Q^T = \begin{bmatrix} -1 & 2 & 1 \\ 1 & 2 & 3 \end{bmatrix}$   
 $P - Q$  is
- (a)  $\begin{bmatrix} 2 & 3 \\ -3 & 0 \\ 0 & -3 \end{bmatrix}$  (b)  $\begin{bmatrix} 4 & 3 \\ -3 & 0 \\ -1 & -2 \end{bmatrix}$   
(c)  $\begin{bmatrix} 4 & 3 \\ -0 & -3 \\ -1 & -2 \end{bmatrix}$  (d)  $\begin{bmatrix} 2 & 3 \\ 0 & -3 \\ 0 & -3 \end{bmatrix}$

### SECTION - B

In this Section attempt any 16 questions out of the Questions 21-40. Each question is of one mark.

21. The function  $f(x) = 2x^3 - 15x^2 + 36x + 6$  is increasing in the interval
- (a)  $(-\infty, 2) \cup (3, \infty)$  (b)  $(-\infty, 2)$   
(c)  $(-\infty, 2] \cup [3, \infty)$  (d)  $[3, \infty)$
22. If  $x = 2\cos \theta - \cos 2\theta$  and  $y = 2\sin \theta - \sin 2\theta$ , then  $\frac{dy}{dx}$  is
- (a)  $\frac{\cos \theta + \cos 2\theta}{\sin \theta - \sin 2\theta}$  (b)  $\frac{\cos \theta - \cos 2\theta}{\sin 2\theta - \sin \theta}$   
(c)  $\frac{\cos \theta - \cos 2\theta}{\sin \theta - \sin 2\theta}$  (d)  $\frac{\cos 2\theta - \cos \theta}{\sin 2\theta + \sin \theta}$
23. What is the domain of the function  $\cos^{-1}(2x - 3)$ ?
- (a)  $[-1, 1]$  (b)  $(1, 2)$   
(c)  $(-1, 1)$  (d)  $[1, 2]$
24. A matrix  $A = [a_{ij}]_{3 \times 3}$  is defined by
- $$a_{ij} = \begin{cases} 2i + 3j & , i < j \\ 5 & , i = j \\ 3i - 2j & , i > j \end{cases}$$

The number of elements in A which are more than 5, is:

- (a) 3 (b) 4  
(c) 5 (d) 6

25. If a function  $f$  defined by

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3 & \text{if } x = \frac{\pi}{2} \end{cases}$$

is continuous at  $x = \frac{\pi}{2}$ , then the value of  $k$  is

- (a) 2 (b) 3  
(c) 6 (d) -6

26. For the matrix  $X = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$ ,  $(X^2 - X)$  is

- (a)  $2I$  (b)  $3I$   
(c)  $I$  (d)  $5I$

27. Let  $X = \{x^2 : x \in \mathbb{N}\}$  and the function  $f : \mathbb{N} \rightarrow X$  is defined by  $f(x) = x^2, x \in \mathbb{N}$ . Then this function is

- (a) injective only (b) not bijective  
(c) surjective only (d) bijective

28. The corner points of the feasible region for a Linear Programming problem are  $P(0, 5)$ ,  $Q(1, 5)$ ,  $R(4, 2)$  and  $S(12, 0)$ . The minimum value of the objective function  $Z = 2x + 5y$  is at the point

- (a) P (b) Q  
(c) R (d) S

29. The equation of the normal to the curve  $ay^2 = x^3$  at the point  $(am^2, am^3)$  is

- (a)  $2y - 3mx + am^3 = 0$   
(b)  $2x + 3my - 3am^4 - am^2 = 0$   
(c)  $2x + 3my + 3am^4 - 2am = 0$   
(d)  $2x + 3my - 3am^4 - 2am^2 = 0$

30. If A is a square matrix of order 3 and  $|A| = -5$ , then  $|\text{adj } A|$  is

- (a) 125 (b) -25  
(c) 25 (d)  $\pm 25$

31. The simplest form of  $\tan^{-1} \left[ \frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right]$  is

- (a)  $\frac{\pi}{4} - \frac{x}{2}$  (b)  $\frac{\pi}{4} + \frac{x}{2}$   
(c)  $\frac{\pi}{4} - \frac{1}{2} \cos^{-1} x$  (d)  $\frac{\pi}{4} + \frac{1}{2} \cos^{-1} x$

32. If for the matrix  $A = \begin{bmatrix} \alpha & -2 \\ -2 & \alpha \end{bmatrix}$ ,  $|A^3| = 125$ , then the

value of  $\alpha$  is

- (a)  $\pm 3$  (b) -3  
(c)  $\pm 1$  (d) 1

33. If  $y = \sin(m \sin^{-1} x)$ , then which one of the following equations is true?

(a)  $(1-x^2) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + m^2 y = 0$

(b)  $(1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + m^2 y = 0$

(c)  $(1+x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - m^2 y = 0$

(d)  $(1+x^2) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - m^2 x = 0$

34. The principal value of  $[\tan^{-1} \sqrt{3} - \cot^{-1}(-\sqrt{3})]$  is

- (a)  $\pi$  (b)  $-\frac{\pi}{2}$   
(c) 0 (d)  $2\sqrt{3}$

35. The maximum value of  $\left(\frac{1}{x}\right)^x$  is

- (a)  $e^{1/e}$  (b)  $e$   
(c)  $\left(\frac{1}{e}\right)^{1/e}$  (d)  $e^e$

36. Let matrix  $X = [x_{ij}]$  is given by  $X = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & -5 \\ 2 & -1 & 3 \end{bmatrix}$ .

Then the matrix  $Y = [m_{ij}]$ , where  $m_{ij} = \text{Minor of } x_{ij}$ , is

(a)  $\begin{bmatrix} 7 & -5 & -3 \\ 19 & 1 & -11 \\ -11 & 1 & 7 \end{bmatrix}$  (b)  $\begin{bmatrix} 7 & -19 & 11 \\ 5 & -1 & -1 \\ 3 & 11 & 7 \end{bmatrix}$

(c)  $\begin{bmatrix} 7 & 19 & -11 \\ -3 & 11 & 7 \\ -2 & -1 & -1 \end{bmatrix}$  (d)  $\begin{bmatrix} 7 & 19 & -11 \\ -1 & -1 & 1 \\ -3 & -11 & 7 \end{bmatrix}$

37. A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = 2 + x^2$  is

- (a) not one-one  
(b) one-one  
(c) not onto  
(d) neither one-one nor onto

38. A Linear Programming Problem is as follows:

Maximise / Minimise objective function  $Z = 2x - y + 5$

Subject to the constraints

$$3x + 4y \leq 60$$

$$x + 3y \leq 30$$

$$x \geq 0, y \geq 0$$

If the corner points of the feasible region are A(0, 10), B(12, 6), C(20, 0) and O(0,0), then which of the following is true ?

- (a) Maximum value of Z is 40  
 (b) Minimum value of Z is - 5  
 (c) Difference of maximum and minimum values of Z is 35  
 (d) At two corner points, value of Z are equal

39. If  $x = -4$  is a root of  $\begin{vmatrix} x & 2 & 3 \\ 1 & x & 1 \\ 3 & 2 & x \end{vmatrix} = 0$ , then the sum of the other two roots is

- (a) 4 (b) -3  
 (c) 2 (d) 5

40. The absolute maximum value of the function  $f(x) = 4x - \frac{1}{2}x^2$  in the interval  $\left[-2, \frac{9}{2}\right]$  is

- (a) 8 (b) 9  
 (c) 6 (d) 10

### SECTION - C

Attempt any 8 questions out of the Questions 41-50. Each question is of one mark.

41. In a sphere of radius  $r$ , a right circular cone of height  $h$  having maximum curved surface area is inscribed. The expression for the square of curved surface of cone is

- (a)  $2\pi^2rh(2rh + h^2)$  (b)  $\pi^2hr(2rh + h^2)$   
 (c)  $2\pi^2r(2rh^2 - h^3)$  (d)  $2\pi^2r^2(2rh - h^2)$

42. The corner points of the feasible region determined by a set of constraints (linear inequalities) are P(0, 5), Q(3, 5), R(5, 0) and S(4, 1) and the objective function is  $Z = ax + 2by$  where  $a, b > 0$ . The condition on  $a$  and  $b$  such that the maximum Z occurs at Q and S is

- (a)  $a - 5b = 0$  (b)  $a - 3b = 0$   
 (c)  $a - 2b = 0$  (d)  $a - 8b = 0$

43. If curves  $y^2 = 4x$  and  $xy = c$  cut at right angles, then the value of  $c$  is


- (a)  $4\sqrt{2}$  (b) 8  
 (c)  $2\sqrt{2}$  (d)  $-4\sqrt{2}$

44. The inverse of the matrix  $X = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$  is

- (a)  $24 \begin{bmatrix} 1/2 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/4 \end{bmatrix}$  (b)  $\frac{1}{24} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(c)  $\frac{1}{24} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$  (d)  $\begin{bmatrix} 1/2 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/4 \end{bmatrix}$

45. For an L.P.P. the objective function is  $Z = 4x + 3y$ , and the feasible region determined by a set of constraints (linear inequations) is shown in the graph.



Which one of the following statements is true ?

- (a) Maximum value of Z is at R.  
 (b) Maximum value of Z is at Q.  
 (c) Value of Z at R is less than the value at P.  
 (d) Value of Z at Q is less than the value at R.

### Case Study

In a residential society comprising of 100 houses, there were 60 children between the ages of 10-15 years. They were inspired by their teachers to start composting to ensure that biodegradable waste is recycled. For this purpose, instead of each child doing it for only his/her house, children convinced the Residents welfare association to do it as a society initiative. For this they identified a square area in the local park. Local authorities charged amount of ₹ 50 per square metre for space so that there is no misuse of the space and Resident welfare association takes it seriously. Association hired a labourer for digging out 250 m<sup>3</sup> and he charged ₹ 400  $x$  (depth)<sup>2</sup>. Association will like to have minimum cost.



46. Let side of square plot is  $x$  m and its depth is  $h$  metres, then cost  $C$  for the pit is

- (a)  $\frac{50}{h} + 400h^2$                       (b)  $\frac{12500}{h} + 400h^2$   
 (c)  $\frac{250}{h} + h^2$                               (d)  $\frac{250}{h} + 400h^2$

47. Value of  $h$  (in m) for which  $\frac{dC}{dh} = 0$  is

- (a) 1.5    (b) 2  
 (c) 2.5    (d) 3

48.  $\frac{d^2C}{dh^2}$  is given by

- (a)  $\frac{25000}{h^3} + 800$                       (b)  $\frac{500}{h^3} + 800$   
 (c)  $\frac{100}{h^3} + 800$                           (d)  $\frac{500}{h^3} + 2$

49. Value of  $x$  (in m) for minimum cost is

- (a) 5    (b)  $10\sqrt{\frac{5}{3}}$   
 (c)  $5\sqrt{5}$                                       (d) 10

50. Total minimum cost of digging the pit (in ₹) is

- (a) 4,100    (b) 7,500  
 (c) 7,850    (d) 3,220

□□□

## SOLUTIONS

### SECTION - A

1. (a)  $\frac{5}{x \log(x^5) \log(\log x^5)}$

*Explanation:* Let  $y = \log[\log(\log x^5)]$

$$\begin{aligned} \therefore \frac{dy}{dx} &= \frac{1}{\log(\log x^5)} \frac{dy}{dx} [\log(\log x^5)] \\ &\quad \text{(By Chain Rule)} \\ &= \frac{1}{\log(\log x^5)} \cdot \frac{1}{\log x^5} \frac{d}{dx} \log x^5 \\ &= \frac{1}{\log(x^5) \log(\log x^5)} \frac{1}{x^5} \frac{d}{dx} (x^5) \\ &= \frac{5}{x \log(x^5) \log(\log x^5)} \end{aligned}$$

2. (c) 64

*Explanation:* The order of the matrix =  $2 \times 3$   
 The number of elements =  $2 \times 3 = 6$

Each place can have either 1 or 2. So, each place can be filled in 2 ways.

Thus, the number of possible matrices =  $2^6 = 64$

3. (d) neither maximum value nor minimum value

*Explanation:* Given,  $f(x) = x^3 + 1$

$$\therefore f'(x) = 3x^2 \text{ and } f''(x) = 6x$$

Put  $f'(x) = 0$

$$\Rightarrow 3x^2 = 0 \Rightarrow x = 0$$

At  $x = 0, f''(x) = 0$

Thus,  $f(x)$  has neither maximum value nor minimum value.

4. (a)  $\frac{\cos a}{\cos^2(a+y)}$

*Explanation:* Given,  $\sin y = x \cos(a+y)$

$$\Rightarrow x = \frac{\sin y}{\cos(a+y)}$$

Differentiating with respect to  $y$ , we get

$$\frac{dx}{dy} = \frac{\cos(a+y) \frac{d}{dy}(\sin y) - \sin y \frac{d}{dy} \{\cos(a+y)\}}{\cos^2(a+y)}$$

$$\Rightarrow \frac{dx}{dy} = \frac{\cos(a+y) \cos y - \sin y [-\sin(a+y)]}{\cos^2(a+y)}$$

$$\Rightarrow \frac{dx}{dy} = \frac{\cos(a+y) \cos y + \sin y \sin(a+y)}{\cos^2(a+y)}$$

$$\Rightarrow \frac{dx}{dy} = \frac{\cos[(a+y) - y]}{\cos^2(a+y)}$$

$$\Rightarrow \frac{dx}{dy} = \frac{\cos a}{\cos^2(a+y)}$$

5. (b)  $(0, \pm 5)$

*Explanation:* The equation of the given curve:

$$\frac{x^2}{9} + \frac{y^2}{25} = 1 \dots(i)$$

On differentiating both sides w.r.t.  $x$ , we get

$$\frac{2x}{9} + \frac{2y}{25} \frac{dy}{dx} = 1$$

$$\Rightarrow \frac{dy}{dx} = \frac{-25x}{9y}$$

Since, tangent is parallel to  $X$ -axis, then the slope of the tangent is zero.

$$\therefore \frac{-25x}{9y} = 0, \text{ which is possible if } x = 0$$

Put  $x = 0$  in eq (i), we get

$$\frac{y^2}{25} = 1 \Rightarrow y^2 = 25 \Rightarrow y = \pm 5$$

Hence, required points are  $(0, \pm 5)$ .

6. (d) 1

**Explanation:** As points are collinear

$\Rightarrow$  area of triangle formed by 3 points is zero.

$$\Rightarrow \frac{1}{2} \begin{vmatrix} x_1 - x_2 & x_2 - x_3 \\ y_1 - y_2 & y_2 - y_3 \end{vmatrix} = 0$$

$$\Rightarrow \frac{1}{2} \begin{vmatrix} 2x - 0 & 0 - (x + 3) \\ (x + 3) - x & x - (x + 6) \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 2x & -(x + 3) \\ 3 & -6 \end{vmatrix} = 0$$

$$\Rightarrow -12x + 3(x + 3) = 0$$

$$\Rightarrow -12x + 3x + 9 = 0$$

$$\Rightarrow -9x = -9$$

$$\Rightarrow x = 1$$

7. (a)  $\frac{\pi}{12}$

**Explanation:**  $\cos^{-1}\left(\frac{1}{2}\right) + \sin^{-1}\left(\frac{-1}{\sqrt{2}}\right)$

$$= \cos^{-1}\left(\cos\frac{\pi}{3}\right) - \sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$$

$$= \frac{\pi}{3} - \sin^{-1}\left(\sin\frac{\pi}{4}\right)$$

$$= \frac{\pi}{3} - \frac{\pi}{4} = \frac{\pi}{12}$$

8. (c)  $\frac{y - 4x(x^2 + y^2)}{4y(x^2 + y^2) - x}$

**Explanation:** Given,  $(x^2 + y^2)^2 = xy$

$$\Rightarrow x^4 + 2x^2y^2 + y^4 - xy = 0$$

Differentiating w.r.t.  $x$ , we get

$$4x^3 + 2 \left[ 2xy^2 + x^2 \cdot 2y \frac{dy}{dx} \right] + 4y^3 \frac{dy}{dx} - \left[ y + x \frac{dy}{dx} \right] = 0$$

$$\frac{dy}{dx} [4x^2y + 4y^3 - x] + [4x^3 + 4xy^2 - y] = 0$$

$$\frac{dy}{dx} = \frac{-(4x^3 + 4xy^2 - y)}{[4x^2y + 4y^3 - x]}$$

$$\text{or } \frac{dy}{dx} = \frac{y - 4x(x^2 + y^2)}{4y(x^2 + y^2) - x}$$

9. (b) Zero square matrix

**Explanation:** If matrix A is symmetric

$$A^T = A$$

If matrix A is skew-symmetric

$$A^T = -A$$

Also, diagonal elements are zero.

Since, it is given that matrix A is both symmetric and skew-symmetric.

$$\therefore A = A^T = -A$$

Which is only possible if A is zero matrix.

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = A^T = -A$$

Thus, if a matrix A is both symmetric and skew-symmetric, then A is necessarily a zero matrix.

10. (c)  $\{(1, 1), (3, 3), (3, 1), (2, 3)\}$

**Explanation:**

(i) R is reflexive if it contains  $\{(1, 1), (2, 2) \text{ and } (3, 3)\}$ .

Since,  $(2, 2) \in R$ . So, we need to add  $(1, 1)$  and  $(2, 2)$  to make R reflexive.


(ii) R is symmetric if it contains  $\{(2, 2), (1, 3), (3, 1), (3, 2), (2, 3)\}$ .

Since,  $\{(2, 2), (1, 3), (3, 2)\} \in R$ . So, we need to add  $(3, 1)$  and  $(2, 3)$ .

Thus, minimum ordered pairs which should be added in relation R to make it reflexive and symmetric are  $\{(1, 1), (3, 3), (3, 1), (2, 3)\}$ .

11. (b) 5 corner points including  $(7, 7)$  and  $(3, 3)$

**Explanation:** On plotting the constraints  $x = 3$ ,  $x = 9$ ,  $x = y$  and  $x + y = 14$ , we get the following graph. From the graph given below it clear that feasible region is ABCDEA, including corner points A(9, 0), B(3, 0), C(3, 3), D(7, 7) and E(9, 5). Thus feasible region has 5 corner points including  $(7, 7)$  and  $(3, 3)$ .



12. (d) 8

**Explanation:** Since,  $f(x)$  is continuous at  $x = 0$ , then

$$\text{LHL} = \text{RHL} = f(0) \quad \text{or} \quad \text{LHL} = \text{RHL} = k$$

$$\begin{aligned} \text{Now, LHL} &= \lim_{h \rightarrow 0} \frac{e^{3(0-h)} - e^{-5(0-h)}}{0-h} \\ &= \lim_{h \rightarrow 0} \frac{e^{-3h} - e^{5h}}{-h} \\ &= \lim_{h \rightarrow 0} \left( \frac{e^{-3h} - 1}{-h} \right) + \lim_{h \rightarrow 0} \left( \frac{e^{5h} - 1}{h} \right) \\ &= 3 \lim_{h \rightarrow 0} \left( \frac{e^{-3h} - 1}{-3h} \right) + 5 \lim_{h \rightarrow 0} \left( \frac{e^{5h} - 1}{5h} \right) \\ &= 3 \times 1 + 5 \times 1 = 8 \end{aligned}$$

Thus,  $k = 8$ .

13. (a) 5

**Explanation:**

$$\text{Here, } C_{31} = (-1)^{3+1} \begin{vmatrix} -1 & 2 \\ 2 & -3 \end{vmatrix} = 3 - 4 = -1$$

$$\text{and } C_{23} = (-1)^{2+3} \begin{vmatrix} 1 & -1 \\ 3 & 2 \end{vmatrix} = -(2 + 3) = -5$$

$$\text{Thus, } C_{31} \cdot C_{23} = (-1)(-5) = 5$$

14. (d)  $(-\infty, 0) \cup (2, \infty)$ .**Explanation:** We have,

$$f(x) = y = x^2 e^{-x}$$

$$\therefore \frac{dy}{dx} = 2x e^{-x} + x^2(-1)e^{-x} = x e^{-x}(2-x)$$



$$\text{Now, put } \frac{dy}{dx} = 0$$

$$\Rightarrow x = 0 \text{ and } x = 2$$

The points  $x = 0$  and  $x = 2$  divide the real line into three disjoint intervals i.e.,  $(-\infty, 0)$ ,  $(0, 2)$  and  $(2, \infty)$

In intervals,  $(-\infty, 0)$  and  $(2, \infty)$ ,  $f'(x) < 0$  as  $e^{-x}$  is always positive.

$\therefore f(x)$  or  $y$  is decreasing in  $(-\infty, 0)$  and  $(2, \infty)$ .

15. (b)  $\{-2, -1, 0, 1, 2\}$ **Explanation:** Given,  $R = \{(x, y) : x, y \in Z, x^2 + y^2 \leq 4\}$ 

$$\text{Let } x = 0, \text{ then } y^2 \leq 4 \Rightarrow y = 0, \pm 1, \pm 2$$

Thus, domain of  $R = \{-2, -1, 0, 1, 2\}$ 16. (d)  $k \neq 5$ **Explanation:** We have,  $5x + ky - 5 = 0$ 

$$\text{and } 3x + 3y - 5 = 0$$

For consistent system

$$\frac{5}{3} \neq \frac{k}{3}$$

$$\Rightarrow k \neq 5$$

17. (c)  $x + 5y = 2$ **Explanation:** Given,  $y(1 + x^2) = 2 - x$  ... (i)

If it cuts X-axis, then y-coordinate is 0.

$$\therefore 0(1 + x^2) = 2 - x$$

$$\Rightarrow x = 2$$

Thus, point of contact is  $(2, 0)$ .Now, differentiating eq.(i) w.r.t.  $x$ , we get

$$y \cdot (2x) + \frac{dy}{dx} (1 + x^2) = -1$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1 - 2xy}{1 + x^2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-(1 + 2xy)}{1 + x^2}$$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{(2,0)} = \frac{-1}{5}$$

Thus, equation of tangent is

$$y - 0 = \frac{-1}{5}(x - 2)$$

$$\Rightarrow 5y + x = 2$$

$$\Rightarrow x + 5y = 2$$

18. (a) 4

**Explanation:** Given,  $\begin{bmatrix} 3c+6 & a-d \\ a+d & 2-3d \end{bmatrix} = \begin{bmatrix} 12 & 2 \\ -8 & -4 \end{bmatrix}$ 

$$\therefore 3c + 6 = 12 \quad \dots \text{(i)}$$

$$a - d = 2 \quad \dots \text{(ii)}$$

$$a + d = -8 \quad \dots \text{(iii)}$$

$$2 - 3d = -4 \quad \dots \text{(iv)}$$

From eq. (i), we get  $c = 2$ 

On solving eqs. (ii) and (iii), we get  $a = -3$  and  $d = -5$  from eq. (iv), we get  $b = 2$

$$\text{Now, } ab - cd = (-3)2 - 2(-5)$$

$$\Rightarrow ab - cd = -6 + 10 = 4$$

19. (a)  $\frac{\pi}{8}$

**Explanation:**  $\tan^{-1}\left(\tan\frac{9\pi}{8}\right) = \tan^{-1}\left(\tan\left(\pi + \frac{\pi}{8}\right)\right)$   
 $= \tan^{-1}\left(\tan\frac{\pi}{8}\right) = \frac{\pi}{8} \quad \left[\because \frac{\pi}{8} \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)\right]$

20. (b)  $\begin{bmatrix} 4 & 3 \\ -3 & 0 \\ -1 & -2 \end{bmatrix}$

**Explanation:**

Here,  $Q = (Q^T)^T = \begin{bmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{bmatrix}$

Now,  $P - Q = \begin{bmatrix} 3 & 4 \\ -1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} -1 & 1 \\ 2 & 2 \\ 1 & 3 \end{bmatrix}$

$$= \begin{bmatrix} 4 & 3 \\ -3 & 0 \\ -1 & -2 \end{bmatrix}$$

**SECTION-B**


21. (c)  $(-\infty, 2] \cup [3, \infty)$

**Explanation:** Given,  $f(x) = 2x^3 - 15x^2 + 36x + 6$

$\therefore f'(x) = 6x^2 - 30x + 36$

It  $f'(x) \geq 0$ , then  $f(x)$  is increasing.

So,  $6x^2 - 30x + 36 \geq 0$



or,  $x^2 - 5x + 6 \geq 0$

or,  $(x-3)(x-2) \geq 0$

$\therefore x \in (-\infty, 2] \cup [3, \infty)$

22. (b)  $\frac{\cos\theta - \cos 2\theta}{\sin 2\theta - \sin\theta}$

**Explanation:** Given,  $x = 2\cos\theta - \cos 2\theta$

and  $y = 2\sin\theta - \sin 2\theta$

Therefore,  $\frac{dx}{d\theta} = -2\sin\theta + 2\sin 2\theta$

and  $\frac{dy}{d\theta} = 2\cos\theta - 2\cos 2\theta$

$$\therefore \frac{dy}{dx} = \frac{2\cos\theta - 2\cos 2\theta}{-2\sin\theta + 2\sin 2\theta}$$

or  $\frac{dy}{dx} = \frac{\cos\theta - \cos 2\theta}{\sin 2\theta - \sin\theta}$

23. (d) [1, 2]

**Explanation:** Let,  $f(x) = \cos^{-1}(2x-3)$

$\therefore -1 \leq 2x-3 \leq 1$

$\Rightarrow 2 \leq 2x \leq 4$

$\Rightarrow 1 \leq x \leq 2$

$\therefore x \in [1, 2]$  or domain of  $x$  is  $[1, 2]$ .

24. (b) 4

**Explanation:** Here,  $A = \begin{bmatrix} 5 & 8 & 11 \\ 4 & 5 & 13 \\ 7 & 5 & 5 \end{bmatrix}$

Thus, number of elements more than 5, is 4.

25. (c) 6

**Explanation:** Since,  $f(x)$  is continuous at  $x = \frac{\pi}{2}$

Therefore,  $\lim_{x \rightarrow \frac{\pi}{2}} f(x) = f\left(\frac{\pi}{2}\right)$

$\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x} = 3$

$\Rightarrow k \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin\left(\frac{\pi}{2} - x\right)}{2\left(\frac{\pi}{2} - x\right)} = 3$

$\Rightarrow \frac{k}{2} \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin\left(\frac{\pi}{2} - x\right)}{\left(\frac{\pi}{2} - x\right)} = 3$

$\Rightarrow \frac{k}{2} \times 1 = 3 \Rightarrow k = 6$

26. (a) 21

**Explanation:**

Here  $X^2 = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

$\Rightarrow X^2 = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$



$$\Rightarrow X^2 = 2 \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} = 2I$$

27. (a) injective only

**Explanation:** Let  $x_1, x_2 \in \mathbb{N}$

$$f(x_1) = f(x_2)$$

$$\Rightarrow x_1^2 = x_2^2$$

$$\Rightarrow x_1^2 - x_2^2 = 0$$

$$\Rightarrow (x_1 + x_2)(x_1 - x_2) = 0$$

$$\Rightarrow x_1 = x_2$$

$$\{x_1 + x_1 \neq 0 \text{ as } x_1, x_2 \in \mathbb{N}\}$$

Hence,  $f(x)$  is injective.

Also, the elements like 2 and 3 have no pre-image in  $\mathbb{N}$ . Thus,  $f(x)$  is not surjective.

28. (c) R

**Explanation:**

Corner Points	Value of $Z = 2x + 5y$
P(0, 5)	$Z = 2(0) + 5(5) = 25$
Q(1, 5)	$Z = 2(1) + 5(5) = 27$
R(4, 2)	$Z = 2(4) + 5(2) = 18$ ® Minimum
S(12, 0)	$Z = 2(12) + 5(0) = 24$

Thus, minimum value of  $Z$  occurs at R(4, 2).

29. (d)  $2x + 3my - 3am^4 - 2am^2 = 0$

**Explanation:**  $ay^2 = x^3$   
Differentiating both sides w.r.t.  $x$ , we get

$$2ay \frac{dy}{dx} = 3x^2$$

$$\Rightarrow \frac{dy}{dx} = \frac{3x^2}{2ay}$$

Slope of the tangent at  $(am^2, am^3)$  is

$$\left(\frac{dy}{dx}\right)_{(am^2, am^3)} = \frac{3(am^2)^2}{2a(am^3)}$$

$$= \frac{3a^2m^4}{2a^2m^3} = \frac{3m}{2}$$

Now, slope of normal at  $(am^2, am^3) = \frac{-1}{\left(\frac{dy}{dx}\right)_{(am^2, am^3)}}$

$$= \frac{-2}{3m}$$

Thus, equation of normal at  $(am^2, am^3)$  is

$$(y - am^3) = \frac{-2}{3m}(x - am^2)$$

$$\Rightarrow 3my - 3am^4 + 2x - 2am^2 = 0$$

$$\Rightarrow 2x + 3my - 3am^4 - 2am^2 = 0$$

30. (c) 25

**Explanation:** We know that,

$$|\text{adj } A| = |A|^{n-1}$$

where  $n$  is the order of the matrix

$$\therefore |\text{adj } A| = (5)^{3-1}$$

$$= 5^2 = 25$$

31. (c)  $\frac{\pi}{4} - \frac{1}{2} \cos^{-1} x$

**Explanation:** We have,

$$\tan^{-1} \left( \frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right)$$

Put  $x = \cos 2\theta$ , so that  $\theta = \frac{1}{2} \cos^{-1} x$

$$\tan^{-1} \left( \frac{\sqrt{1+\cos 2\theta} - \sqrt{1-\cos 2\theta}}{\sqrt{1+\cos 2\theta} + \sqrt{1-\cos 2\theta}} \right)$$

$$= \tan^{-1} \left( \frac{\sqrt{2\cos^2 \theta} - \sqrt{2\sin^2 \theta}}{\sqrt{2\cos^2 \theta} + \sqrt{2\sin^2 \theta}} \right)$$

$$= \tan^{-1} \left( \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta} \right)$$

$$= \tan^{-1} \left( \frac{1 - \tan \theta}{1 + \tan \theta} \right)$$

$$= \tan^{-1}(1) - \tan^{-1}(\tan \theta)$$

$$\left[ \because \tan^{-1} \left( \frac{x-y}{1+xy} \right) = \tan^{-1} x - \tan^{-1} y \right]$$

$$= \tan^{-1} \left( \tan \frac{\pi}{4} \right) - \theta$$

$$= \frac{\pi}{4} - \frac{1}{2} \cos^{-1} x$$

32. (a)  $\pm 3$

**Explanation:** Given,  $A = \begin{bmatrix} \alpha & -2 \\ -2 & \alpha \end{bmatrix}$

$$\Rightarrow |A| = \alpha^2 - 4 \quad \dots(i)$$

Also, given  $|A^3| = 125$

$$\begin{aligned} \Rightarrow & |A|^3 = 125 \\ \Rightarrow & |A| = 5 \\ \Rightarrow & \alpha^2 - 4 = 5 \quad [\text{from eq. (i)}] \\ \Rightarrow & \alpha^2 = 9 \\ \Rightarrow & \alpha = \pm 3 \end{aligned}$$

33. (b)  $(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + m^2y = 0$

**Explanation:** Given,  $y = \sin(m\sin^{-1}x)$  ... (i)

Differentiating both sides w.r.t.  $x$ , we get

$$\frac{dy}{dx} = \cos(m\sin^{-1}x) \times \frac{m}{\sqrt{1-x^2}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{m\cos(m\sin^{-1}x)}{\sqrt{1-x^2}} \quad \dots(\text{ii})$$

$$\Rightarrow y' = \frac{m\cos(m\sin^{-1}x)}{\sqrt{1-x^2}} \quad \dots(\text{ii})$$

$$\Rightarrow (\sqrt{1-x^2})y' = m\cos(m\sin^{-1}x)$$

Differentiating again w.r.t. ' $x$ ', we get

$$\begin{aligned} y''(\sqrt{1-x^2}) + y' \frac{(-2x)}{2\sqrt{1-x^2}} \\ = -m^2 \sin(m\sin^{-1}x) \frac{1}{\sqrt{1-x^2}} \end{aligned}$$

$$\Rightarrow y''(1-x^2) - xy' = -m^2y$$

$$\Rightarrow y''(1-x^2) - xy' + m^2y = 0$$

or,  $(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + m^2y = 0$

34. (b)  $-\frac{\pi}{2}$

**Explanation:** We have,

$$\begin{aligned} \tan^{-1}\sqrt{3} - \cot^{-1}(-\sqrt{3}) \\ = \tan^{-1}\left(\tan\frac{\pi}{3}\right) - \pi + \cot^{-1}\cot\sqrt{3} \\ = \frac{\pi}{3} - \left(\pi - \frac{\pi}{6}\right) \\ = -\frac{\pi}{2} \end{aligned}$$

35. (a)  $e^{1/e}$

**Explanation:** Let  $y = \left(\frac{1}{x}\right)^x$

Then,  $\log y = x \log\left(\frac{1}{x}\right) = -x \log x \quad \dots(\text{i})$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \therefore \frac{1}{y} \frac{dy}{dx} &= -\left[x \cdot \frac{1}{x} + \log x\right] \\ &= -(1 + \log x) \quad \dots(\text{ii}) \end{aligned}$$

On differentiating again eq. (ii), we get

$$\begin{aligned} \frac{1}{y} \frac{d^2y}{dx^2} - \frac{1}{y^2} \left(\frac{dy}{dx}\right)^2 \\ = \frac{-1}{x} \quad \dots(\text{iii}) \end{aligned}$$

From eq. (ii), we get

$$\begin{aligned} \frac{dy}{dx} &= -y(1 + \log x) \\ &= -\left(\frac{1}{x}\right)^x (1 + \log x) \end{aligned}$$

For maximum or minimum values of  $y$ , put  $\frac{dy}{dx} = 0$

Therefore,  $\left(\frac{1}{x}\right)^x (1 + \log x) = 0$

However,  $\left(\frac{1}{x}\right)^x \neq 0$  for any value of  $x$ . Therefore

$$1 + \log x = 0$$

$$\Rightarrow \log x = -1 \Rightarrow x = e^{-1} \Rightarrow x = \frac{1}{e}$$

When  $x = \frac{1}{e}$ , from eq. (iii)

$$\frac{1}{y} \frac{d^2y}{dx^2} - 0 = -e$$

$$\Rightarrow \frac{d^2y}{dx^2} = -e(e)^{1/e} < 0$$

Hence,  $y$  is maximum when  $x = \frac{1}{e}$  and maximum value of  $y = e^{1/e}$ .

36. (d)  $\begin{bmatrix} 7 & 19 & -11 \\ -1 & -1 & 1 \\ -3 & -11 & 7 \end{bmatrix}$

**Explanation:**  $m_{11} = \begin{vmatrix} 4 & -5 \\ -1 & 3 \end{vmatrix} = 12 - 5 = 7$

$$m_{12} = \begin{vmatrix} 3 & -5 \\ 2 & 3 \end{vmatrix} = 9 + 10 = 19$$

$$m_{13} = \begin{vmatrix} 3 & 4 \\ 2 & -1 \end{vmatrix} = -3 - 8 = -11$$

$$m_{21} = \begin{vmatrix} -1 & 2 \\ -1 & 3 \end{vmatrix} = -3 + 2 = -1$$

$$m_{22} = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = 3 - 4 = -1$$

$$m_{23} = \begin{vmatrix} 1 & -1 \\ 2 & -1 \end{vmatrix} = -1 + 2 = 1$$

$$m_{31} = \begin{vmatrix} -1 & 2 \\ 4 & -5 \end{vmatrix} = 5 - 8 = -3$$

$$m_{32} = \begin{vmatrix} 1 & 2 \\ 3 & -5 \end{vmatrix} = -5 - 6 = -11$$

$$m_{33} = \begin{vmatrix} 1 & -1 \\ 3 & 4 \end{vmatrix} = 4 + 3 = 7$$

$$\therefore Y = \begin{bmatrix} 7 & 19 & -11 \\ -1 & -1 & 1 \\ -3 & -11 & 7 \end{bmatrix}$$

**37. (d) neither one-one nor onto**

**Explanation:** Given,  $f(x) = 2 + x^2$

For one-one,  $f(x_1) = f(x_2)$

$$\Rightarrow 2 + x_1^2 = 2 + x_2^2$$

$$\Rightarrow x_1^2 = x_2^2$$

$$\Rightarrow x_1 = \pm x_2$$

$$\Rightarrow x_1 = x_2$$

or  $x_1 = -x_2$

Thus,  $f(x)$  is not one-one.

For onto

Let  $f(x) = y$  such that  $y \in R$

$$\therefore x^2 = y - 2$$

$$\Rightarrow x = \pm\sqrt{y-2}$$

Put  $y = -3$ , we get

$$x = \pm\sqrt{-3-2} = \pm\sqrt{-5}$$

Which is not possible as root of negative is not a real number.

Hence,  $x$  is not real.

So,  $f(x)$  is not onto.

**38. (b) Minimum value of Z is -5**

**Explanation:**

Corner Points	Value of $Z = 2x - y + 5$
A(0, 10)	$Z = 2(0) - 10 + 5 = -5$ (Minimum)
B(12, 6)	$Z = 2(12) - 6 + 5 = 23$
C(20, 0)	$Z = 2(20) - 0 + 5 = 45$ (Maximum)
O(0, 0)	$Z = 0(0) - 0 + 5 = 5$

So the minimum value of Z is -5.

**39. (a) 4**

**Explanation:** Given,  $\begin{vmatrix} x & 2 & 3 \\ 1 & x & 1 \\ 3 & 2 & x \end{vmatrix} = 0$

$$\Rightarrow x(x^2 - 2) - 2(x - 3) + 3(2 - 3x) = 0$$

$$\Rightarrow x^3 - 13x + 13 = 0$$

Since  $(x + 4)$  is one root of above cubic equation.

$$\text{Sum roots} = 0$$

$$\therefore \text{Sum of two roots} + (-4) = 0$$

$$\text{Sum of two roots} = 4$$

**40. (a) 8**

**Explanation:** Given,  $f(x) = 4x - \frac{1}{2}x^2$

$$\therefore f'(x) = 4 - \frac{1}{2}(2x) = 4 - x$$

put  $f'(x) = 0$

$$\Rightarrow 4 - x = 0$$

$$\Rightarrow x = 4$$

Then, we evaluate the  $f$  at critical point  $x = 4$  and at the end points of the interval  $\left[-2, \frac{9}{2}\right]$ .

$$f(4) = 16 - \frac{1}{2}(16) = 16 - 8 = 8$$

$$f(-2) = -8 - \frac{1}{2}(4)$$


$$= -8 - 2 = -10$$

$$f\left(\frac{9}{2}\right) = 4\left(\frac{9}{2}\right) - \frac{1}{2}\left(\frac{9}{2}\right)^2$$

$$= 18 - \frac{81}{8} = 7.875$$

Thus, the absolute maximum value of  $f$  on  $\left[-2, \frac{9}{2}\right]$  is

8 occurring at  $x = 4$ .

**SECTION - C**41. (c)  $2\pi^2 r(2rh^2 - h^3)$ **Explanation:**Here, CSA of cone =  $\pi Rl$ Radius of sphere =  $r$ height of cone =  $h$ In  $\triangle AOC$ ,

$$AO^2 = AC^2 + OC^2$$

$$\Rightarrow r^2 = R^2 + (h-r)^2$$

$$\Rightarrow R^2 = 2hr - h^2$$

$$\therefore \text{Radius of cone, } R = \sqrt{2hr - h^2} \quad \dots(i)$$

In  $\triangle ABC$ ,

$$AB^2 = AC^2 + BC^2$$

$$\Rightarrow l^2 = R^2 + h^2$$

$$\Rightarrow l^2 = 2hr - h^2 + h^2$$

$$\therefore \text{slant height} = \sqrt{2hr} \quad \dots(ii)$$

CSA of cone =  $\pi Rl$ 

$$= \pi \sqrt{2hr - h^2} \sqrt{2hr}$$

$$(\text{CSA of cone})^2 = \pi^2 (2hr - h^2)(2hr)$$

$$= 2\pi^2 hr(2hr - h^2)$$

$$= 2\pi^2 r(2rh^2 - h^3)$$

42. (d)  $a - 8b = 0$ **Explanation:** Given, Max.  $Z = ax + 2by$ Max. value of  $Z$  on  $Q(3, 5) = \text{Max. value of } Z \text{ on } S(4, 1)$ 

$$\Rightarrow 3a + 10b = 4a + 2b$$

$$\Rightarrow a - 8b = 0$$

43. (a)  $4\sqrt{2}$ **Explanation:** Given curves,  $y^2 = 4x$  and  $xy = c$  cuts orthogonally.Let them intersect at  $(x_1, y_1)$ .

Now,

$$y^2 = 4x$$

$$\therefore 2y \frac{dy}{dx} = 4$$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{y}$$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{(x_1, y_1)} = \frac{2}{y_1} \quad \dots(i)$$

and  $xy = c$ 

$$\therefore x \frac{dy}{dx} + y = 0$$

$$\therefore \frac{dy}{dx} = \frac{-y}{x}$$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{(x_1, y_1)} = \frac{-y_1}{x_1} \quad \dots(ii)$$

From eqs. (i) and (ii)

$$\frac{2}{y_1} \times \left( \frac{-y_1}{x_1} \right) = -1 \quad [\because m_1 m_2 = -1]$$

$$\Rightarrow x_1 = 2$$

Put  $x_1 = 2$  in  $y_1^2 = 4x_1$ , we get

$$y_1^2 = 4(2) = 8$$

$$\Rightarrow y_1 = 2\sqrt{2}$$

Now, put value of  $x_1$  and  $y_1$  in  $x_1 y_1 = c$ , we get

$$c = 2(2\sqrt{2}) = 4\sqrt{2}$$

$$44. (d) \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$$

**Explanation:** The inverse of a diagonal matrix is obtained by replacing each element in the diagonal with its reciprocal.

$$\text{Since, } X = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

$$\text{Therefore } X^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$$

45. (b) Maximum value of  $Z$  is at  $Q$ .

*Explanation:*

Corner Points of Feasible Region	Value of $z = (z = 4x + 3y)$
O(0, 0)	$Z = 4(0) + 3(0) = 0$
P(0, 40)	$Z = 4(0) + 3(40) = 120$
Q(30, 20)	$Z = 4(30) + 3(20) = 180$ (Maximum)
R(40, 0)	$Z = 4(40) + 3(0) = 160$

Thus, Maximum value of  $z$  is at  $Q$ , which is 180.

46. (b)  $\frac{12500}{h} + 400h^2$

*Explanation:*  $C = \frac{250 \times 50}{h} + 400 \times h^2$

$$\Rightarrow C = \frac{12500}{h} + 400h^2$$

47. (c) 2.5 m

*Explanation:* Since,

$$C = \frac{12500}{h} + 400h^2$$

$$\therefore \frac{dC}{dh} = \frac{-12500}{h^2} + 800h$$

Put  $\frac{dC}{dh} = 0$

$$\therefore \frac{-12500}{h^2} + 800h = 0$$

$$\Rightarrow 800h^3 = 12500$$

$$\Rightarrow h^3 = \frac{125}{8}$$

$$\Rightarrow h = \sqrt[3]{\frac{125}{8}} = \frac{5}{2} = 2.5 \text{ m}$$

48. (a)  $\frac{25000}{h^3} + 800$

*Explanation:* Since,

$$\therefore \frac{dC}{dh} = \frac{-12500}{h^2} + 800h$$

$$\therefore \frac{d^2C}{dh^2} = \frac{-(-2) \times 12500}{h^3} + 800$$

$$\Rightarrow \frac{d^2C}{dh^2} = \frac{25000}{h^3} + 800$$

49. (d) 10 m

*Explanation:* For minimum cost, put  $\frac{dC}{dh} = 0$ , we get

$$h = 2.5 \text{ m}$$

At  $h = 2.5$ ,  $\frac{d^2C}{dh^2} > 0$

(Hence, minimum)

Value of  $x$  at minimum cost

$$x = \frac{400 \times (2.5)^2}{250}$$

$$= \frac{2500}{250} = 10 \text{ m}$$

50. (b) 7,500

*Explanation:* Total minimum cost,

$$C = \frac{12500}{h} + 400h^2 \quad (\text{At } 2.5)$$

$$\Rightarrow C = \frac{12500}{2.5} + 400(2.5)^2$$

$$\Rightarrow C = 5000 + 2500$$

$$\Rightarrow C = ₹7500$$

□□□

Term – I

OMR SHEET

Booklet Series

A

Use English Numbers / Letters only. Use Blue / Black Ball Point Pen to write in box.

Booklet Series

- (A)
- (B)
- (C)
- (D)

Roll Number

- (0) (0) (0) (0) (0) (0) (0) (0) (0)
- (1) (1) (1) (1) (1) (1) (1) (1) (1)
- (2) (2) (2) (2) (2) (2) (2) (2) (2)
- (3) (3) (3) (3) (3) (3) (3) (3) (3)
- (4) (4) (4) (4) (4) (4) (4) (4) (4)
- (5) (5) (5) (5) (5) (5) (5) (5) (5)
- (6) (6) (6) (6) (6) (6) (6) (6) (6)
- (7) (7) (7) (7) (7) (7) (7) (7) (7)
- (8) (8) (8) (8) (8) (8) (8) (8) (8)
- (9) (9) (9) (9) (9) (9) (9) (9) (9)

Name

Test Date

Invigilator's Signature

Student's Signature

Certified that all the entries in this section have been properly filled by the student.

Proper Marking

The OMR Sheet will be computer checked. Fill the circles completely and dark enough for proper detection. Use ballpen (black or blue) for marking.

- (A) (B) (C) (D)

Avoid Improper Marking

- Partially Filled

- Lightly Filled

Test Center Code

- (0) (0)
- (1) (1)
- (2) (2)
- (3) (3)
- (4) (4)
- (5) (5)
- (6) (6)
- (7) (7)
- (8) (8)
- (9) (9)

Subject

IMPORTANT

The candidate should check that the Test Book Series printed on the OMR Sheet is the same as printed on the Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the invigilator for replacement of both the Test Booklet and the Answer Sheet.

Darken the circle for each question.

Q.No.	Response	Q.No.	Response	Q.No.	Response	Q.No.	Response
01	(A) (B) (C) (D)	16	(A) (B) (C) (D)	31	(A) (B) (C) (D)	46	(A) (B) (C) (D)
02	(A) (B) (C) (D)	17	(A) (B) (C) (D)	32	(A) (B) (C) (D)	47	(A) (B) (C) (D)
03	(A) (B) (C) (D)	18	(A) (B) (C) (D)	33	(A) (B) (C) (D)	48	(A) (B) (C) (D)
04	(A) (B) (C) (D)	19	(A) (B) (C) (D)	34	(A) (B) (C) (D)	49	(A) (B) (C) (D)
05	(A) (B) (C) (D)	20	(A) (B) (C) (D)	35	(A) (B) (C) (D)	50	(A) (B) (C) (D)
06	(A) (B) (C) (D)	21	(A) (B) (C) (D)	36	(A) (B) (C) (D)	51	(A) (B) (C) (D)
07	(A) (B) (C) (D)	22	(A) (B) (C) (D)	37	(A) (B) (C) (D)	52	(A) (B) (C) (D)
08	(A) (B) (C) (D)	23	(A) (B) (C) (D)	38	(A) (B) (C) (D)	53	(A) (B) (C) (D)
09	(A) (B) (C) (D)	24	(A) (B) (C) (D)	39	(A) (B) (C) (D)	54	(A) (B) (C) (D)
10	(A) (B) (C) (D)	25	(A) (B) (C) (D)	40	(A) (B) (C) (D)	55	(A) (B) (C) (D)
11	(A) (B) (C) (D)	26	(A) (B) (C) (D)	41	(A) (B) (C) (D)	56	(A) (B) (C) (D)
12	(A) (B) (C) (D)	27	(A) (B) (C) (D)	42	(A) (B) (C) (D)	57	(A) (B) (C) (D)
13	(A) (B) (C) (D)	28	(A) (B) (C) (D)	43	(A) (B) (C) (D)	58	(A) (B) (C) (D)
14	(A) (B) (C) (D)	29	(A) (B) (C) (D)	44	(A) (B) (C) (D)	59	(A) (B) (C) (D)
15	(A) (B) (C) (D)	30	(A) (B) (C) (D)	45	(A) (B) (C) (D)	60	(A) (B) (C) (D)