

AB-NK025-4

CDC2-CT (CDK1-1) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase CDK1 (CDC2)



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Target Protein

Name Long:	Cyclin-dependent protein-serine kinase 1; Cell division control protein 2 homologue
Alias:	Cdc2; CDC28; CDC28A; CDC2A; Cell division control protein 2; Cell division cycle 2, G1 to S and G2 to M; Cyclin-dependent kinase 1; P34 protein kinase; Kinase Cdc2; MPF; DKFZp686L20222; MGC111195; ENSG00000170312
UniProt ID:	P06493
Sequence Predicted Mass (KDa):	34.095 (297 AA; P06493); 27.503 (240 AA; P06493-2)
Observed SDS-PAGE Mass (KDa):	30-35

Immunogen

Antibody Immunogen Source:	Human CDK1 (CDC2) sequence peptide
Antibody Immunogen Sequence:	CFLSKMLVYDPAKRISGKMALKHPYFDDLDNQIKKM
Location in Target:	Corresponds to amino acid residues F263 to M297; C-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced on an A.B.I. automated peptide synthesizer. The peptide was cleaved from the synthesis resin with HF and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by amino acid analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 µg
Antibody Concentration:	1 mg/ml
Lot Number:	10320
Storage Buffer:	100 mM Tris-glycine, pH 7.0
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months, but either 0.1% sodium azide or 0.05% Thimerosal should be added. Avoid repeated freeze-thaw cycles.

AB-NK025-4

CDC2-CT (CDK1-1) Antibody



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Cross Reactivities:	In rat tissues it cross reacts with 46 and 70 KDa proteins.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit www.kinexusproducts.ca or contact us at 1-866-KINEXUS(546-3987)