

AB-NK042-1

COT (COT-1) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase COT (MAP3K8, TPL2)



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Target Protein

Name Long:	Osaka thyroid oncogene protein-serine kinase (Tpl2); Mitogen-activated protein kinase kinase kinase 8
Alias:	FLJ10486; M3K8; MAP3K8; Tpl2; TPL2; TPL-2; tumour progression locus 2; COT; ESTF; EST; Tpl-2; c-COT; ENSG00000107968
UniProt ID:	P41279
Sequence Predicted Mass (kDa):	52.925 (467 AA; P41279); 49.620 (438 AA; P41279-2)
Observed SDS-PAGE Mass (kDa):	52-57

Immunogen

Antibody Immunogen Source:	Human COT (MAP3K8, TPL2) sequence peptide
Antibody Immunogen Sequence:	MEYMSTGSDNKKEIDC
Location in Target:	Corresponds to amino acid residues M1 to D15; N-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced on an A.B.I. automated peptide synthesizer. The peptide was cleaved from the synthesis resin with HF and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by amino acid analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 µg
Antibody Concentration:	1 mg/ml
Lot Number:	10320
Storage Buffer:	100 mM Tris-glycine, pH 7.0
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months, but either 0.1% sodium azide or 0.05% Thimerosal should be added. Avoid repeated freeze-thaw cycles.

AB-NK042-1

COT (COT-1) Antibody



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Strong immunoreactivity with recombinant human Cot expressed in E. coli as a glutathione S transferase fusion protein.
Antibody Cross Reactivities:	Cross-reacts with an unidentified 30 KDa protein in rat skeletal muscle.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit www.kinexusproducts.ca or contact us at 1-866-KINEXUS(546-3987)