

Whole lysate from HeLa cells that were cultivated to 70-90% confluency and deprived of serum for 18-20 hours prior to harvesting



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Production	
Product Name Long:	Human epidermoid carcinoma HeLa cells - Whole lysate
Production Method:	HeLa cells were cultivated to 70-90% confluency and deprived of serum for 18-20 hours prior to harvesting. Lysates were prepared from scrapped cells that were homogenized by sonication in buffer formulated with 1% Triton X-100, 60 mM β -glycerophosphate, pH 7.2, 20 mM MOPS, 20 mM sodium pyrophosphate, 30 mM sodium fluoride, 5 mM EDTA, 3 mM benzamidine, 2 mM EGTA, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonylfluoride, 1 mM dithiothreitol, 10 μ M leupeptin, and 5 μ M pepstatin. Lysates were prepared following sonication and 30 min ultracentrifugation at 100,000 rpm. Lysates were further diluted in SDS-PAGE sample buffer at a final concentration of 2 mg/ml.
Amount:	200 µg
Protein Concentration:	2 mg/ml
Storage Stability:	1 year at -70°C
Applications	

Lysate Use Description: For testing antibodies by immunoblotting.

This product is for in vitro research use only and is not intended for use in humans or animals.