

47 W Polk St. STE 100-241 Chicago, IL 60605 liftmode@liftmode.com

Lift \\ode

20-06-2022

Icariin

(Epimedium P.E. 98%)

Material Lot #: 171010 Test Date:

Country of Origin: China Re-Test Date: 17-06-2025

Analysis Claim Result

Icariin ≥98% 96.9%*

Te	st Specification	Result
Icariin (HPLO	∑) ≥98%	%96.9
Lead Mercury Cadmium Arsenic	≤0.5ppm ≤0.5ppm ≤0.5ppm ≤0.5 ppm	0.053 ppm <0.001 ppm 0.090 ppm 0.101 ppm
Total Aerobic Plate Coun Yeast & Moulds Moulds Escherichia coli Coliforms Salmonella	<100 cfu/g <100 cfu/g Negative <10 cfu/g	<100 cfu/g <100 cfu/g <10 cfu/g Negative <10 cfu/g Negative

Icariin should be stored at or below room temperature in a tightly sealed durable container. Icariin should be protected from excess heat, direct sunlight, excess humidity, and moisture. Icariin has a retesting period of 3 years from the date of analysis when properly stored.

*Icariin HPLC results are within acceptable +/-2% margin of error.



Main Benefits

Icariin is a phyochemical found in Horny Goat Weed, is primarily grown in China. Also known as Yin Yang Huo, this plant has a long history of use for sexual health.

Benefits include supporting a healthy libido, and offer support for bones and cartilage, and supporting a healthy metabolism.

Overall, Horny Goat Weed extract can be recommended for both men and women who are looking for a powerful aphrodisiac supplement.

Main Cautions

- Side effects have been researched thoroughly and it has been found that moderate use of Horny Goat extracts very rarely results in negative side effects.
- At lager servings, possible side effects may include dizziness, dry mouth, thirstiness and nausea.
- Icariin may be unsafe for pregnant women and people with bleeding disorders, low blood pressure or hormone sensitive cancers.

Usage Tips

A 0.15 cc measuring scoop is included. One rounded scoop contains approximately one serving, or approximately **50mg of Icariin**. As a dietary supplement, take 1-4 servings 1-2 times per day. Start at the lower suggested quantity to assess response.

The negative effects of Icariin are dependent on the amount taken. Use of a scale with 10mg/0.01g accuracy or better is highly recommended.

Given the potency, Icariin can be very bitter and astringent. Use of capsules, or mixing with tea, yogurt, apple sauce, or oatmeal may help make the powder easier to tolerate.

- This supplement is not intended to treat, diagnose, prevent, or cure any diseases. Consult your healthcare provider before use if you have a medical condition or if you are taking any prescription medications.
- It is safe to stack Icariin with other supplements, so long as the amount consumed does not exceed the suggested serving size.
- The benefits of Icariin are most effective when they are supported by a healthy diet and plenty of exercise



24 June 2022

Job Number:	26026a
PO Number:	verba

Synaptent LLC 47 W. Polk Street #100-241 Chicago, Illinois 60605

REPORT OF ANALYSIS

One small jar labeled "Icariin 98% #171010" was received on 16 June 2022. The contents of the jar were analyzed for purity using high pressure liquid chromatography (HPLC). The detector was set to $195 \, \mathrm{nm}$. The results are summarized in the table below.

Sample	Chromatographic Purity (% area)		
Icariin 98% #171010	96.9		

Neil E. Spingam, Ph.D. Lab Director

2021 East Fourth Street, Suite 112

Santa Ana, California 92705

(714) 543-2211



812 Meadow Lark Lane, Godlettsville, TN 37072 Telephone: 615-239-8604

Certificate of Analysis

Synaptent LLC 47 W Polk Street, 100-241 Chicago, IL 60654

Product Name	Icariins	Client Lot Number	171010
Report Date	01/03/18	Laboratory #	9674

Test	Method	Result	
Assay	CA-081 (HPLC)	99.7%	
Lead	ICP-MS USP <730>	0.053 ppm	
Arsenic	ICP-MS USP <730>	0.101 ppm	
Cadmium	ICP-MS USP <730>	0.090 ppm	
Mercury	ICP-MS USP <730>	<0.001 ppm	
Total Aerobic Count	Biolumix	<100 CFU/G	
Yeast and Mold	Biolumix	<100 CFU/G	
Salmonella	Biolumix	Negative	
E. Coli	Biolumix	Negative .	
Coliform	Biolumix	<10 CFU/G	


Collin Thomas MDC
Laboratory Manager

01/03/2018 1/3 /18 Date

The result(s) stated in this report is only for the sample submitted. This report may not be reproduced in whole or in part, nor may any reference be made to the work, the result, or the company in any news release, public announcements or advertising without our prior written consent.

Instrument Annex-1 Sequence 26019, 26025- MK677, RAD140, NPT (Noopepi), loanin Pulities

Chromatogram and Results				
njection Details				
njection Name:	100ppm Icariin lot171010 dup	Run Time (min):	22.00	
Vial Number:	RB7	Injection Volume:	10.00	
njection Type:	Unknown	Channel:	UV_VIS_1	
Calibration Level:		Wavelength:	195	
nstrument Method:	AB 150mm Max RP 22min Method	Bandwidth:	10	
Processing Method:	Processing Method	Dilution Factor:	1.0000	
niection Date/Time:	20/Jun/22 18:14	Sample Weight:	1.0000	



	gration Results			A CONTRACTOR OF THE PARTY OF TH	and the same of th	And the second of	
No.	Peak Name	Retention Time min	Area mAU*min	Height mAU	Relative Area %	Relative Height %	Amount
1	FERRIT	10.077	0.479	8.735	0.84	0.91	n.a.
2		10.180	0.045	0.966	0.08	0.10	n.a.
3		10.333	0.015	0.349	0.03	0.04	n.a.
4		10.430	0.016	0.409	0.03	0.04	n.a.
5		10.567	0.052	0.637	0.09	0.07	n.a.
6	Icanin	10.790	55.117	936.798	96.92	97.09	n.a.
1	1200	11.070	0.069	1.327	0.12	0.14	n.a.
В		11.407	0.032	0.603	0.06	0.06	n.a.
9		11.527	0.012	0.285	0.02	0.03	п.а.
10		12.300	0.975	13.897	1.71	1.44	n.a.
11_		12.710	0.055	0.918	0.10	0.10	n.a.
Total	l:		56.868	964.925	100.00	100.00	

Report Template/Integration

Chromeleon (c) Dionex Version 7.2.10.23925