

Solutions

Chapter 1 Exponents and Scientific Notation

Basic Practice

1. Evaluate the following without using a calculator.

Solution

(a) $(4^5 + 7^9)^0 = 1$

(b) $6^8 \times 6^7 \div 6^{13} = 6^{8+7-13}$
 $= 6^{15-13}$
 $= 6^2$
 $= 36$

(c) $2^{15} \times 2^5 \div (4)^8 = 2^{15+5} \div (2^2)^8$
 $= 2^{20} \div 2^{16}$
 $= 2^{20-16}$
 $= 2^4$
 $= 16$

(d) $(3^2)^2 = 3^4$
 $= 81$

(e) $(5^0 + 5^1) \times 5^2 = (1 + 5) \times 25$
 $= 6 \times 25$
 $= 150$

(f) $(2^3)^2 + (2^3 \times 2^2) = [2^6 + (2^3 \times 2^2)]$
 $= 64 + (8 \times 4)$
 $= 64 + 32$
 $= 96$

(g) $9^{\frac{1}{2}} + 9^2 + 9^{-1} = 3 + 81 + \frac{1}{9}$
 $= 84\frac{1}{9}$

(h) $27^{\frac{2}{3}} = (3^3)^{\frac{2}{3}}$
 $= 3^2$
 $= 9$

2. Evaluate the following without using a calculator.

Solution

(a) $16^{10} \times 16^{-8} \div \sqrt{16} = 16^2 \div 16^{\frac{1}{2}}$
 $= 16^{\frac{3}{2}}$
 $= (4^2)^{\frac{3}{2}}$
 $= 4^3$
 $= 64$

(b) $5^{\frac{5}{2}} \times 5^2 \div 5^{\frac{3}{2}} = 5^{\frac{5}{2}+2-\frac{3}{2}}$
 $= 5^3$
 $= 125$

(c) $25^{\frac{3}{2}} = (5^2)^{\frac{3}{2}}$
 $= 5^3$
 $= 125$

(d) $(\sqrt{64})^{\frac{2}{3}} = 8^{\frac{2}{3}}$
 $= (2^3)^{\frac{2}{3}}$
 $= 2^2$
 $= 4$

(e) $2^{-1} - 2^{-2} + 2^{-3} = \frac{1}{2^1} - \frac{1}{2^2} + \frac{1}{2^3}$
 $= \frac{1}{2} - \frac{1}{4} + \frac{1}{8}$
 $= \frac{3}{8}$

(f) $\left(\frac{2}{3}\right)^{-3} = \left(\frac{3}{2}\right)^3$
 $= \frac{27}{8}$
 $= 3\frac{3}{8}$

(g) $(-3)^2 + 3^{-2} = 9 + \frac{1}{9}$
 $= 9\frac{1}{9}$

(h) $\left(8^{\frac{2}{3}}\right)^{-2} = \left[(2^3)^{\frac{2}{3}}\right]^{-2}$
 $= (2^2)^{-2}$
 $= 4^{-2}$
 $= \frac{1}{16}$

3. Simplify the following and express your answers with positive exponents.

Solution

(a) $(3x^4)^2 = 9x^8$

(b) $5x^3 \times 3x^2 = (5 \times 3) \times (x^3 \times x^2)$
 $= 15x^5$

(c) $24y^3 \div 8y^2 = \frac{24}{8}y^{3-2}$
 $= 3y$

(d) $(3x^5)^0 = 1$

(e) $\left(a^{\frac{2}{3}}\right)^6 = a^4$

(f) $\left(\frac{z^8}{z^2}\right)^{\frac{1}{3}} = (z^6)^{\frac{1}{3}}$
 $= z^2$

(g) $w^{-4} = \frac{1}{w^4}$

(h) $(c^4)^{-2} = c^{-8}$
 $= \frac{1}{c^8}$

4. Simplify the following and express your answers with positive exponents.

Solution

$$(a) \quad 15x^3y \div 3xy^4 = \frac{5x^3y}{xy^4}$$

$$= \frac{5x^2}{y^3}$$

$$(b) \quad (p^2)^{-1} \times (q^3)^2 = p^{-2} \times q^6$$

$$= \frac{q^6}{p^2}$$

$$(c) \quad \sqrt[3]{27s^6t^9} = 3s^2t^3$$

$$(d) \quad (3x^3y^2)^2 \times x^2y^4 = 9x^6y^4 \times x^2y^4$$

$$= 9x^8y^8$$

$$(e) \quad (2a^4b^{-3})^2(a^{-1}b)^5 = 4a^8b^{-6}a^{-5}b^5$$

$$= \frac{4a^3}{b}$$

$$(f) \quad \frac{(n^2)^2}{m^6 \times n^7} = \frac{n^4}{m^6n^7}$$

$$= \frac{1}{m^6n^3}$$

$$(g) \quad \frac{6p^4 \times 7q^3}{14q^6 \times 3p^2} = \frac{2p^4q^3}{2q^6p^2}$$

$$= \frac{p^2}{q^3}$$

$$(h) \quad \left(\frac{x^4}{9y^6}\right)^{\frac{1}{2}} = \frac{x^2}{3y^3}$$

5. Solve the following equations.

Solution

$$(a) \quad 6^x = 1$$

$$6^x = 6^0$$

$$x = 0$$

$$(b) \quad 3^x = 27$$

$$3^x = 3^3$$

$$x = 3$$

$$(c) \quad 2^x = \frac{1}{16}$$

$$2^x = \frac{1}{2^4}$$

$$2^x = 2^{-4}$$

$$x = -4$$

$$(d) \quad 4^x = 2^{15}$$

$$2^{2x} = 2^{15}$$

$$2x = 15$$

$$x = 7\frac{1}{2}$$

$$(e) \quad 5^x = 25^{-8}$$

$$5^x = (5^2)^{-8}$$

$$5^x = 5^{-16}$$

$$x = -16$$

$$(f) \quad \sqrt{7^x} = 49$$

$$7^{\frac{x}{2}} = 7^2$$

$$\frac{x}{2} = 2$$

$$x = 4$$

6. Solve the following equations.

Solution

$$(a) \quad 3^{4x} = 9^{12}$$

$$3^{4x} = (3^2)^{12}$$

$$3^{4x} = 3^{24}$$

$$4x = 24$$

$$x = 6$$

$$(b) \quad 6^{2-x} = 36^4$$

$$6^{2-x} = (6^2)^4$$

$$6^{2-x} = 6^8$$

$$2 - x = 8$$

$$x = -6$$

$$(c) \quad 5^2 \times 5^{2x} = 5^2$$

$$5^{2+2x} = 5^2$$

$$2 + 2x = 2$$

$$2x = 0$$

$$x = 0$$

$$(d) \quad 2^x \div 32 = 2^{-x}$$

$$2^x \div 2^5 = 2^{-x}$$

$$2^{x-5} = 2^{-x}$$

$$x - 5 = -x$$

$$2x = 5$$

$$x = 2\frac{1}{2}$$

$$(e) \quad \sqrt[3]{7^2} = 7^6$$

$$7^{\frac{2}{3}} = 7^6$$

$$\frac{2}{3} = 6$$

$$x = \frac{1}{3}$$

$$(f) \quad 4^x - 1 = 0$$

$$4^x = 1$$

$$4^x = 4^0$$

$$x = 0$$

7. Express each of the following in scientific notation correct to 3 significant figures.

Solution

$$(a) \quad 3,245 = 3.245 \times 10^3$$

$$= 3.25 \times 10^3 \text{ (correct to 3 sig. fig.)}$$

$$(b) \quad 6,782,450 = 6.78245 \times 10^6$$

$$= 6.78 \times 10^6 \text{ (correct to 3 sig. fig.)}$$

$$(c) \quad 0.03463 \times 10^7 = 3.463 \times 10^{-2} \times 10^7$$

$$= 3.46 \times 10^5 \text{ (correct to 3 sig. fig.)}$$

$$(d) \quad 279,825 \div 10^2 = 2,798.25$$

$$= 2.79825 \times 10^3$$

$$= 2.80 \times 10^3 \text{ (correct to 3 sig. fig.)}$$

$$(e) \quad 0.006752 = 6.752 \times 10^{-3}$$

$$= 6.75 \times 10^{-3} \text{ (correct to 3 sig. fig.)}$$

$$(f) \quad 0.0000464 = 4.64 \times 10^{-5}$$

$$(g) \quad 0.03463 \times 10^{-5} = 3.463 \times 10^{-2} \times 10^{-5}$$

$$= 3.46 \times 10^{-7} \text{ (correct to 3 sig. fig.)}$$

$$(h) \quad 4,295 \div 10^{-8} = 4.295 \times 10^3 \div 10^{-8}$$

$$= 4.295 \times 10^{11}$$

$$= 4.30 \times 10^{11} \text{ (correct to 3 sig. fig.)}$$

22. (a) Simplify by factorization,
 (i) $(x + 1)^2 - (x - 1)^2$,
 (ii) $(x + 2)^2 - (x - 2)^2$,
 (iii) $(x + 3)^2 - (x - 3)^2$.
 (b) Hence, simplify $(x + n)^2 - (x - n)^2$.
 (c) Use the answer in (b) to
 (i) evaluate $(345 + 29)^2 - (345 - 29)^2$,
 (ii) solve the equation $(x + 100)^2 - (x - 100)^2 = 640$.

Solution

(a) (i) $(x + 1)^2 - (x - 1)^2$
 $= [x + 1 + x - 1][x + 1 - (x - 1)]$
 $= 2x(2)$
 $= 4x$
 (ii) $(x + 2)^2 - (x - 2)^2$
 $= (x + 2 + x - 2)[x + 2 - (x - 2)]$
 $= 2x(4)$
 $= 8x$
 (iii) $(x + 3)^2 - (x - 3)^2$
 $= (x + 3 + x - 3)[x + 3 - (x - 3)]$
 $= 2x(6)$
 $= 12x$

(b) By inspection of the answers in (a),
 $(x + n)^2 - (x - n)^2 = 2x(2n)$
 $= 4nx$

(c) (i) $(345 + 29)^2 - (345 - 29)^2$
 $= 4(29)(345)$
 $= 40,020$
 (ii) $(x + 100)^2 - (x - 100)^2 = 400$
 $\therefore 4(100)x = 640$
 $400x = 640$
 $x = 1.6$

Challenging Practice

23. (a) The diameter of a circle is $(6r + 16s)$ cm, where r and s are positive numbers.
 Find and simplify, in terms of r , s , and π ,
 (i) the circumference of the circle,
 (ii) the area of the circle.
 (b) Suppose that the circle is the base of a solid prism and the height of the prism is twice the diameter of its base.
 Find and simplify, in terms of r , s , and π ,
 (i) the volume of the prism,
 (ii) the total surface area of the prism.

Solution

(a) (i) Diameter $= (6r + 16s)$ cm
 Circumference $= (6r + 16s)\pi$ cm
 (ii) Radius $= (3r + 8s)$ cm
 \therefore area $= \pi(3r + 8s)^2$
 $= (9r^2 + 48rs + 64s^2)\pi$ cm²
 (b) (i) Volume of prism
 $=$ base area \times height
 $= \pi(9r^2 + 48rs + 64s^2) \times 2(6r + 16s)$
 (ii) Total surface area of prism
 $= 2 \times$ base area $+ \text{circumference} \times \text{height}$
 $= 2(9r^2 + 48rs + 64s^2)\pi + (6r + 16s)\pi \times 2(6r + 16s)$
 $= (18r^2 + 96rs + 128s^2)\pi + 2\pi(6r + 16s)^2$
 $= [18r^2 + 96rs + 128s^2 + 2(36r^2 + 192rs + 256s^2)]\pi$
 $= (90r^2 + 480rs + 640s^2)\pi$ cm²

24. (a) Evaluate $(997 - w)^2 + (993 - w)^2$ if $(997 - w)(993 - w) = 21$.
 (b) Find the value of $(1 + x)(2 + x)(3 + x)^2(4 + x)(5 + x)$ if $x^2 + 6x = 2$.

Solution


(a) $(997 - w)^2 + (993 - w)^2$
 $= [(997 - w)^2 - 2(997 - w)(993 - w) + (993 - w)^2] + 2(997 - w)(993 - w)$
 $= [(997 - w) - (993 - w)]^2 + 2(21)$
 $= (997 - 993)^2 + 42$
 $= 16 + 42$
 $= 58$
 (b) $(1 + x)(2 + x)(3 + x)^2(4 + x)(5 + x)$
 $= (1 + x)(5 + x)(2 + x)(4 + x)(3 + x)^2$
 $= (x^2 + 6x + 5)(x^2 + 6x + 8)(x^2 + 6x + 9)$ ----- (1)
 Since $x^2 + 6x - 2 = 0$
 $x^2 + 6x = 2$ ----- (2)
 Putting (2) into (1), $(2 + 5)(2 + 8)(2 + 9) = 770$
 $\therefore (1 + x)(2 + x)(3 + x)^2(4 + x)(5 + x) = 770$

25. A stone is tossed from a point W into the air. The height y meters above ground level of the stone at time t seconds is given by $y = -t^2 + 4t + 5$.
 (a) How far above the ground level is W ?
 (b) (i) If $-t^2 + 4t + 5$ can be expressed as $-(t - 2)^2 + P$, find the value of P .
 (ii) Hence, deduce the maximum vertical distance of the stone above point W and the time taken to reach this distance.

Solution

(a) When $t = 0$,
 $y = -(0)^2 + 4(0) + 5$
 $= 5$
 $\therefore W$ is 5 m above the ground level.
 (b) (i) Method 1
 $-(t - 2)^2 + P = -(t^2 - 4t + 4) + P$
 $= -t^2 + 4t - 4 + P$
 Given $-t^2 + 4t - 4 + P = -t^2 + 4t + 5$
 $-4 + P = 5$
 $\therefore P = 9$
Method 2
 Given $-t^2 + 4t + 5 = -(t - 2)^2 + P$
 $-t^2 + 4t + 5 = -(t^2 - 4t + 4) + P$
 $-t^2 + 4t + 5 = -t^2 + 4t - 4 + P$
 $\therefore P = 9$
 (ii) $y = -t^2 + 4t + 5$
 $= -(t - 2)^2 + 9$
 $(t - 2)^2 \geq 0$
 $\therefore -(t - 2)^2 \leq 0$
 $\therefore y$ is maximized when $t = 2$.
 maximum value of $y = 9$
 \therefore maximum vertical distance of the stone above point $W = 9 - 5$
 $= 4$ m
 Time taken to reach this distance $= 2$ s

33. In the diagram, the right-angled triangle DEF is an enlargement of the triangle ABC about the centre P which is not shown. B and D are points on the line AE , $AB = 2$ units, $BC = 1$ unit, $BD = 4$ units, and $DE = 6$ units.




- State the scale factor of the enlargement.
- Find the length of EF .
- By calculation or accurate scale drawing on a sheet of graph paper, find the distance of A from the centre of enlargement P .

Solution

(a) Scale factor = $\frac{DE}{AB}$
 $= \frac{6}{2}$
 $= 3$

(b) $\frac{EF}{BC} = \frac{DE}{AB}$
 $= 3$


$\therefore \frac{EF}{1} = 3$
 $EF = 3$ units




From the diagram, the distance of A from the center of enlargement P is 3 units.

Enrichment


34. The diagram shows a figure and a point X drawn on the grid lines. Divide the figure into 4 congruent figures along the grid lines only. The four congruent figures must not all meet at the point X .



Solution



35. Two squares, $ABCD$ and $PQRS$, are drawn in the diagram.



Describe completely the transformation that moves

- $ABCD$ to $PQRS$,
- $ABCD$ to $QPSR$,
- $ABCD$ to $QRSP$,
- $ABCD$ to $RSPQ$.

Solution

- $ABCD$ is translated to $PQRS$ by 15 units down.
- $ABCD$ is reflected in the line $y = 15$ to $QPSR$.
- $ABCD$ is rotated through 90° clockwise about the point $(5, 15)$ to $QRSP$.
- $ABCD$ is rotated 180° about the point $(12.5, 15)$ to $RSPQ$.