


Orego-Stim Liquid



Composition

Orego-Stim Liquid is a high quality eubiotic containing a 100% natural oregano essential oil.

Animal focus

Designed for the use in all livestock and specific companion animals.

Main benefits

- ✓ Non-synthetic, consistent single source of natural oregano oil
- ✓ Flavouring compound, shown to increase voluntary intakes
- ✓ Helps maintain a healthy appetite through the critical stages of growth such as weaning
- ✓ Oregano essential oil contains anti-oxidants that help neutralise free radicals. Free radicals can cause cell damage and levels of these may be increased in times of stress.
- ✓ Oregano essential oil helps support animals natural defences
- ✓ Oregano essential oil can help maintain a state of eubiosis. Eubiosis is defined as well balanced and diverse microbiota.

Directions for use

Orego-Stim Liquid inclusions vary according to conditions. Please see reverse for details.

For specific requirements, please contact your local representative.



Typical specifications

Physical analysis	Milky off white liquid with characteristic smell
Packaging	Available in 500ml, 1L, 5L, 25L, 200L, 1000L
Shelf life	36 months from date of manufacture
Storage	Store in cool, dry area away from heat and damp conditions
Precautionary classification	Irritant
Accreditation status	All products are FEMAS approved



Directions for use

Orego-Stim Liquid inclusions vary according to conditions:

Poultry, Pigeons & Avian

For general use: 150 ml per 1000 L of water (1.5 ml / 10 L of water)
During critical periods: 300 ml per 1000 L of water (3.0 ml / 10 L of water)

Sheep & Goats

For first 28 days: 500 ml per 1000 L of water (5.0 ml / 10 L of water)
Maintenance rate: 125 ml per 1000 L of water (1.25 ml / 10 L of water)

Swine

Creep feeding up to 15 kg bodyweight: 500 ml per 1000 L of water (5.0 ml / 10 L of water)
15 kg to 25 kg bodyweight: 250 ml per 1000 L of water (2.5 ml / 10 L of water)
25 kg bodyweight to slaughter: 125 ml per 1000 L of water (1.25 ml / 10 L of water)
Sows & Boars: 250 ml per 1000 L of water (2.5 ml / 10 L of water)

Calves, Beef & Dairy Cattle

Calves (birth to weaning): An initial 10 ml/head/day added to milk for first 10 days followed by 2 ml/head/day until weaning
Beef cattle: An initial 500 ml per 1000 L of water (5.0 ml / 10 L of water) for 28 days followed by a maintenance rate of 125 ml per 1000 L of water (1.25 ml / 10 L of water)
Dairy cattle: 15 ml/head/day for 28 days

Rabbits

For routine use: 500 ml per 1000 L of water (5.0 ml / 10 L of water)